

NASA TECHNICAL NOTE



NASA TN D-2874

NASA TN D-2874

FACILITY FORM 602

N65-27346 (ACCESSION NUMBER)	(THRU)
15 (PAGES)	1 (CODE)
(NASA CR OR TMX OR AD NUMBER)	29 (CATEGORY)

GPO PRICE \$ _____
OTS PRICE(S) \$ _____
Hard copy (HC) 1.00
Microfiche (MF) .50

A COMPUTER PROGRAM TO CALCULATE ARTIFICIAL RADIATION BELT DECAY FACTORS

by E. G. Stassinopoulos
Goddard Space Flight Center
Greenbelt, Md.

A COMPUTER PROGRAM TO CALCULATE ARTIFICIAL
RADIATION BELT DECAY FACTORS

By E. G. Stassinopoulos

Goddard Space Flight Center
Greenbelt, Md.

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

For sale by the Clearinghouse for Federal Scientific and Technical Information
Springfield, Virginia 22151 - Price \$1.00

A COMPUTER PROGRAM TO CALCULATE ARTIFICIAL RADIATION BELT DECAY FACTORS

by

E. G. Stassinopoulos

Goddard Space Flight Center

SUMMARY

The computer code presented here is a FORTRAN II, Monitor-Controlled program for the IBM 7090-7094, and has been developed for calculating artificial radiation-belt decay factors to evaluate the time-dependent change in the intensity of the Starfish electrons injected into the upper atmosphere on July 9, 1962. The program (called subroutine DECAY) has been written in subroutine form as part of an orbital flux code for calculating omnidirectional, vehicle-encountered fluxes. Execution time for the program varies, but even in the worst case it is still insignificantly small (a fraction of a second). By means of this code, anticipated fluxes at any epoch may be predicted from known radiation levels at a given time.

CONTENTS

Summary	iii
INTRODUCTION	1
METHOD OF COMPUTATION	2
USAGE AND LIMITATIONS.	3
CONCLUSION	4
References	4
Appendix A—Subroutine.	5
Appendix B—List of Symbols	11

A COMPUTER PROGRAM TO CALCULATE ARTIFICIAL RADIATION BELT DECAY FACTORS

by
E. G. Stassinopoulos
Goddard Space Flight Center

27346

INTRODUCTION

A significant loss of Starfish electrons has occurred since their injection into the upper atmosphere on July 9, 1962. It is assumed that this artificial radiation-belt population decreases exponentially with time (References 1 and 2), thus, the remaining particle intensity may be adequately approximated by a decay factor of the form,

$$\exp\left(-\frac{\Delta t}{\tau}\right), \quad (1)$$

where τ is the mean lifetime of electrons with energies greater than 1.2 Mev (Reference 3) and Δt is the specified time difference ($t_2 - t_1$). If the flux at time t_2 is then related to that of time t_1 by this decay factor, the expression,

$$\Phi(t_2) = \Phi(t_1) \exp\left(-\frac{(t_2 - t_1)}{\tau}\right), \quad (2)$$

is obtained which may be used readily in a computer calculation.

Measurements of mean lifetime, τ , (by Van Allen and Lin with INJUN III and by Bostrom and Williams of APL with 1963-38c, References 1 and 2) have shown, that the decay constant is mainly a function of the magnetic shell parameter, L , and is almost independent of the field strength, B .

It should be noted, that the 1963-38c measurements were taken at a later epoch than the INJUN III and hence are consistently lower (Table 1 and Figure 1).

Table 1

Subroutine-DECAY Mean Lifetime Measurements of Artificial Radiation-Belt Electrons.

Measurement No.	Magnetic-Shell Parameter, L (earth radii)	Magnetic Field Strength, B (gauss)	Mean Lifetime, τ_1^\dagger (days)	Mean Lifetime, τ_2^\ddagger (days)
1	(1.15)	-----	(0)	(0)
2	1.20	0.185-0.205	130	120
3	1.23	0.170-0.205	190	165
4	1.30	0.100-0.230	320	235
5	1.40	0.165-0.210	500	390
6	1.45*	-----	565*	445*
7	1.50	0.175-0.215	590	460
8	1.55*	-----	585*	445*
9	1.60	0.180-0.225	(500)	360
10	1.65*	-----	275*	225*
11	(1.70)	-----	(100)	(100)
12	1.725*	-----	50*	65*
13	1.75*	-----	25*	40*
14	(1.80)	-----	(0)	(0)

All values in parentheses are estimates not contained in original information.

*Values indicated were obtained by fitting the τ -curve (Figure 1) to the source and to the estimates.

$\dagger\tau_1$ measurements made by Van Allen and Lin with INJUN III.

$\ddagger\tau_2$ measurements made by Bostrom and Williams with 1963-38c.

METHOD OF COMPUTATION

The subroutine first calculates the time difference $\Delta t = t_2 - t_1$, where t_1 , contained internally, defines the epoch of the electron grid used* and t_2 , an input quantity, pertains to a given later date, for which flux information is being sought.

If $\Delta t \leq 0$, the subroutine will bypass the rest of the program and, since no decay is involved in that case, it will set the decay factor equal to unity and return to the main routine.

If $\Delta t > 0$ the subroutine continues, it then initiates computation of the decay factor by




Figure 1—Function $\tau(L)$ fitted to measurements by Van Allen and Lin (τ_1) Bostrom and Williams (τ_2) for $L \leq 1.6$ and to estimates by Hess for $L \geq 1.6$.

*Presently the code is programmed for the E8 grid of November, 1962.

establishing once, in its first passage, the τ -to-L dependence in the form of a look-up table where $\tau(L) = 0$ for $L > 1.8$ (see Table 1). Subsequently, it tests the L of the orbital position under consideration and if $1.15 < L < 1.8$ it proceeds to obtain τ by a linear interpolation from both sources available (i.e., Van Allen and Lin, and Bostrom and Williams) and then either computes or defines the proper decay factor, $\exp(-\Delta t/\tau)$.

Finally, the subroutine sets up suitable control parameters in order to avoid, during its repetitive use by the Orbital Flux Code, recomputation of any quantity that is both invariant with respect to a given trajectory and permanently established for that orbit during the first passage. In this manner the subroutine also passes over the once initialized look-up table.

USAGE AND LIMITATIONS

No actual limitations exist with respect to computation of a decay factor by the subroutine. It is programmed to accept any epoch t_2 including those that fall on, or before, its critical or effective date t_1 .

Epoch t_2 must be given in the form of year, month and day, and in that order. The subroutine will test each quantity separately, in the same order, and convert into days if necessary.

There is no difficulty in substituting another effective date t_1 (for different flux data) into the code as long as appropriate reference relations are included.

Similarly, the mean lifetimes may be changed easily without major modifications by simply inserting new tables of τ to L values between statements 551 and 550 and by adjusting the limits of statements 550 and Dimension (if necessary).

Due to the IBM 7094 floating point limitations, it is to be expected that the decay factors will be zero whenever the ratio $\Delta t/\tau$ becomes greater than that value, which will make the characteristic exceed the storage capacity of the register. When this happens, underflow occurs. Figure 2 shows the relationship of time difference, Δt , to mean lifetime, τ , where the shaded area indicates combinations causing such underflow. This however does not affect the results in any way. Contributions of such small magnitudes can be neglected. A sample computation with the τ_1 decay constant for various constant flux ratios (ϕ/ϕ_0) is shown in Figure 3.




Figure 2—Underflow relationship of mean lifetime τ and time difference Δt . (Shaded area shows combinations causing decay factors to become zero.)

The constant τ_1 is that measured by Van Allen and Lin with INJUN III.

CONCLUSION

Whenever the orbital flux code is employed and its output is utilized, caution must be exercised in regard to the following shortcomings and restrictions:

1. The subroutine applies to Starfish electrons only. It does not consider or include natural electrons.*
2. The mean lifetime (and hence the flux) has been set equal to zero for $L \geq 1.8$.
3. The decay constants τ_1 and τ_2 were obtained from measurement of $E > 1.2$ -MeV electrons. If used for all energy electrons, the resulting spectra may not be very accurate (values possibly inflated for low energy fluxes).*
4. Since increasing solar activity in the years ahead will greatly affect the lifetimes of the Starfish electrons, the use of this routine for epochs later than 1966 is not advisable.*




Figure 3—Contours of constant flux ratio Φ/Φ_0 as functions of L and t for Starfish electrons. (Each curve traces a given fixed percentage by which an initial flux is requested to decay.)

REFERENCES

1. Van Allen, J. A., "Lifetimes of Geometrically Trapped Electrons of Several MeV Energy," *Nature*, vol. 203, no. 4949, pp. 1006-1007, London: Macmillan and Company, Ltd., September 1964.
2. Bostrom, C. O., Williams, D. J., "Time Decay of the Artificial Radiation Belt," *Journal of Geophysical Research*, vol. 70, no. 1, pp. 240-242, Richmond: The American Geophysical Union, January 1965.

*Private communication with Dr. Wilmot N. Hess, GSFC, "Circular to users of E8 grid," October 20, 1964.

Appendix A

Subroutine

Arguments and Parameters, Flow
Chart and Listing

1. Arguments and Parameters

A. Input:

- IN - Control parameter; governs the calculation of Δt ; may have values 0, 1 or 2; at the start it is set equal to zero in main routine and remains unaffected by it thereafter; it is redefined in subroutine; it performs as follows:
- 0: attempts to compute Δt ,
 - 1: Δt is well defined, hence it bypasses section computing Δt and proceeds to obtain decay factors,
 - 2: Δt cannot be defined, hence it bypasses section computing Δt , sets decay factors equal to one.
- DL - Magnetic shell parameter L associated with orbital position under consideration.
- YEAR - Year of date for which decay factor is to be calculated.
- AMONTH - Month of date for which decay factor is to be calculated.
- DAY - Day of date for which decay factor is to be calculated.
- NOT - Control parameter; governs the initialization of the τ -L tables; may be -1, 0, +1; at the start it is set equal to zero in main routine and remains unaffected by it thereafter; it is redefined in subroutine; it performs as follows:
- 1: skips initialization of tables
 - 0, 1: initializes tables

B. Internal:

- T1 - year(s) in days
- T2 - month(s) in days
- T3 - day(s)
- } These arguments establish the time difference $\Delta t = (t_2 - t_1)$
- T - time difference Δt in days, i.e., the sum of T1, T2, T3
- TAL(I) - L_i values of the τ -L tables, where $i = 1, \dots, 14$
- TA1(I) - τ_{1_i} values of the τ -L tables, where $i = 1, \dots, 14$
- TA2(I) - τ_{2_i} values of the τ -L tables, where $i = 1, \dots, 14$
- KK - Control parameter within L-testing loop; defines special cases.
- TAU1 - decay constant τ_1
- TAU2 - decay constant τ_2


C. Output:

DFCTR1 - decay factor with respect to τ_1

DFCTR2 - decay factor with respect to τ_2

Note: All arguments are in floating point notation except the parameters IN, NOT, KK which are fixed point quantities.

2. Flow Chart



3. Listing

```
      SUBROUTINE DECAY(IN,DL,YEAR,AMONTH,DAY,DFCTR1,DFCTR2,NOT)
C      THE MEAN LIFE TIME TAU1 IS OBTAINED FROM MEASUREMENTS MADE WITH
C      INJUN III
C      THE MEAN LIFE TIME TAU2 IS OBTAINED FROM MEASUREMENTS MADE WITH
C      1963 38C
      DIMENSION TAL(20),TA1(20),TA2(20)
      IF(IN - 1)315,320,310
315  T1 = 0.
      T2 = 0.
      T3 = 0.
      T = 0.
      IF(YEAR - 1962.)310,100,105
100  T1 = -365.
      GO TO 110
105  T1 = 365.*(YEAR - 1963.)
110  IF(AMONTH - 11.)204,200,202
200  T2 = 364.
      GO TO 210
202  T2 = 394.
      GO TO 210
204  MONTH = AMONTH
      GO TO (90,91,92,93,94,95,96,97,98,99),MONTH
 90  T2 = 60.
      GO TO 210
 91  T2 = 91.
      GO TO 210
 92  T2 = 119.
      GO TO 210
 93  T2 = 150.
      GO TO 210
 94  T2 = 180.
      GO TO 210
 95  T2 = 211.
      GO TO 210
 96  T2 = 241.
      GO TO 210
 97  T2 = 272.
      GO TO 210
 98  T2 = 303.
      GO TO 210
 99  T2 = 333.
210  IF(DAY - 1.)300,300,305
300  T3 = 1.
      GO TO 290
305  T3 = DAY
290  T = T1 + T2 + T3
      IF(T - 484.)295,296,296
296  T = T + 1.
295  IF(T)310,310,320
310  DFCTR1 = 1.
      DFCTR2 = 1.
      IN = 2
      RETURN
320  IF(NOT)550,551,551
551  TAL(1) = 1.15
      TAL(2) = 1.20
      TAL(3) = 1.23
      TAL(4) = 1.30
```

```

TAL(5) = 1.40
TAL(6) = 1.45
TAL(7) = 1.50
TAL(8) = 1.55
TAL(9) = 1.60
TAL(10) = 1.65
TAL(11) = 1.70
TAL(12) = 1.725
TAL(13) = 1.75
TAL(14) = 1.80
TA1(1) = 0.
TA1(2) = 130.
TA1(3) = 190.
TA1(4) = 320.
TA1(5) = 500.
TA1(6) = 565.
TA1(7) = 590.
TA1(8) = 585.
TA1(9) = 500.
TA1(10) = 275.
TA1(11) = 100.
TA1(12) = 50.
TA1(13) = 25.
TA1(14) = 0.
TA2(1) = 0.
TA2(2) = 120.
TA2(3) = 165.
TA2(4) = 235.
TA2(5) = 390.
TA2(6) = 445.
TA2(7) = 460.
TA2(8) = 445.
TA2(9) = 360.
TA2(10) = 225.
TA2(11) = 100.
TA2(12) = 65.
TA2(13) = 40.
TA2(14) = 0.
NOT = -1
550 DO 900 K=1,14
    KK = K
    IF(TAL(K) - DL)900,500,501
900 CONTINUE
    GO TO 502
501 IF(KK-1)502,502,321
502 DFCTR1 = 0.
    DFCTR2 = 0.
    GO TO 370
500 TAU1 = TA1(KK)
    TAU2 = TA2(KK)
    GO TO 322
321 TAU1 = TA1(KK) - ((TA1(KK) - TA1(KK-1)) / (TAL(KK) - TAL(KK-1))) * (TAL(KK) -
IDL)
    TAU2 = TA2(KK) - ((TA2(KK) - TA2(KK-1)) / (TAL(KK) - TAL(KK-1))) * (TAL(KK) -
IDL)
322 DFCTR1 = EXPF(-T/TAU1)
    DFCTR2 = EXPF(-T/TAU2)
370 IN=1
    RETURN
    END

```

Appendix B

List of Symbols

Δt = Time difference in days = $(t_2 - t_1)$

t_1 = November 1, 1962, effective date relating to epoch of electron data

t_2 = Any other date

τ = Mean lifetime of Starfish electrons in days

τ_1 = Mean lifetime by Van Allen and Lin from measurements with INJUN III

τ_2 = Mean lifetime by Bostrom and Williams from measurements with 1963-38c

L = Magnetic shell parameter in earth radii

B = Magnetic field strength in gauss

Φ = Omnidirectional flux in particles per square centimeter per second (particles/cm²-second)