https://ntrs.nasa.gov/search.jsp?R=19660004431 2018-07-25T23:35:41+00:00Z

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.

- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.

- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.

- This document is paginated as submitted by the original source.

- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

Produced by the NASA Center for Aerospace Information (CASI)

AD621864

ORC 65-30
AUGUST 1965

CONSERVATIVE TOLERANCE AND CONFIDENCE LIMITS

by
Richard E. Barlow

CLEARINGHOUSE
FOR FEDERAL SCIENTIFIC AND
TECHNICAL INFORMATION

Hardcopy	Microfiche		
$ 1.00	$ 0.50	24 pp	as

ARCHIVE COPY

DDC
OCT 14 1965
TISIA D

OPERATIONS RESEARCH CENTER

COLLEGE OF ENGINEERING

UNIVERSITY OF CALIFORNIA - BERKELEY

CONSERVATIVE TOLERANCE AND CONFIDENCE LIMITS

by

Richard E. Barlow
University of California, Berkeley

September, 1965 ORC 65-30

This research has been supported by the Office of Naval Research Contract Nonr-3656(18) and the Army Research Office Contract DA-31-124-ARO-D-331 with the University of California and the National Aeronautics and Space Administration under contract with the RAND Corporation.

ABSTRACT

This paper extends the validity of exponential tolerance and confidence limits, under certain restrictions, to the class of distributions with monotone failure rate. In particular, the usual exponential lower tolerance limit is shown to be conservative for the increasing failure rate class of distributions in the range of population coverages and confidence coefficients of practical interest. Conservative confidence limits are also obtained on tail probabilities and moments.

CONSERVATIVE TOLERANCE AND CONFIDENCE LIMITS

1. Introduction

A fundamental problem in statistical reliability theory and life testing is to obtain lower tolerance limits as a function of sample data, say $\underline{X} = (X_1, X_2, \ldots, X_n)$. That is, if X denotes the time to failure of an item with distribution F, then we seek a function $L(\underline{X})$ such that

$$P\{1 - F[L(\underline{X})] \geq 1 - q\} \geq 1 - \alpha .$$

We call $1 - q$ the population coverage for the interval $[L(\underline{X}), \infty]$ and $1 - \alpha$ the confidence coefficient. Another important problem is to obtain a function $M(\underline{X})$ such that

$$P\{1 - F(T) \geq M(\underline{X})\} \geq 1 - \alpha$$

for a specified time $T \geq 0$. Related problems are those of obtaining confidence limits on moments and percentiles.

Early papers in life testing (e.g. Epstein and Sobel (1953)) derived confidence limits assuming an exponential life distribution. Goodman and Madansky (1962) examine various criteria for goodness of tolerance intervals and certain optimum properties of the usual exponential tolerance limits are demonstrated. Recently, a great deal of effort has been devoted to obtaining various confidence limits for the Weibull distribution. Dubey (1962) obtains asymptotic confidence limits on $1 - F(T)$ and the failure rate for the class of Weibull distributions with non-decreasing failure rate. He also studies the properties of various estimators for Weibull parameters

(Dubey (1963)). Johns and Lieberman (1965) present a method for obtaining exact lower confidence limits for $1 - F(T)$ when F is the Weibull distribution with both scale and shape parameters unknown. Unlike Dubey, they do not require that the Weibull distribution in question have a non-decreasing failure rate. These confidence limits are obtained both for the censored and non-censored cases and are asymptotically efficient. Hanson and Koopmans (1964) obtain upper tolerance limits for the class of distributions with increasing hazard rate and lower tolerance limits for the class of distributions with PF_2 density, f (i.e. $\log f(x)$ is concave where finite). However, they do not assume non-negative random variables.

Assuming that the sample data arises from a distribution with monotone failure rate (either non-decreasing or non-increasing and $F(0^-) = 0$) we obtain conservative confidence limits for most reliability parameters of interest. These confidence limits are, in part, derived from the exponential distribution. Since in many cases these are optimum confidence limits when the failure distribution is exponential (Goodman and Madansky (1962)), they are, in this sense, best possible for the class of distributions with monotone failure rate. (See Barlow and Proschan (1965) Chapter 2 and Appendix 2 for a discussion of distributions with monotone failure rate and a test for its validity.) They also have the advantage that they are convenient to compute and are <u>not</u> based on a strong, non-verifiable, parametric assumption. Since these confidence limits are derived in part from the exponential distribution this paper, in a sense, represents a new justification for the use of exponential confidence limits in reliability theory.

2. Summary and Discussion of Results.

Let $X_1 \leq X_2 \leq \dots \leq X_r \leq \dots \leq X_n$ denote an ordered sample from a life

distribution F. We shall only allow the possibility of censorship on the right. Our methods will be used to obtain confidence bounds for more general types of censorship in another paper.

We say that a distribution F is IFR (DFR) if and only if $\ln [1 - F(x)]$ is concave where finite (convex on $[0,\infty]$). If F with density f is IFR (DFR) then the failure rate $\frac{f(t)}{1 - F(t)}$ is non-decreasing (non-increasing) in t. Barlow and Proschan (1964) obtain inequalities for expected values of statistics based on the exponential assumption when in fact the true distribution has a monotone hazard rate.

IFR Results

Let

$$\hat{\theta}_{r,n} = \frac{\sum_1^r X_i + (n-r) X_r}{r}$$

and

$$C_{1-\alpha,q}(r) = \begin{cases} \dfrac{-2r \ln(1-q)}{\chi^2_{1-\alpha}(2r)} & \text{if } \chi^2_{1-\alpha}(2r) > -2n \ln(1-q) \\[2ex] \dfrac{r}{n} & \text{if } \chi^2_{1-\alpha}(2r) \leq -2n \ln(1-q) \end{cases}$$

where $\chi^2_{1-\alpha}(2r)$ is the $(1-\alpha)$-th percentage point of a chi-square distribution with $2r$ degrees of freedom.

THEOREM 1. If F is IFR, $F(0^-) = 0$, $\zeta_q = \sup \{ x \;\; F(x) \leq q \}$ then

(1) $P \{ 1 - F[C_{1-\alpha,q}(r) \hat{\theta}_{r,n}] \geq 1 - q \} \geq 1 - \alpha$

(2) $P \{ \zeta_q \geq C_{1-\alpha,q}(r) \hat{\theta}_{r,n} \} \geq 1 - \alpha$.

Mathematically (1) and (2) are equivalent statements. When $\chi^2_{1-\alpha}(2r) > -2n \ln(1 - q)$, the lower tolerance limit provided by (1) is identical with the exponential tolerance limit. Amazingly enough, the exponential lower tolerance limits provide conservative tolerance limits for most cases of practical interest. For example, if $1 - \alpha > 1 - e^{-1} \sim .633$ and $1 - q > e^{-r/n}$, then the inequality $\chi^2_{1-\alpha}(2r) > -2n \ln(1 - q)$ holds. In the sense of being "most stable" (see Goodman and Madansky)1962)) this is the best lower tolerance limit for the exponential distribution and hence a "sharp" conservative tolerance limit. If the full sample is known this is "best" for $r = n$.

Let

$$C^*_{\alpha,q}(r) = \begin{cases} \dfrac{-2r \ln(1 - q)}{\chi^2_\alpha(2r)} & \text{if } \chi^2_\alpha(2r) < -2 \ln(1-q) \\ r & \text{if } \chi^2_\alpha(2r) \geq -2 \ln(1-q). \end{cases}$$

<u>THEOREM</u> 2. If F is IFR, $F(0^-) = 0$, $\zeta_q = \sup \{ x \mid F(x) \leq q \}$ then

(3) $P \{ F[C^*_{\alpha,q}(r) \hat{\theta}_{r,n}] \geq q \} \geq 1 - \alpha$

(4) $P \{ \zeta_q \leq C^*_{\alpha,q}(r) \hat{\theta}_{r,n} \} \geq 1 - \alpha$.

In this case the exponential upper tolerance limits are valid when $\chi^2_\alpha(2r) \leq -2 \ln(1 - q)$. Unfortunately this inequality does not hold for all

values of r $(1 \leq r \leq n)$ in the range of population coverage values, q, of greatest practical interest. A table follows which gives the largest values of r such that

$$\chi^2_\alpha(2r) \leq -2 \ln(1 - q) .$$

Table 1

Largest values of r such that the exponential upper tolerance limit is a conservative upper tolerance limit for the IFR class.

(i.e. $\chi^2_\alpha(2r) \leq -2 \ln(1 - q)$)

q	$1 - \alpha = .90$	$1 - \alpha = .95$	$1 - \alpha = .99$
	r	r	r
.70	3	3	4
.75	3	3	5
.80	3	4	5
.85	4	4	6
.90	4	5	6
.95	5	6	8
.97	6	7	8
.98	6	7	9
.99	7	8	10
.999	10	11	14

The upper tolerance limit given in (3) is a significant improvement over the tolerance limit given by Hanson and Koopmans (1964) for the IFR class. However they do not restrict attention to non-negative random variables. Also they do not obtain a lower tolerance limit for the IFR class.

<u>THEOREM</u> 3. If F is IFR, $F(0^-) = 0$ and $T > 0$ is specified, then

$$(5) \qquad P\left\{1 - F(T) \geq \delta\left(\frac{r}{n}\hat{\theta}_{r,n} - T\right) \exp\left[-\frac{\chi^2_{1-\alpha}(2r)\, T}{2r\,\hat{\theta}_{r,n}}\right]\right\} \geq 1 - \alpha$$

where

$$\delta(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 . \end{cases}$$

Johns and Lieberman (1965) study the problem of obtaining lower confidence limits on $1 - F(T)$ for the Weibull distribution. (5) is more convenient than their result. However, if $\frac{r}{n}\hat{\theta}_{r,n} < T$ our result is trivial. If $\frac{r}{n}\hat{\theta}_{r,n} > T$, then it is identical with the exponential lower confidence limit. In reliability applications where it is desired to establish high reliability the mean, hopefully, will far exceed T and therefore it seems quite likely that $\hat{\theta}_{r,n}$ will also.

<u>THEOREM</u> 4. If F is IFR, $F(0^-) = 0$ and $\theta = \int_0^\infty x\, dF(x)$, then

$$(6) \qquad P\left\{\theta \leq k_{\alpha,r}\,\hat{\theta}_{r,n}\right\} \geq 1 - \alpha$$

$$(7) \qquad P \left\{ \theta \geq \frac{\left[1 - \exp(- \frac{\chi^2_\alpha(2r)}{2n})\right]}{\chi^2_\alpha(2r)} \; 2r \; \hat{\theta}_{r,n} \right\} \geq 1 - \alpha ,$$

where

$$k_{\alpha,r} = \begin{cases} r & \text{if } \chi^2_\alpha(2r) \geq 2 \\ \dfrac{2r}{\chi^2_\alpha(2r)} & \text{if } \chi^2_\alpha(2r) < 2 . \end{cases}$$

Similar confidence limits can be obtained for higher order moments.

The upper confidence limit on θ in (6) is the usual exponential confidence limit when $\chi^2_\alpha(2r) < 2$. Unfortunately this condition is not satisfied for values of r greater than 3 or 4 at the usual significance levels.

In acceptance sampling the following hypothesis testing problem is considered:

$$H_o : \theta = \theta_o$$

$$\text{versus} \qquad H_1 : \theta < \theta_o .$$

The rejection region for the exponential case is of the form:

$$\text{Reject } H_o \text{ if } \quad \hat{\theta}_{r,n} \leq \frac{\theta_o \; \chi^2_\alpha(2r)}{2r} .$$

If $\chi^2_\alpha(2r) < 2$, then by (6) this test is also a size α test for the IFR case; i.e.

$$P \left\{ \hat{\theta}_{r,n} \leq \frac{\theta_o \; \chi^2_\alpha(2r)}{2r} \;\middle|\; F \text{ IFR} ; \; \theta > \theta_o \right\} \leq \alpha .$$

DFR Results

As we might expect, if a useful exponential confidence limit exists for a problem relative to IFR distributions, then no useful exponential confidence limit exists for the same problem relative to DFR distributions and conversely.

THEOREM 5. If F is DFR, $F(0^-) = 0$, $\zeta_q = \sup\{ x \mid F(x) \leq q \}$ and $\chi^2_{1-\alpha}(2r) < -2n \ln(1 - q)$, then

$$(8) \quad P \left\{ 1 - F\left[\frac{-2r \ln(1 - q)}{\chi^2_{1-\alpha}(2r)} \hat{\theta}_{r,n} \right] \geq 1 - q \right\} \geq 1 - \alpha$$

$$(9) \quad P \left\{ \zeta_q \geq \frac{-2r \ln(1 - q)}{\chi^2_{1-\alpha}(2r)} \hat{\theta}_{r,n} \right\} \geq 1 - \alpha .$$

If $\chi^2_{1-\alpha}(2r) \geq -2n \ln(1 - q)$ we can only make the trivial statement

$$P \{ \zeta_q \geq 0 \} \geq 1 - \alpha .$$

For most cases of practical interest -- high confidence and high population coverage -- (8) is not a useful result.

THEOREM 6. If F is DFR, $F(0^-) = 0$, $\zeta_q = \sup\{ x \mid F(x) \leq q \}$ and $\chi^2_{\alpha}(2r) > -2 \ln(1 - q)$, then

$$(10) \quad P \left\{ F\left[\frac{-2r \ln(1 - q)}{\chi^2_{\alpha}(2r)} \hat{\theta}_{r,n} \right] \geq q \right\} \geq 1 - \alpha$$

$$(11) \qquad P\left\{ \zeta_q \le \frac{-2r\ \ln(1-q)}{\chi^2_\alpha(2r)}\ \hat{\theta}_{r,n} \right\} \ge 1 - \alpha\ .$$

The upper confidence limit is trivial when $\chi^2_\alpha(2r) \le -2\ \ln(1 - q)$.

Table 2

Smallest values of r such that the exponential upper tolerance limit is a conservative upper tolerance limit for the DFR class.

(i.e. $\chi^2_\alpha(2r) > -2\ \ln(1 - q)$)

q	$1 - \alpha = .90$	$1 - \alpha = .95$	$1 - \alpha = .99$
	r	r	r
.70	4	4	5
.75	4	5	6
.80	4	5	6
.85	5	5	7
.90	5	6	7
.95	6	7	9
.97	7	8	10
.98	8	9	10
.99	8	9	11
.999	11	12	15

THEOREM 7. If F is DFR, $F(0^-) = 0$ and $T > 0$ is specified, then

$$(12) \qquad P\left\{1 - F(T) \geq \delta(T - r\hat{\theta}_{r,n}) \exp\left[\frac{-\chi^2_{1-\alpha}(2r)\ T}{2r\ \hat{\theta}_{r,n}}\right]\right\} \geq 1 - \alpha$$

where

$$\delta(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$$

as before.

THEOREM 8. If F is DFR, $F(0^-) = 0$, $\int_0^\infty x\, dF(x) = \theta$ and $\chi^2_\alpha(2r) < 2n$, then

$$(13) \qquad P\left\{\theta \geq \frac{2r\ \hat{\theta}_{r,n}}{\chi^2_\alpha(2r)}\right\} \geq 1 - \alpha$$

while if $\chi^2_\alpha(2r) \geq 2n$, then

$$(14) \qquad P\left\{\theta \geq \frac{r}{n}\hat{\theta}_{r,n} \exp\left[1 - \frac{\chi^2_\alpha(2r)}{2n}\right]\right\} \geq 1 - \alpha .$$

(13) holds for significance levels of practical interest when $r = n$.

3. Proofs of Theorems in Section 2.

Let Y denote a random variable with distribution G. If X has a continuous distribution F, note that $Y = G^{-1}F(X)$ has distribution G. We will repeatedly use the following lemma.

Lemma. If $G^{-1}F(x)$ is convex non-decreasing for $x \geq 0$, $G^{-1}F(0) = 0$ and $Y_i = G^{-1}F(X_i)$, then

$$(15) \qquad F\left[\sum_{i=1}^{n} a_i X_i\right] \underset{(\geq)}{\leq} G\left[\sum_{i=1}^{n} a_i Y_i\right]$$

when $a_i \geq 0$ and $\sum_{i=1}^{n} a_i = 1$ $(a_i \geq 1$ or $a_i = 0$, $i = 1,2,\ldots,n)$.

The proof is obvious.

In what follows it will be convenient to let

$$G(x) = \begin{cases} 1 - e^{-x} & x \geq 0 \\ 0 & x < 0 . \end{cases}$$

<u>Proof of Theorem 1.</u>

Since (1) and (2) are mathematically equivalent we need only prove (2). By the lemma we have

$$G\left[\sum_{i=1}^{r} \frac{Y_i + (n-r)Y_r}{n}\right] \geq F\left[\sum_{i=1}^{r} \frac{X_i + (n-r)X_r}{n}\right]$$

since $G^{-1}F(x)$ is convex when F is IFR. Now choose $k_{1-\alpha}$ so that

$$P\left\{G\left[\sum_{i=1}^{r} \frac{Y_i + (n-r)Y_r}{n}\right] \leq k_{1-\alpha}\right\} = 1 - \alpha$$

i.e. $\ln(1 - k_{1-\alpha}) = \frac{-\chi^2_{1-\alpha}(2r)}{2n}$. Since F is IFR we know (Barlow and Proschan (1965), p. 27) that

$$F(t;\zeta_q) \geq b(t;\zeta_q) = \begin{cases} 0 & t < \zeta_q \\ 1 - (1-q)^{t/\zeta_q} & t \geq \zeta_q \end{cases}$$

where ζ_q is the (unknown) q-th quantile.

Hence

$$G\left[\frac{\sum_{1}^{r} Y_i + (n-r)Y_r}{n}\right] \geq F(\tfrac{r}{n}\hat{\theta}_{r,n};\ \zeta_q) \geq b(\tfrac{r}{n}\hat{\theta}_{r,n};\ \zeta_q)$$

and

$$P\ \{\ b\ (\tfrac{r}{n}\hat{\theta}_{r,n};\ \zeta_q) \leq k_{1-\alpha}\ \} \geq 1-\alpha.$$

Since $b(t;\ \zeta_q)$ is non-increasing in ζ_q we have

$$P\ \{\ \zeta_q \geq b^{-1}(\tfrac{r}{n}\hat{\theta}_{r,n};\ k_{1-\alpha})\ \} \geq 1-\alpha$$

where the inverse is taken with respect to ζ_q .

<u>Case 1</u>. $k_{1-\alpha} \geq q$ (i.e. $\chi^2_{1-\alpha}(2r) \geq -2n\ \ln(1-q)$). From the following figure



we see that $b(\frac{r}{n}\hat{\theta}_{r,n};\ \zeta_q) \leq k_{1-\alpha}$ if and only if

$$\zeta_q \geq \frac{-2r\ \ln(1-q)\ \hat{\theta}_{r,n}}{\chi^2_{1-\alpha}(2r)}\ .$$

<u>Case 2</u>. $k_{1-\alpha} < q$ (i.e. $\chi^2_{1-\alpha}(2r) \leq -2n\ \ln(1-q)$). In this case

$b(\frac{r}{n}\hat{\theta}_{r,n};\ \zeta_q) \leq k_{1-\alpha}$ if and only if $\zeta_q > \frac{r}{n}\hat{\theta}_{r,n}$.

In either case we have

$$P\ \{\ \zeta_q \geq C_{1-\alpha,\ q}(r)\ \hat{\theta}_{r,n}\ \} \geq 1-\alpha$$

where

$$C_{1-\alpha,q}(r) = \begin{cases} \dfrac{-2r\ \ln(1-q)}{\chi^2_{1-\alpha}(2r)} & \text{if } \chi^2_{1-\alpha}(2r) > -2n\ \ln(1-q) \\ \\ \dfrac{r}{n} & \text{if } \chi^2_{1-\alpha}(2r) \leq -2n\ \ln(1-q) \end{cases}$$

Proof of Theorem 2. Again we need only prove statement (4). We use the following inequality which follows from the IFR assumption and the lemma:

$$G[\sum_1^r Y_i + (n-r)\ Y_r\ ] \leq F[\sum_1^r X_i + (n-r)\ X_r\ ] .$$

We choose k_α so that

$$P\ \{\ G\ [\ \sum_1^r Y_i + (n-r)\ Y_r\ ] \geq k_\alpha\ \} = 1 - \alpha$$

i.e. $\ln(1 - k_\alpha) = \dfrac{-\chi^2_\alpha(2r)}{2}$. From Barlow and Proschan (1965) p. 27 we have the sharp bound

$$F(t;\ \zeta_q) \leq B(t;\ \zeta_q) = \begin{cases} 1 - (1-q)^{t/\zeta_q} & t \leq \zeta_q \\ \\ 1 & t > \zeta_q . \end{cases}$$

Since $B(t;\ \zeta_q)$ is decreasing in ζ_q we have

$$P\left\{B(r\ \hat{\theta}_{r,n};\ \zeta_q) \geq k_\alpha\right\} = P\left\{\zeta_q \leq B^{-1}(r\ \hat{\theta}_{r,n};\ k_\alpha)\right\} \geq 1 - \alpha .$$

Case 1. $k_\alpha > q$ (i.e. $\chi^2_\alpha(2r) \geq -2\ \ln(1-q)$). From the following figure we see that $B(r\ \hat{\theta}_{r,n};\ \zeta_q) \geq k_\alpha$ if and only if $\zeta_q < r\ \hat{\theta}_{r,n}$.



<u>Case 2</u>. $k_\alpha \le q$ (i.e. $\chi^2_\alpha(2r) < -2\ln(1-q)$). In this case

$B(r\,\hat{\theta}_{r,n};\,\zeta_q) \ge k_\alpha$ if and only if $\zeta_q \le \dfrac{-2r\ln(1-q)}{\chi^2_\alpha(2r)}$.

Cases 1 and 2 together establish statement (4). ||

<u>Proof of Theorem 3</u>. Again we use the inequality

$$G\left[\frac{\sum_1^r Y_i + (n-r)\,Y_r}{n}\right] \ge F\left[\frac{\sum_1^r X_i + (n-r)\,X_r}{n}\right]$$

and choose $k_{1-\alpha}$ so that

$$P\left\{ G\left[\frac{\sum_1^r Y_i + (n-r)\,Y_r}{n}\right] \le k_{1-\alpha}\right\} = 1-\alpha$$

i.e. $\ln(1-k_{1-\alpha}) = \dfrac{\chi^2_{1-\alpha}(2r)}{2n}$. Let $p = F(T)$. We again use the sharp bound

$$F(t;\,p) \ge b(t;\,p) = \begin{cases} 0 & t < T \\ 1-(1-p)^{t/T} & t \ge T\,. \end{cases}$$

Then

$$P\left\{ b(\tfrac{r}{n}\,\hat{\theta}_{r,n};\,p) \le k_{1-\alpha}\right\} \ge 1-\alpha\,.$$

Since $b(t;\,p)$ is increasing in p , we have

$$P\left\{ 1-F(T) \ge 1-b^{-1}(\tfrac{r}{n}\,\hat{\theta}_{r,n};\,k_{1-\alpha})\right\} \ge 1-\alpha$$

(5) follows when we recall that $b(\frac{r}{n}\hat{\theta}_{r,n};\ p) = 0$ when $T > \frac{r}{n}\hat{\theta}_{r,n}$. ||

<u>Proof of Theorem 4</u>. To show (6) use the sharp bound

$$F(t;\ \theta) \leq B(t;\ \theta) = \begin{cases} 1 - e^{-t/\theta} & t \leq \theta \\ 1 & t > \theta \end{cases}$$

(Barlow and Proschan (1965), p. 27)

together with

$$G\ [\ \sum_1^r Y_i + (n - r)\ Y_r\ ] \leq F(r\ \hat{\theta}_{r,n};\ \theta) \leq B(r\ \hat{\theta}_{r,n};\ \theta)$$

to establish

$$P\ \{\ B(r\ \hat{\theta}_{r,n};\ \theta) \geq k_\alpha\ \} \geq 1 - \alpha$$

where $\ln(1 - k_\alpha) = \frac{-\chi^2_\alpha(2r)}{2}$ as before.

<u>Case 1</u>. $k_\alpha > 1 - e^{-1}$ (i.e. $\chi^2_\alpha(2r) > 2$). From the following figure we see that $B(r\ \hat{\theta}_{r,n};\ \theta) \geq k_\alpha$ if and only if $\theta < r\ \hat{\theta}_{r,n}$.



Case 2. $k_\alpha \le 1 - e^{-1}$ (i.e. $\chi^2_\alpha(2r) \le 2$). Also we see that

$B(r\,\hat{\theta}_{r,n};\ \theta) \ge k_\alpha$ if and only if $\theta \le \dfrac{r\,\hat{\theta}_{r,n}}{\chi^2_\alpha(2r)}$. Hence the result.

To show (7). Use the sharp bound

$$F(t;\ \theta) \ge b(t;\ \theta) = \begin{cases} 0 & t < \theta \\ 1 - e^{-wt} & t \ge \theta \end{cases}$$

where w depends on t and satifies

$$(16) \qquad \int_0^t e^{-wx}\,dx = \theta\ ,$$

(see Barlow and Proschan (1965) p. 28),

together with

$$G\left[\frac{\sum_1^r Y_i + (n-r)\,Y_r}{n}\right] \ge F(\tfrac{r}{n}\,\hat{\theta}_{r,n};\ \theta) \ge b(\tfrac{r}{n}\,\hat{\theta}_{r,n};\ \theta)$$

to assert

$$P\left\{ b(\tfrac{r}{n}\,\hat{\theta}_{r,n};\ \theta) \le k_{1-\alpha} \right\} \ge 1 - \alpha$$

where $\ln(1 - k_{1-\alpha}) = \dfrac{-\chi^2_\alpha(2r)}{2n}$ as before. Notice that $w = w(\theta)$ is a function of θ and is decreasing in θ. Hence

$$P\left\{ 1 - \exp\left[-w(\theta)\,\tfrac{r}{n}\,\hat{\theta}_{r,n}\right] \le k_{1-\alpha} \right\}$$

$$= P\left\{ w(\theta) \le \frac{-\ln(1 - k_{1-\alpha})}{\frac{r}{n}\,\hat{\theta}_{r,n}} \right\} \ge 1 - \alpha$$

or

$$P\left\{\theta \geq w^{-1}\left[\frac{-\ln(1-k_{1-\alpha})}{\frac{r}{n}\hat{\theta}_{r,n}}\right]\right\} \geq 1-\alpha .$$

Since $\theta = \dfrac{1-\exp\left[-w\frac{r}{n}\hat{\theta}_{r,n}\right]}{w}$ by (16) we have

$$w^{-1}\left[\frac{-\ln(1-k_{1-\alpha})}{\frac{r}{n}\hat{\theta}_{r,n}}\right] = \frac{\left[1-\exp\left(\frac{-\chi^2_\alpha(2r)}{2n}\right)\right]2r\,\hat{\theta}_{r,n}}{\chi^2_\alpha(2r)}$$

which establishes (7).

We omit proofs of the DFR results since they are a straightforward application of the same techniques applied to bounds on DFR distributions.

REFERENCES

BARLOW, R.E., and F. PROSCHAN. (1964). Exponential life test procedures when the distribution has monotone failure rate. Operations Research Center Report ORC 64-29, University of California, Berkeley.

BARLOW, R.E. and F. PROSCHAN. (1965). Mathematical Theory of Reliability. J. Wiley and Sons, New York.

DUBEY, S.D. (1962). On some statistical inferences for Weibull laws. Unpublished paper presented at the Annual Meeting of the American Statistical Association, Minneapolis, Minn., Sept. 10, 1962.

DUBEY, S.D. (1963). Asymptotic efficiencies of the moment estimators for the parameters of the Weibull laws. Unpublished paper presented at the Annual Meeting of the Institute of Mathematical Statistics, Cambridge, Mass. May 6-7.

EPSTEIN, B. and M. SOBEL. (1953). Life testing. J. Amer. Statist. Assoc., 48, 486-502.

GOODMAN, L. and A. MADANSKY. (1962). Parameter-free and nonparametric tolerance limits: the exponential case. Technometrics, 4, 75-95.

HANSON, D.L. and L.H. KOOPMANS. (1964). Tolerance limits for the class of distributions with increasing hazard rates. Ann. Math. Statist., 35, 1561-1570.

JOHNS, M.V. and G.J. LIEBERMAN. (1965). An exact asymptotically efficient confidence bound for reliability in the case of the Weibull distribution. Stanford University Technical Report.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY *(Corporate author)* University of California, Berkeley	2a. REPORT SECURITY CLASSIFICATION Unclassified 2b. GROUP

3. REPORT TITLE

Conservative Tolerance and Confidence Limits

4. DESCRIPTIVE NOTES *(Type of report and inclusive dates)*

Research Report

5. AUTHOR(S) *(Last name, first name, initial)*

Barlow, Richard E.

6. REPORT DATE September 1965	7a. TOTAL NO. OF PAGES 19	7b. NO. OF REFS 8

8a. CONTRACT OR GRANT NO. Nonr-3656(18) b. PROJECT NO. NR 042 238 c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) ORC 65-30 9b. OTHER REPORT NO(S) *(Any other numbers that may be assigned this report)*

10. AVAILABILITY/LIMITATION NOTICES Available upon request through:	Operations Research Center University of California Berkeley, California 94720
11. SUPPLEMENTARY NOTES Also sponsored by Army Research Office Contract DA-31-124-ARO-D-331	12. SPONSORING MILITARY ACTIVITY Logistics and Math. Stat. Branch

13. ABSTRACT

This paper extends the validity of exponential tolerance and confidence limits, under certain restrictions, to the class of distributions with monotone failure rate. In particular, the usual exponential lower tolerance limit is shown to be conservative for the increasing failure rate class of distributions in the range of population coverages and confidence coefficients of practical interest. Conservative confidence limits are also obtained on tail probabilities and moments.

DD FORM 1 JAN 64 1473

Unclassified

Unclassified

Security Classification

14. KEY WORDS	LINK A ROLE	LINK A WT	LINK B ROLE	LINK B WT	LINK C ROLE	LINK C WT
Conservative tolerance						
Confidence limits						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization *(corporate author)* issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers *(either by the originator or by the sponsor)*, also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

(1) "Qualified requesters may obtain copies of this report from DDC."

(2) "Foreign announcement and dissemination of this report by DDC is not authorized."

(3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through

_____."

(4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through

_____."

(5) "All distribution of this report is controlled. Qualified DDC users shall request through

_____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring *(paying for)* the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as *(TS)*, *(S)*, *(C)*, or *(U)*.

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.