https://ntrs.nasa.gov/search.jsp?R=19660014530 2018-07-25T00:27:22+00:00Z

ATLAS of PROTEIN SEQUENC and STRUCTURE 1965

N66-23819
(ACCESSION NUMBER)
(THRU)
111
(PAGES)
1
(CODE)
CR-71805
(NASA CR OR TMX OR AD NUMBER)
04
(CATEGORY)
FACILITY FORM 602

GPO PRICE $
CFSTI PRICE(S) $
Hard copy (HC) 4.00
Microfiche (MF) .75
ff 653 July 65

Margaret O. Dayhoff
Richard V. Eck
rie A. Chang
R. Sochard

NATIONAL BIOMEDICAL RESEARCH FOUNDATION
8600 16TH STREET
Silver Spring, Maryland

ATLAS of PROTEIN SEQUENCE and STRUCTURE 1965

Margaret O. Dayhoff
Richard V. Eck
Marie A. Chang
Minnie R. Sochard

NBR

NATIONAL BIOMEDICAL RESEARCH FOUNDATION
8600 16TH STREET
Silver Spring, Maryland

ATLAS OF PROTEIN SEQUENCE AND STRUCTURE

1965

Library of Congress Card Catalogue Number 65-29342

© Copyright 1965

by

The National Biomedical Research Foundation

Extra copies may be purchased from:

The National Biomedical Research Foundation

8600 16th Street, Silver Spring, Maryland

DEDICATION

To all the investigators who have developed the techniques necessary for the grand accomplishments represented by this tabulation, and to all those who have spent so much tedious effort in their application.

We would gratefully appreciate receiving suggestions, corrections, new data (even if fragmentary or provisional), and references to any data omitted from this volume.

M. O. D.
R. V. E.
M. A. C.
M. R. S.

PREFACE

This Atlas voluminously illustrates the triumph of experimental technique over the secretiveness of nature. Perhaps nowhere has the power of the scientific method been more brilliantly demonstrated than in the development of procedures for the study of the chemistry of life. As recently as twenty years ago, it was customary for biologists to have a hopeless attitude about biochemistry. Some details might be elicited, perhaps, but living things were thought to be so very complex and intricate that there surely was no hope of fully "understanding" them in all their chemical detail. Who, if he really comprehended the difficulty of the problem, would dare to think of man's ever knowing the detailed structure of a protein, for example, much less be able to synthesize it? Who would ever understand the mechanism of an enzyme as clearly as a chemist understands the details of an inorganic reaction? How could we ever hope to know the atomic details of a protein crystal?

Today some of these ambitions have already been attained, and the others no longer seem out of reach. We now rationally hope to be able to discover and understand the finest chemical details of living processes. These accomplishments and hopes have been made possible by the combined effect of several new approaches.

Techniques which permit the separation of chemically similar compounds have been developed for microgram samples. Among these are ion-exchange columns, paper chromatography, electrophoresis, and counter-current distribution. Radioactive tracer techniques and other micro-quantitative analytical procedures, often dependent on electronics and automation, have aided the analyses. X-ray crystallography, starting with the art of protein crystal production and ending with the processing of great numbers of experimental observations in the high-speed computer, has permitted a glimpse of three-dimensional structure.

Confidence in our understanding of experimental procedures and relationships among proteins has grown so great that sequences of amino acids are inferred from those found in homologous proteins. This technique requires only a small proportion of the analytical work needed to sequence a protein with no known relatives. The effectiveness of laboratory effort is thus magnified.

Some of the insights which have been developed cannot be attributed to any particular worker or school. Perhaps the greatest of these insights is that nature always uses "building blocks." A living cell is extremely complex and almost unimaginably intricate in detail. But it consists of a limited, understandable number of types of processes, reduplicated with variations. To understand the cell, we must have a few examples of each type of process, from which we can see the overall principles. For understanding, we need not work out the details of all the variations on these principles, although we may eventually choose to do so for medical or other practical reasons. Similarly, the analysis of such large, complex chemical molecules as proteins has been made possible by the recognition of their essential modularity, their building-block nature. Proteins are precise chemical structures built from regular subunits,

not statistical mixtures or hopelessly intricate molecular conglomerates as was once thought. It is by means of the discovery and utilization of such building block principles, combined with the large-scale application of new and improved techniques, that we now dare hope to make all of living nature accessible to our understanding.

Hidden in the amino acid sequence of a protein is the chemical information that produces its three-dimensional structure. In the case of an enzyme, this structure forms locks into which the proper keys—cell chemicals—fit. By these locks, the enzymes bring the proper reactants together quickly, efficiently, and selectively. Uncatalyzed reactions cannot complete with such specifically catalyzed reactions; therefore, the presence of enzymes determines which reactions can take place in living chemistry. In many cases, if not all, this three-dimensional structure is fully determined by the information in the one-dimensional sequence. The folding is the thermodynamically most stable result of all the possible intermolecular forces, such as hydrogen bonds and hydrophobic bonds, which can form between the various links of the chain. In principle, if we knew these forces in detail, and if we had appropriate computer routines, we should then be able to determine the three-dimensional structure of a protein, given only its amino acid sequence.

Also hidden in the sequences is information about the genes which directed their synthesis. For each amino acid there are a small number of possible corresponding nucleotide triplets in the gene. That is, each protein sequence corresponds to a limited number of possible nucleotide sequences. When nucleotide mutations occur, the substitution of alternative amino acids is not random. Analysis of amino acid sequence data, considered as a mathematical puzzle, can help elucidate both the mathematical details of the genetic code and the structural aspects of the genetic mechanism.

Hidden in each family of homologous sequences is the story of its evolution. Simple organisms, caught in their primitive ecological niches, still preserve even today enzymes performing primeval functions, held relatively fixed by natural selection. Even the older proteins of man are preserved as living "fossils" in his metabolism.

Enmeshed also in homologous sequences are the records of the many thousands of mutational steps by which we can quantify a phylogenetic tree. Each amino acid link is a trait by which we can trace species evolution. By comparison, the traditional taxonomic criteria are extremely vague and uncertain. In the case of distant relationships, they often break down completely. A truly quantitative and inclusive system of phylogenetic classification would be of great help to comparative physiologists and other students of evolution.

Conspicuous in comparative human protein sequences is information of great medical-diagnostic value. A long series of abnormalities has been found to be attributable to single amino acid replacements. One such tragically serious disease is sickle-cell anemia.

To facilitate the theoretical study of the protein sequences which have already been so ingeniously and laboriously determined, we have undertaken this compilation.

The information is kept in a compact, uniform format on punched cards. New information and corrections are easily inserted, while the text is kept accurate.

It is our intention to include the currently accepted amino acid sequence of every protein for which complete or substantial data is available. Uusally, only the definitive report giving the complete sequence from each laboratory will be referenced. If a substantial amount of work has been done on the same protein in other laboratories, their reports will also be referenced. We have also included some smaller peptides that have come to our attention. Unusual polypeptides which are presumably not produced by the genetic code have been omitted.

The format in which the Atlas is kept on punched cards is suitable for direct use in our computer programs. We use a three-letter code, which is a slight modification of the conventional notation, and also a mnemonic one-letter code which is clearer and much more suitable for certain comparative studies. We use a system of punctuation to describe the degree of confidence in each bond. Brief remarks are also included about the nature and function of the protein, and additional structural information such as the attachment of prosthetic groups, the location of S-S bonds, amino acids involved in active sites, and three-dimensional structures. In later editions we intend to include a section in which the alignment of all sequences of each family is given. Possibly we will also have sections on alleles and on mathematical methods and computer programs to treat the information.

This first edition is incomplete and imperfect and is intended mainly for distribution to investigators who have published protein sequence analyses, to acquaint them with the existence of this Atlas. We would gratefully appreciate their cooperation in making corrections, additions and suggestions for future editions. Since sequences are being reported in great numbers, we plan to bring out supplementary material in six months and a second edition in a year.

We thank all those who have assisted with this compilation, particularly Mr. Javier Albarran for his help with the computer aspects and Miss Lorrie Goldstein for her design of the cover.

The tabulations and computations were made at the University of Maryland Computer Science Center, College Park.

This work was supported by Grants GM-08710 and GM-12168 from the National Institutes of Health to the National Biomedical Research Foundation.

TABLE OF CONTENTS

		PAGE
PREFACE		0.001
CONTENTS		0.002
EXPLANATION OF NOTATION		0.010
I. CYTOCHROME C		1.000
BAKER'S YEAST	CY BY	1.001
CHICKEN	CY CH	1.002
HORSE	CY HO	1.003
HUMAN	CY HU	1.004
PIG AND BOVINE	CY PG	1.005
PSEUDOMONAS	CY PS	1.006
TUNA FISH	CY TF	1.007
BOMBYX MORI (SILKWORM)	CY SW	1.008
RATTLESNAKE	CY RS	1.009
RHODOSPIRILLUM RUBRUM	CY RR	1.010
SALMON	CY SM	1.011
II. GLOBINS		2.000
HEMOGLOBINS		2.000
HUMAN ALPHA	GL HUHA	2.001
HUMAN BETA	GL HUHB	2.002
HUMAN GAMMA	GL HUHG	2.003
GORILLA BETA	GL GOHB	2.004
HORSE BETA	GL HOHB	2.005
HORSE ALPHA	GL HOHA	2.006
LEMUR FULVUS BETA	GL LEHB	2.007
ABNORMAL HUMAN HEMOGLOBINS	GL HUH	2.020
MYOGLOBINS		2.100
WHALE	GL WHMY	2.101
III. OTHER RESPIRATORY PROTEINS		3.000
DIHEME PEPTIDE - CHROMATIUM	DH CH	3.001
FERREDOXIN - CLOSTRIDIUM PASTEURIANUM	FE CP	3.002
AZURIN - PSEUDOMONAS	AZ PS	3.003
IV. RIBONUCLEASE		4.000
BOVINE	RN BO	4.001
V. INHIBITORS		5.000
TRYPSIN INHIBITOR - BOVINE	TI BOPA	5.001

PAGE

VI. TOBACCO MOSAIC VIRUS 6.000

TOBACCO MOSAIC VIRUS . . . . . TM TM 6.001
TOBACCO MOSAIC VIRUS DAHLMENSE. . . . . TM TMD 6.002

VII. DIGESTIVE ENZYMES 7.000

CHYMOTRYPSINOGEN-A - BOVINE . . . . . TR BOCH 7.001
TRYPSINOGEN - BOVINE. . . . . TR BOTR 7.002
PAPAIN. . . . . PA PA 7.101
LYSOZYME - CHICKEN. . . . . LS CH 7.201

VIII. HORMONES 8.000

GLUCAGON

BOVINE . . . . . GN BO 8.001

PRESSINS

VASOPRESSIN ARGININE . . . . . PR BOAR 8.101
VASOPRESSIN LYSINE . . . . . PR PGLS 8.102
OXYTOCIN . . . . . PR BOOX 8.103
HYPERTENSIN . . . . . PR BOHY 8.104

CORTICOIDS

ALPHA MELANOCYTE-STIMULATING HORMONE
BOVINE, PIG, AND HORSE. . . . . TN BPAM 8.201
BETA MELANOCYTE-STIMULATING HORMONE
BOVINE . . . . . TN BOBM 8.202
PIG . . . . . TN PGBM 8.203
HORSE . . . . . TN HOBM 8.204
HUMAN . . . . . TN HUBM 8.205

BETA CORTICOTROPIN - PIG. . . . . TN PGAC 8.206
ALPHA CORTICOTROPIN - SHEEP AND BOVINE. . . . . TN SBAC 8.207

INSULINS

INSULIN A
BOVINE . . . . . IS BOA 8.301
BONITO. . . . . IS BNA 8.302
HORSE . . . . . IS HOA 8.303
SHEEP . . . . . IS SHA 8.304
SPERM WHALE, FIN-WHALE, PIG, AND HUMAN. . . . IS WPA 8.305
SEI-WHALE . . . . . IH WHA 8.306

INSULIN B
BOVINE, SHEEP, HORSE, HUMAN, PIG, AND
SPERM WHALE. . . . . IS BOB 8.321
BONITO . . . . . IS BNB 8.322

PAGE

IX. PLASMA PROTEINS 9.000

FIBRINOPEPTIDE A

BOVINE. . . . . FB BOA 9.001
SHEEP . . . . . FB SHA 9.002
GOAT. . . . . FB GTA 9.003
REINDEER. . . . . FB RDA 9.004
PIG . . . . . FB PGA 9.005
HUMAN . . . . . FB HUA 9.006
RABBIT. . . . . FB RTA 9.007

FIBRINOPEPTIDE B

BOVINE. . . . . FB BOB 9.101
SHEEP . . . . . FB SHB 9.102
GOAT. . . . . FB GTB 9.103
REINDEER. . . . . FB RDB 9.104
PIG . . . . . FB PGB 9.105
HUMAN . . . . . FB HUB 9.106
RABBIT. . . . . FB RTB 9.107

X. GLOBULINS 10.000

BENCE-JONES . . . . . BJ HU 10.001

AUTHOR INDEX 100.000

THE MEANING OF THE PUNCTUATION IS AS FOLLOWS.

BLANK	SEQUENCE OF AMINO ACIDS HAS BEEN DETERMINED.
( )	ENCLOSE PORTION OF SEQUENCE NOT SPECIFICALLY DETERMINED. TO PRESERVE PROPER SPACING,
=	IS USED INSTEAD OF)(
,	SEPARATES AMINO ACIDS WITHIN PARENTHESES,
BUT .	SEPARATES AMINO ACIDS, WHERE THE SEQUENCE IS PRESUMED BY HOMOLOGY WITH A KNOWN SEQUENCE.
/ OR ///	FRAGMENT, CONNECTION UNDETERMINED
* OR ***	CARBOXYL END OF PROTEIN

* ASTERISK BEFORE REFERENCE INDICATES THAT THE SEQUENCE WAS COPIED FROM, AND PROOFREAD AGAINST, THE ORIGINAL ARTICLE.

= BEFORE REFERENCE INDICATES THAT WE HAVE NOT SEEN THE ORIGINAL ARTICLE.

NO MARK BEFORE REFERENCE INDICATES OTHER GROUPS WHICH HAVE ALSO REPORTED WORK ON THE SAME PROTEIN.

BOTH SINGLE- AND THREE-LETTER NOTATIONS ARE USED, AS FOLLOWS.

A = ALA = ALANINE	M = MET = METHIONINE
C = CYS = CYSTEINE	N = ASN = ASPARAGINE
D = ASP = ASPARTIC ACID	O = TYR = TYROSINE
E = GLU = GLUTAMIC ACID	P = PRO = PROLINE
F = PHE = PHENYLALANINE	Q = GLN = GLUTAMINE
G = GLY = GLYCINE	R = ARG = ARGININE
H = HIS = HISTIDINE	S = SER = SERINE
I = ILU = ISOLEUCINE	T = THR = THREONINE
K = LYS = LYSINE	W = TRP = TRYPTOPHAN
L = LEU = LEUCINE	V = VAL = VALINE

B = ASX = ASPARTIC ACID OR ASPARAGINE
Z = GLX = GLUTAMIC ACID OR GLUTAMINE
X = XXX = UNDETERMINED OR OTHERWISE UNUSUAL

MNEMONICS OF THE ONE-LETTER CODE

IF POSSIBLE, THE INITIAL LETTER OF THE AMINO ACID IS USED.
IF MORE THAN ONE AMINO ACID BEGINS WITH THE SAME LETTER,
THE MOST COMMONLY-OCCURRING ONE IS ASSIGNED THE INITIAL.

A = ALANINE	I = ISOLEUCINE	S = SERINE
C = CYSTEINE	L = LEUCINE	T = THREONINE
G = GLYCINE	M = METHIONINE	V = VALINE
H = HISTIDINE	P = PROLINE	

SOME OF THE OTHERS ARE PHONETICALLY SUGGESTIVE.

F = PHENYLALANINE
R = ARGININE
O = TYROSINE

DOUBLE RING IN THE SIDE CHAIN.

W = TRYPTOPHAN

THE TWO ACIDS ARE ADJACENT, IN ALPHABETICAL ORDER.

D = ASPARTIC ACID
E = GLUTAMIC ACID

THE TWO AMINES HAVE LETTERS FROM THE MIDDLE OF THE ALPHABET.

N = ASPARAGINE (CONTAINS N)
Q = GLUTAMINE ('Q-TAMINE')

NON-INITIAL LETTER AS CLOSE AS POSSIBLE TO ITS INITIAL.

K = LYSINE

CYTOCHROME C - BAKER'S YEAST

HEME BONDED TO CYSTEINES AT POSITIONS 19 AND 22.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 T E F K A G S A K K G A T L F K T R C E L C H T V E K G G P
31 H K V G P N L H G I F G R H S G Q A Q G O S O T D A N I K K
61 N V L W D E N N M S E O L T N P K K O I P G T K M A F G G L
91 K K E K D R N D L I T O L K K A C E *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 THR GLU PHE LYS ALA GLY SER ALA LYS LYS GLY ALA THR LEU PHE
   LYS THR ARG CYS GLU LEU CYS HIS THR VAL GLU LYS GLY GLY PRO
31 HIS LYS VAL GLY PRO ASN LEU HIS GLY ILU PHE GLY ARG HIS SER
   GLY GLN ALA GLN GLY TYR SER TYR THR ASP ALA ASN ILU LYS LYS
61 ASN VAL LEU TRP ASP GLU ASN ASN MET SER GLU TYR LEU THR ASN
   PRO LYS LYS TYR ILU PRO GLY THR LYS MET ALA PHE GLY GLY LEU
91 LYS LYS GLU LYS ASP ARG ASN ASP LEU ILU THR TYR LEU LYS LYS
   ALA CYS GLU ***
```

COMPOSITION

7	ALA	A	2	GLN	Q	8	LEU	L	4	SER	S
3	ARG	R	7	GLU	E	16	LYS	K	8	THR	T
7	ASN	N	12	GLY	G	2	MET	M	1	TRP	W
4	ASP	D	4	HIS	H	4	PHE	F	5	TYR	O
3	CYS	C	4	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 108

* NARITA,K.,TITANI,K.,YAOI,Y.,MURAKAMI,H., BIOCHIM. BIOPHYS. ACTA, VOL. 77, PP.688-690, 1963

CYTOCHROME C - CHICKEN

ACETYL AT AMINO END.
HEME BONDED TO CYSTEINES AT POSITIONS 14 AND 17.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D I E K G K K I F V Q K C S Q C H T V E K G G K H K T G P
31 N L H G L F G R K T G Q A E G F S O T D A N K N K G I T W G
61 E D T L M E O L E N P K K O I P G T K M I F A G I K K K S E
91 R V D L I A O L K K A T N S *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP ILU GLU LYS GLY LYS LYS ILU PHE VAL GLN LYS CYS SER
   GLN CYS HIS THR VAL GLU LYS GLY GLY LYS HIS LYS THR GLY PRO
31 ASN LEU HIS GLY LEU PHE GLY ARG LYS THR GLY GLN ALA GLU GLY
   PHE SER TYR THR ASP ALA ASN LYS ASN LYS GLY ILU THR TRP GLY
61 GLU ASP THR LEU MET GLU TYR LEU GLU ASN PRO LYS LYS TYR ILU
   PRO GLY THR LYS MET ILU PHE ALA GLY ILU LYS LYS LYS SER GLU
91 ARG VAL ASP LEU ILU ALA TYR LEU LYS LYS ALA THR ASN SER ***
```

COMPOSITION

5	ALA	A	3	GLN	Q	6	LEU	L	4	SER	S
2	ARG	R	7	GLU	E	18	LYS	K	8	THR	T
5	ASN	N	13	GLY	G	2	MET	M	1	TRP	W
4	ASP	D	3	HIS	H	4	PHE	F	4	TYR	O
2	CYS	C	7	ILU	I	3	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 104

* MARGOLIASH,E.,NEEDLEMAN,S.B. AND STEWART,J.W., ACTA CHEM. SCAND., VOL.17, SUPPL.1, PP.250-256, 1963

CYTOCHROME C - HORSE

ACETYL AT AMINO END.
HEME BONDED TO CYSTEINES AT POSITIONS 14 AND 17.
OXIDATION-REDUCTION POTENTIAL EQUALS .250 V.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D V E K G K K I F V Q K C A Q C H T V E K G G K H K T G P
31 N L H G L F G R K T G Q A P G F T O T D A N K N K G I T W K
61 E E T L M E O L E N P K K O I P G T K M I F A G I K K K T E
91 R E D L I A O L K K A T N E *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP VAL GLU LYS GLY LYS LYS ILU PHE VAL GLN LYS CYS ALA
   GLN CYS HIS THR VAL GLU LYS GLY GLY LYS HIS LYS THR GLY PRO
31 ASN LEU HIS GLY LEU PHE GLY ARG LYS THR GLY GLN ALA PRO GLY
   PHE THR TYR THR ASP ALA ASN LYS ASN LYS GLY ILU THR TRP LYS
61 GLU GLU THR LEU MET GLU TYR LEU GLU ASN PRO LYS LYS TYR ILU
   PRO GLY THR LYS MET ILU PHE ALA GLY ILU LYS LYS LYS THR GLU
91 ARG GLU ASP LEU ILU ALA TYR LEU LYS LYS ALA THR ASN GLU ***
```

COMPOSITION

6	ALA	A	3	GLN	Q	6	LEU	L	0	SER	S
2	ARG	R	9	GLU	E	19	LYS	K	10	THR	T
5	ASN	N	12	GLY	G	2	MET	M	1	TRP	W
3	ASP	D	3	HIS	H	4	PHE	F	4	TYR	O
2	CYS	C	6	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 104

* MARGOLIASH,E., SMITH,E.L., KREIL,G.,AND TUPPY, H., NATURE, VOL.192, NO.4808, PP.1121-1127, DEC.23, 1961

CYTOCHROME C - HUMAN
ACETYL AT AMINO END.
HEME BONDED TO CYSTEINES AT POSITIONS 14 AND 17.
LEU (L) REPLACES MET (M) AT POSITION 65 IN 10 PERCENT
YIELD IN POOLED PROTEIN.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D V E K G K K I F I M K C S Q C H T V E K G G K H K T G P
31 N L H G L F G R K T G Q A P G O S O T A A N K N K G I I W G
61 E D T L M E O L E N P K K O I P G T K M I F V G I K K K E E
91 R A D L I A O L K K A T N E *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP VAL GLU LYS GLY LYS LYS ILU PHE ILU MET LYS CYS SER
   GLN CYS HIS THR VAL GLU LYS GLY GLY LYS HIS LYS THR GLY PRO
31 ASN LEU HIS GLY LEU PHE GLY ARG LYS THR GLY GLN ALA PRO GLY
   TYR SER TYR THR ALA ALA ASN LYS ASN LYS GLY ILU ILU TRP GLY
61 GLU ASP THR LEU MET GLU TYR LEU GLU ASN PRO LYS LYS TYR ILU
   PRO GLY THR LYS MET ILU PHE VAL GLY ILU LYS LYS LYS GLU GLU
91 ARG ALA ASP LEU ILU ALA TYR LEU LYS LYS ALA THR ASN GLU ***
```

COMPOSITION

6	ALA	A	2	GLN	Q	6	LEU	L	2	SER	S
2	ARG	R	8	GLU	E	18	LYS	K	7	THR	T
5	ASN	N	13	GLY	G	3	MET	M	1	TRP	W
3	ASP	D	3	HIS	H	3	PHE	F	5	TYR	O
2	CYS	C	8	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 104

* MATSUBARA, H., AND SMITH,E.L., J. BIOL. CHEM., VOL. 237, NO.11, PC3575-PC3576, NOV., 1962

CYTOCHROME - C PIG AND BOVINE

ACETYL AT AMINO END.
HEME BONDED TO CYSTEINES AT POSITIONS 14 AND 17.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D V E K G K K I F V Q K C A Q C H T V E K G G K H K T G P
31 N L H G L F G R K T G Q A P G F S O T D A N K N K G I T W G
61 E E T L M E O L E N P K K O I P G T K M I F A G I K K K G E
91 R E D L I A O L K K A T N E *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP VAL GLU LYS GLY LYS LYS ILU PHE VAL GLN LYS CYS ALA
   GLN CYS HIS THR VAL GLU LYS GLY GLY LYS HIS LYS THR GLY PRO
31 ASN LEU HIS GLY LEU PHE GLY ARG LYS THR GLY GLN ALA PRO GLY
   PHE SER TYR THR ASP ALA ASN LYS ASN LYS GLY ILU THR TRP GLY
61 GLU GLU THR LEU MET GLU TYR LEU GLU ASN PRO LYS LYS TYR ILU
   PRO GLY THR LYS MET ILU PHE ALA GLY ILU LYS LYS LYS GLY GLU
91 ARG GLU ASP LEU ILU ALA TYR LEU LYS LYS ALA THR ASN GLU ***
```

COMPOSITION

6	ALA	A	3	GLN	Q	6	LEU	L	1	SER	S
2	ARG	R	9	GLU	E	18	LYS	K	8	THR	T
5	ASN	N	14	GLY	G	2	MET	M	1	TRP	W
3	ASP	D	3	HIS	H	4	PHE	F	4	TYR	O
2	CYS	C	6	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 104

* MARGOLIASH,E.,NEEDLEMAN,S.B. AND STEWART,J.W., ACTA CHEM. SCAND., VOL.17, SUPPL.1, PP.250-256, 1963 (PIG)

TUPPY H., AND PALEUS, S., ACTA CHEM. SCAND., VOL. 13, NO.4, PP. 641-646, 1959, (HEME ATTACHEMENT REGION ONLY - BOVINE)

YASUNOBU, K. T., NAKASHIMA, T., HIGA, H., MATSUBARA, H., AND BENSON, A. , BIOCHIM. BIOPHYS. ACTA VOL. 78, PN1324 PP. 791-794, 1963 (BOVINE)

CYTOCHROME C - PSEUDOMONAS

HEME BONDED TO CYSTEINES AT POSITIONS 12 AND 15.
THE AMINO END IS NOT ACETYLATED.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 E D P E V L F K N K G C V A C H A I D T K M V G P A O K D V
31 A A K F A G Q A G A E A E L A Q R I K N G S Q G V W G P I P
61 M P P N A V S D D E A Q T L A K W V L S Q K *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLU ASP PRO GLU VAL LEU PHE LYS ASN LYS GLY CYS VAL ALA CYS
   HIS ALA ILU ASP THR LYS MET VAL GLY PRO ALA TYR LYS ASP VAL
31 ALA ALA LYS PHE ALA GLY GLN ALA GLY ALA GLU ALA GLU LEU ALA
   GLN ARG ILU LYS ASN GLY SER GLN GLY VAL TRP GLY PRO ILU PRO
61 MET PRO PRO ASN ALA VAL SER ASP ASP GLU ALA GLN THR LEU ALA
   LYS TRP VAL LEU SER GLN LYS ***
```

COMPOSITION

13	ALA	A	5	GLN	Q	4	LEU	L	3	SER	S
1	ARG	R	5	GLU	E	8	LYS	K	2	THR	T
3	ASN	N	7	GLY	G	2	MET	M	2	TRP	W
5	ASP	D	1	HIS	H	2	PHE	F	1	TYR	O
2	CYS	C	3	ILU	I	6	PRO	P	7	VAL	V

TOTAL NO. OF ACIDS = 82

* AMBLER, R. P., BIOCHEM. J., VOL.89, P.349-378. 1963

CYTOCHROME C - TUNA FISH

ACETYL AT AMINO END.
HEME BONDED TO CYSTEINES AT POSITIONS 14 AND 17.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D V A K.G K K.T F V Q K.C A Q(C.H)T V E N G G K.H K(V.G.P.
31 N)L W.G L F.G R.K T(G.Q)A E G O.S O T(D.A.N)K.S K.G I V W(N,
61 N,D)T L M E O.L E N P K K.O(I.P.G)T K(M.I)F.A G I K K.K G E
91 R.Q D L(V.A)O.L K S T A S •
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP VAL ALA LYS.GLY LYS LYS.THR PHE VAL GLN LYS.CYS ALA
   GLN(CYS.HIS)THR VAL GLU ASN GLY GLY LYS.HIS LYS(VAL.GLY.PRO.
31 ASN)LEU TRP.GLY LEU PHE.GLY ARG.LYS THR(GLY.GLN)ALA GLU GLY
   TYR.SER TYR THR(ASP.ALA.ASN)LYS.SER LYS.GLY ILU VAL TRP(ASN,
61 ASN,ASP)THR LEU MET GLU TYR.LEU GLU ASN PRO LYS LYS.TYR(ILU.
   PRO.GLY)THR LYS.MET.ILU)PHE.ALA GLY ILU LYS LYS.LYS GLY GLU
91 ARG.GLN ASP LEU(VAL.ALA)TYR.LEU LYS SER THR ALA SER ***
```

COMPOSITION

7	ALA	A	4	GLN	Q	6	LEU	L	4	SER	S
2	ARG	R	5	GLU	E	16	LYS	K	7	THR	T
6	ASN	N	13	GLY	G	2	MET	M	2	TRP	W
4	ASP	D	2	HIS	H	3	PHE	F	5	TYR	O
2	CYS	C	4	ILU	I	3	PRO	P	6	VAL	V

TOTAL NO. OF ACIDS = 103

* KREIL,G., Z. PHYSIOL. CHEM., BD. 334, PP.154-166, 1963

CYTOCHROME C - BOMBYX MORI (SILKWORM)

HEME BONDED TO CYSTEINES AT POSITIONS 4 AND 7 OF FRAGMENT.

```
  1 2 3 4 5 6 7 8 9 0 1
/ V Q R C A Q C H T(V,E)/
```

```
     1   2   3   4   5   6   7   8   9  10  11
/// VAL GLN ARG CYS ALA GLN CYS HIS THR(VAL,GLU)///
```

COMPOSITION OF FRAGMENT

1	ALA	A	2	GLN	Q	0	LEU	L	0	SER	S
1	ARG	R	1	GLU	E	0	LYS	K	1	THR	T
0	ASN	N	0	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	1	HIS	H	0	PHE	F	0	TYR	O
2	CYS	C	0	ILU	I	0	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS IN FRAGMENT = 11

* TUPPY H., Z. NATURFORSCH., VOL.12, PP.784-788, 1957

CYTOCHROME C - RATTLESNAKE

ACETYL AT AMINO END.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 G D V E K G K K I F(I.T.K.C.S.Q.C.H.T.V.E.K.G.G.K.H)K T G P
31 N L H.G L F.G R K T G Q A V G O.S O.T A A N K N.K G I I W.G
61 D D T L M E O.L E N P K K O.I P G T K M.V F.T G L.S K K K E
91 R T N L.I A O.L K E K T A A *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 GLY ASP VAL GLU LYS GLY LYS LYS ILU PHE(ILU.THR.LYS.CYS.SER.
   GLN.CYS.HIS.THR.VAL.GLU.LYS.GLY.GLY.LYS.HIS)LYS THR GLY PRO
31 ASN LEU HIS.GLY LEU PHE.GLY ARG LYS THR GLY GLN ALA VAL GLY
   TYR.SER TYR.THR ALA ALA ASN LYS ASN.LYS GLY ILU ILU TRP.GLY
61 ASP ASP THR LEU MET GLU TYR.LEU GLU ASN PRO LYS LYS TYR.ILU
   PRO GLY THR LYS MET.VAL PHE.THR GLY LEU.SER LYS LYS LYS GLU
91 ARG THR ASN LEU.ILU ALA TYR.LEU LYS GLU LYS THR ALA ALA ***
```

COMPOSITION

6	ALA	A	2	GLN	Q	7	LEU	L	3	SER	S
2	ARG	R	6	GLU	E	18	LYS	K	10	THR	T
5	ASN	N	13	GLY	G	2	MET	M	1	TRP	W
3	ASP	D	3	HIS	H	3	PHE	F	5	TYR	O
2	CYS	C	6	ILU	I	3	PRO	P	4	VAL	V

TOTAL NO. OF ACIDS = 104

* BAHL, O. P. AND SMITH, E. L., J. BIOL. CHEM., VOL.240, NO.9, PP.3585-3593, SEPT., 1965

CYTOCHROME - C RHODOSPIRILLUM RUBRUM

HEME BONDED TO CYSTEINES AT POSITIONS 1 AND 4 OF FRAGMENT.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3
/ C L A C H T F B Z G A N K /

     1   2   3   4   5   6   7   8   9  10  11  12  13
/// CYS LEU ALA CYS HIS THR PHE ASX GLX GLY ALA ASN LYS ///
```

COMPOSITION OF FRAGMENT

2	ALA	A	0	GLN	Q	1	LEU	L	0	SER	S
0	ARG	R	0	GLU	E	1	LYS	K	1	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	1	HIS	H	1	PHE	F	0	TYR	O
2	CYS	C	0	ILU	I	0	PRO	P	0	VAL	V
1	ASX	B	1	GLX	Z						

TOTAL NO. OF ACIDS IN FRAGMENT = 13

* TUPPY H., AND PALEUS, S., ACTA CHEM. SCAND., VOL. 13, NO.4, PP. 641-646, 1959

CYTOCHROME C - SALMON

HEME BONDED TO CYSTEINES AT POSITIONS 4 AND 7 OF FRAGMENT.

```
1 2 3 4 5 6 7 8 9 0 1
/ V Q K C A Q H C T(V,E)/
```

```
     1   2   3   4   5   6   7   8   9  10  11
/// VAL GLN LYS CYS ALA GLN CYS HIS THR(VAL,GLU)///
```

COMPOSITION OF FRAGMENT

1	ALA	A	2	GLN	Q	0	LEU	L	0	SER	S
0	ARG	R	1	GLU	E	1	LYS	K	1	THR	T
0	ASN	N	0	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	1	HIS	H	0	PHE	F	0	TYR	O
2	CYS	C	0	ILU	I	0	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS IN FRAGMENT = 11

* TUPPY H., AND PALEUS, S., ACTA CHEM. SCAND., VOL. 9, P.353-364, 1955

HEMOGLOBIN ALPHA - HUMAN

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 V L S P A D K T N V K A A W G K V G A H A G E O G A E A L E
 31 R M F L S F P T T K T O F P H F D L S H G S A Q V K G H G K
 61 K V A D A L T N A V A H V D D M P N A L S A L S D L H A H K
 91 L R V D P V N F K L L S H C L L V T L A A H L P A E F T P A
121 V H A S L D K F L A S V S T V L T S K O R *

     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 VAL LEU SER PRO ALA ASP LYS THR ASN VAL LYS ALA ALA TRP GLY
    LYS VAL GLY ALA HIS ALA GLY GLU TYR GLY ALA GLU ALA LEU GLU
 31 ARG MET PHE LEU SER PHE PRO THR THR LYS THR TYR PHE PRO HIS
    PHE ASP LEU SER HIS GLY SER ALA GLN VAL LYS GLY HIS GLY LYS
 61 LYS VAL ALA ASP ALA LEU THR ASN ALA VAL ALA HIS VAL ASP ASP
    MET PRO ASN ALA LEU SER ALA LEU SER ASP LEU HIS ALA HIS LYS
 91 LEU ARG VAL ASP PRO VAL ASN PHE LYS LEU LEU SER HIS CYS LEU
    LEU VAL THR LEU ALA ALA HIS LEU PRO ALA GLU PHE THR PRO ALA
121 VAL HIS ALA SER LEU ASP LYS PHE LEU ALA SER VAL SER THR VAL
    LEU THR SER LYS TYR ARG ***
```

COMPOSITION

21	ALA	A	1	GLN	Q	18	LEU	L	11	SER	S
3	ARG	R	4	GLU	E	11	LYS	K	9	THR	T
4	ASN	N	7	GLY	G	2	MET	M	1	TRP	W
8	ASP	D	10	HIS	H	7	PHE	F	3	TYR	O
1	CYS	C	0	ILU	I	7	PRO	P	13	VAL	V

TOTAL NO. OF ACIDS = 141

* HILL,R.J., AND KONIGSBERG, W., J. BIOL. CHEM., VOL. 237, NO.10, PP. 3151-3156, OCT., 1962

BRAUNITZER, G., GEHRING-MULLER, R., HILSCHMANN, N., HILSE, K., HOBOM, G., RUDLOFF, V., AND WITTMANN-LIEBOLD, B., Z. PHYSIOL. CHEM., VOL. BD 325, PP.283-286, 1961

THE SAME SEQUENCE, WITHOUT DISTINGUISHING AMINES, ALSO REPORTED IN THE ARTICLE.

SCHROEDER,W.A., J.R.SHELTON, J.B.SHELTON, AND J.CORMICK BIOCHEMISTRY, VOL. 2, NO.6, PP.1353-1357, NOV.-DEC., 1963

FETAL ALPHA CHAIN IS VERY PROBABLY IDENTICAL WITH ADULT ALPHA CHAIN. TRYPTIC AND CHYMOTRYPTIC PEPTIDES, MOST OF WHICH WERE COMPLETELY SEQUENCED, WERE SHOWN TO FIT EXACTLY INTO THE ADULT ALPHA CHAIN SEQUENCE.

HEMOGLOBIN BETA - HUMAN

```
     1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
   1 V H L T P E E K S A V T A L W G K V D V D E V G G E A L G R
  31 L L V V O P W T E R F F E S F G D L S T P D A V M G D P K V
  61 K A H G K K V L G A F S D G L A H L D D L K G T F A T L S E
  91 L H C D K L H V D P E D F R L L G D V L V C V L A H H F G K
 121 E F T P P V E A A O E K V V A G V A D A L A H K O H *
      1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
   1 VAL HIS LEU THR PRO GLU GLU LYS SER ALA VAL THR ALA LEU TRP
     GLY LYS VAL ASP VAL ASP GLU VAL GLY GLY GLU ALA LEU GLY ARG
  31 LEU LEU VAL VAL TYR PRO TRP THR GLU ARG PHE PHE GLU SER PHE
     GLY ASP LEU SER THR PRO ASP ALA VAL MET GLY ASP PRO LYS VAL
  61 LYS ALA HIS GLY LYS LYS VAL LEU GLY ALA PHE SER ASP GLY LEU
     ALA HIS LEU ASP ASP LEU LYS GLY THR PHE ALA THR LEU SER GLU
  91 LEU HIS CYS ASP LYS LEU HIS VAL ASP PRO GLU ASP PHE ARG LEU
     LEU GLY ASP VAL LEU VAL CYS VAL LEU ALA HIS HIS PHE GLY LYS
 121 GLU PHE THR PRO PRO VAL GLU ALA ALA TYR GLU LYS VAL VAL ALA
     GLY VAL ALA ASP ALA LEU ALA HIS LYS TYR HIS ***
```

COMPOSITION

15	ALA	A	0	GLN	Q	18	LEU	L	5	SER	S
3	ARG	R	11	GLU	E	11	LYS	K	7	THR	T
0	ASN	N	13	GLY	G	1	MET	M	2	TRP	W
13	ASP	D	9	HIS	H	8	PHE	F	3	TYR	O
2	CYS	C	0	ILU	I	7	PRO	P	18	VAL	V

TOTAL NO. OF ACIDS = 146

• BRAUNITZER, G., GEHRING-MULLER, R., HILSCHMANN, N., HILSE, K., HOBOM, G., RUDLOFF, V., AND WITTMANN-LIEBOLD, B., Z. PHYSIOL. CHEM., VOL. BD 325, PP.283-286, 1961

HEMOGLOBIN GAMMA - HUMAN

```
     1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
   1 G H F T E E D K A T I T S L W G K V N V E D A G G E T L G R
  31 L L V V O P W T Q R F F D S F G N L S S A S A I M G N P K V
  61 K A H G K K V L T S L G D A I K H L D D L K G T F A Q L S E
  91 L H C D K L H V D P E N F K L L G N V L V T V L A I H F G K
 121 E F T P E V Q A S W Q K M V T G V A S A L S S R O H *
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
   1 GLY HIS PHE THR GLU GLU ASP LYS ALA THR ILU THR SER LEU TRP
     GLY LYS VAL ASN VAL GLU ASP ALA GLY GLY GLU THR LEU GLY ARG
  31 LEU LEU VAL VAL TYR PRO TRP THR GLN ARG PHE PHE ASP SER PHE
     GLY ASN LEU SER SER ALA SER ALA ILU MET GLY ASN PRO LYS VAL
  61 LYS ALA HIS GLY LYS LYS VAL LEU THR SER LEU GLY ASP ALA ILU
     LYS HIS LEU ASP ASP LEU LYS GLY THR PHE ALA GLN LEU SER GLU
  91 LEU HIS CYS ASP LYS LEU HIS VAL ASP PRO GLU ASN PHE LYS LEU
     LEU GLY ASN VAL LEU VAL THR VAL LEU ALA ILU HIS PHE GLY LYS
 121 GLU PHE THR PRO GLU VAL GLN ALA SER TRP GLN LYS MET VAL THR
     GLY VAL ALA SER ALA LEU SER SER ARG TYR HIS ***
```

COMPOSITION

11	ALA	A	4	GLN	Q	17	LEU	L	11	SER	S
3	ARG	R	8	GLU	E	12	LYS	K	10	THR	T
5	ASN	N	13	GLY	G	2	MET	M	3	TRP	W
8	ASP	D	7	HIS	H	8	PHE	F	2	TYR	O
1	CYS	C	4	ILU	I	4	PRO	P	13	VAL	V

TOTAL NO. OF ACIDS = 146

* SCHROEDER,W.A., SHELTON, J.R., SHELTON,J.B., CORMICK, J., AND JONES,R.T., BIOCHEMISTRY, VOL.2, NO.5, PP.992-1008, SEPT.-OCT., 1963

HEMOGLOBIN BETA - GORILLA

```
     1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
   1 V H L T P E E K S A V T A L W G K V D V D E V G G E A L G R
  31 L L V V O P W T E R F F E S F G D L S T P D A V M G D P K V
  61 K A H G K K V L G A F S D G L A H L D D L K G T F A T L S E
  91 L H C D K L H V D P E D F L L L G D V L V C V L A H H F G K
 121 E F T P P V E A A O E K V V A G V A D A L A H K O H *
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
   1 VAL HIS LEU THR PRO GLU GLU LYS SER ALA VAL THR ALA LEU TRP
     GLY LYS VAL ASP VAL ASP GLU VAL GLY GLY GLU ALA LEU GLY ARG
  31 LEU LEU VAL VAL TYR PRO TRP THR GLU ARG PHE PHE GLU SER PHE
     GLY ASP LEU SER THR PRO ASP ALA VAL MET GLY ASP PRO LYS VAL
  61 LYS ALA HIS GLY LYS LYS VAL LEU GLY ALA PHE SER ASP GLY LEU
     ALA HIS LEU ASP ASP LEU LYS GLY THR PHE ALA THR LEU SER GLU
  91 LEU HIS CYS ASP LYS LEU HIS VAL ASP PRO GLU ASP PHE LEU LEU
     LEU GLY ASP VAL LEU VAL CYS VAL LEU ALA HIS HIS PHE GLY LYS
 121 GLU PHE THR PRO PRO VAL GLU ALA ALA TYR GLU LYS VAL VAL ALA
     GLY VAL ALA ASP ALA LEU ALA HIS LYS TYR HIS ***
```

COMPOSITION

15	ALA	A	0	GLN	Q	19	LEU	L	5	SER	S
2	ARG	R	11	GLU	E	11	LYS	K	7	THR	T
0	ASN	N	13	GLY	G	1	MET	M	2	TRP	W
13	ASP	D	9	HIS	H	8	PHE	F	3	TYR	O
2	CYS	C	0	ILU	I	7	PRO	P	18	VAL	V

TOTAL NO. OF ACIDS = 146

= ZUCKERKANDL, E., SCIENTIFIC AMERICAN, VOL. 212, NO. 5,
PP. 110-118, MAY 1965

HEMOGLOBIN BETA - HORSE

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 V E L S G E E K A A L(V,A,L,W,D)K V D E E E V G(G.E.A)L G R
 31 L L V V O P W T E R F(F.E.S.F.G.D.L.S.G.P.D.A.V)M(G.D.P)K V
 61 K A H G K K V L H S F G E G V H H(L.D.D.L)K G T F A(A.L.S.E.
 91 L.H.C.D.K.L.H.V.D.P.E.D.F)R L L G D V L A L V V A R H F G K
121 D F T P E L E A S O E K V V A G V A D A L A H K O H *

     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 VAL GLU LEU SER GLY GLU GLU LYS ALA ALA LEU(VAL,ALA,LEU,TRP,
    ASP)LYS VAL ASP GLU GLU GLU VAL GLY(GLY.GLU.ALA)LEU GLY ARG
 31 LEU LEU VAL VAL TYR PRO TRP THR GLU ARG PHE(PHE.GLU.SER.PHE.
    GLY.ASP.LEU.SER.GLY.PRO.ASP.ALA.VAL)MET(GLY.ASP.PRO)LYS VAL
 61 LYS ALA HIS GLY LYS LYS VAL LEU HIS SER PHE GLY GLU GLY VAL
    HIS HIS(LEU.ASP.ASP.LEU)LYS GLY THR PHE ALA(ALA.LEU.SER.GLU.
 91 LEU.HIS.CYS.ASP.LYS.LEU.HIS.VAL.ASP.PRO.GLU.ASP.PHE)ARG LEU
    LEU GLY ASP VAL LEU ALA LEU VAL VAL ALA ARG HIS PHE GLY LYS
121 ASP PHE THR PRO GLU LEU GLU ALA SER TYR GLU LYS VAL VAL ALA
    GLY VAL ALA ASP ALA LEU ALA HIS LYS TYR HIS ***
```

COMPOSITION

15	ALA	A	0	GLN	Q	19	LEU	L	6	SER	S
4	ARG	R	15	GLU	E	11	LYS	K	3	THR	T
0	ASN	N	14	GLY	G	1	MET	M	2	TRP	W
13	ASP	D	9	HIS	H	8	PHE	F	3	TYR	O
1	CYS	C	0	ILU	I	5	PRO	P	17	VAL	V

TOTAL NO. OF ACIDS = 146

* SMITH, D. B., CAN. J. BIOCHEM., VOL.42, NO.5, PP.755-762, 1964

HEMOGLOBIN ALPHA - HORSE

```
     1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
   1 V L S A A D K T N V K A A W S K V G G H A G E O G A E A L E
  31 R M F L G F P T T K T O F P H F D L S H G S A Q V K A H G K
  61 K V A D G L T L A V G H L D D L P G A L S N L S D L H A H K
  91 L R V D P V N F K L L S H C L L S T L A V H L P N D F T P A
 121 V H A S L D K F L S S V S T V L T S K O R *

      1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
   1 VAL LEU SER ALA ALA ASP LYS THR ASN VAL LYS ALA ALA TRP SER
     LYS VAL GLY GLY HIS ALA GLY GLU TYR GLY ALA GLU ALA LEU GLU
  31 ARG MET PHE LEU GLY PHE PRO THR THR LYS THR TYR PHE PRO HIS
     PHE ASP LEU SER HIS GLY SER ALA GLN VAL LYS ALA HIS GLY LYS
  61 LYS VAL ALA ASP GLY LEU THR LEU ALA VAL GLY HIS LEU ASP ASP
     LEU PRO GLY ALA LEU SER ASN LEU SER ASP LEU HIS ALA HIS LYS
  91 LEU ARG VAL ASP PRO VAL ASN PHE LYS LEU LEU SER HIS CYS LEU
     LEU SER THR LEU ALA VAL HIS LEU PRO ASN ASP PHE THR PRO ALA
 121 VAL HIS ALA SER LEU ASP LYS PHE LEU SER SER VAL SER THR VAL
     LEU THR SER LYS TYR ARG ***
```

COMPOSITION

16	ALA	A	1	GLN	Q	21	LEU	L	13	SER	S
3	ARG	R	3	GLU	E	11	LYS	K	9	THR	T
4	ASN	N	10	GLY	G	1	MET	M	1	TRP	W
9	ASP	D	10	HIS	H	7	PHE	F	3	TYR	O
1	CYS	C	0	ILU	I	6	PRO	P	12	VAL	V

TOTAL NO. OF ACIDS = 141

* BRAUNITZER, G. AND MATSUDA, G., J. BIOCHEM. (TOKYO), VOL.53, NO.3, PP.262-263, 1963

THIS SEQUENCE WAS DETERMINED PARTIALLY BY HOMOLOGY WITH HUMAN ALPHA.

HEMOGLOBIN BETA - LEMUR FULVUS

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 T L L S A E E D A H V T S L W G K V N V E K V G G E A L G R
 31 L L V V(O,P,W,T,E,R,F,F,E,S,F,G,D=L,S,S,P,S,A,V,M,G,D,P,K,V,
 61 K,A,H,G,K,K,V,L,S,A,F,S,E,G=L,H,H,L,D,D,L,K,G,T,F,A,A,L,S,E,
 91 L,H,C,V,A,L,H,V,D,P,E,D,F,K,L,L,G,D,S,L,S,D,V,L,A,D,H,F,G,K)
121 X X X X X X X X X X X X V V A G V(A,D,A,L,A,H,K,O,H)*
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 THR LEU LEU SER ALA GLU GLU ASP ALA HIS VAL THR SER LEU TRP
    GLY LYS VAL ASN VAL GLU LYS VAL GLY GLY GLU ALA LEU GLY ARG
 31 LEU LEU VAL VAL(TYR,PRO,TRP,THR,GLU,ARG,PHE,PHE,GLU,SER,PHE,
    GLY,ASP=LEU,SER,SER,PRO,SER,ALA,VAL,MET,GLY,ASP,PRO,LYS,VAL,
 61 LYS,ALA,HIS,GLY,LYS,LYS,VAL,LEU,SER,ALA,PHE,SER,GLU,GLY=LEU,
    HIS,HIS,LEU,ASP,ASP,LEU,LYS,GLY,THR,PHE,ALA,ALA,LEU,SER,GLU,
 91 LEU,HIS,CYS,VAL,ALA,LEU,HIS,VAL,ASP,PRO,GLU,ASP,PHE,LYS,LEU,
    LEU,GLY,ASP,SER,LEU,SER,ASP,VAL,LEU,ALA,ASP,HIS,PHE,GLY,LYS)
121 XXX XXX XXX XXX XXX XXX XXX XXX XXX XXX XXX XXX VAL VAL ALA
    GLY VAL(ALA,ASP,ALA,LEU,ALA,HIS,LYS,TYR,HIS)***
```

COMPOSITION

14	ALA	A	0	GLN	Q	19	LEU	L	11	SER	S
2	ARG	R	9	GLU	E	10	LYS	K	4	THR	T
1	ASN	N	12	GLY	G	1	MET	M	2	TRP	W
11	ASP	D	9	HIS	H	7	PHE	F	2	TYR	O
1	CYS	C	0	ILU	I	4	PRO	P	15	VAL	V
12	XXX	X									

TOTAL NO. OF ACIDS = 146

* BUETTNER-JANUSCH, J. AND HILL, R. L., SCIENCE, VOL. 147, PP. 836-842, FEB. 19, 1965

ABNORMAL HUMAN HEMOGLOBIN

Normal adult human hemoglobin (hemoglobin A) contains two pairs of polypeptide chains, termed alpha and beta. Each pair is identical. Some modified beta chains have been given other Greek letters, for example, normal fetal hemoglobin is composed of two alpha chains and two "gamma" chains. Usually, however, altered hemoglobins are different in only a single amino acid. A number of hemoglobins bearing these altered amino acid sequences in their polypeptide chains have been described. For example, one of the early reports by Ingram (1957) shows the chemical difference between normal human hemoglobin and sickle cell hemoglobin. By comparison of amino acid sequences of tryptic peptide digests of the two hemoglobins, it was established that hemoglobin A (normal) contains a GLU residue in the locus where hemoglobin S (sickle cell) contains VAL. This replacement of two charged GLU residues for two uncharged VAL residues in the hemoglobin tetramer is sufficient to account for the "sickling" phenomenon in the abnormal hemoglobin. Listed below are a number of known amino acid replacements in abnormal human hemoglobins.

HEMOGLOBIN	CHANGES			REFERENCE
NAME	CHAIN	POS.	FROM TO	
A NORMAL				
F NORMAL FETAL	BETA (CALLED GAMMA)			1
I	ALPHA	16	LYS-ASP	2
NORFOLK	ALPHA	57	GLY-ASP	3
M BOSTON	ALPHA	58	HIS-TYR	4
M SASKATOON	BETA	63	HIS-TYR	4
M MILWAUKEE	BETA	67	VAL-GLU	4
D PUNJAB	BETA	121	GLU-GLN	5
G SAN JOSE	BETA	7	GLU-GLY	6
ZURICH	BETA	63	HIS-ARG	7
C	BETA	6	GLU-LYS	8
O ARABIA	BETA	121	GLU-LYS	9
O INDONESIA	ALPHA	116	GLU-LYS	9
X	ALPHA and BETA	68 6	ASN-LYS GLU-LYS	10
S	BETA	6	GLU-VAL	11
D IBADAN	BETA	87	THR-LYS	12
F TEXAS	GAMMA	5 or 6	GLU-LYS	13
KENWOOD	BETA	143	HIS-ASP	14
G	BETA	7	GLU-GLY	15

1. Rhinesmith, H. W., Schroeder, W. A., and Pauling, L., J. Am. Chem. Soc., Vol. 79, p. 4682, 1957

   Rhinesmith, H. W., Schroeder, W. A., and Martin, N., J. Am. Chem. Soc., Vol. 80, p. 3358, 1958

2. Murayama, M., Fed. Proc., Vol. 19, p. 78, 1960

3. Baglioni, C., J. Biol. Chem., Vol. 237, pp. 69-74, 1962

4. Gerald, P. S. and Efron, M. L., Proc. Natl. Acad. Sci. U.S., Vol. 47, pp. 1758-1767, 1958

5. Baglioni, C., Biochim. Biophys. Acta, Vol. 59, pp. 437-440, 1962

6. Hill, R. L. and Schwartz, H. C., Nature, Vol. 184, pp. 641-642, 1959

7. Muller, C. J. and Kingma, S., Biochim. Biophys. Acta, Vol. 50, p. 595, 1961

8. Hunt, J. A. and Ingram, V. M., Nature, Vol. 184, p. 640, 1959

   Ingram, V. M., Nature, Vol. 180, pp. 326-328, 1957

9. Baglioni, C. and Lehmann, H., Nature, Vol. 196, pp. 229-232, 1962

10. Baglioni, C. and Ingram, V. M., Nature, Vol. 189, pp. 465-467, 1961

11. Ingram, V. M., Nature, Vol. 180, pp. 326-328, 1957

12. Watson-Williams, E. J., Nature, Vol. 205, pp. 1273-1276, 1965

13. Schneider, R. G., Science, Vol. 148, pp. 240-242, 1965

14. Beale, D. and Lehmann, H., Nature, Vol. 207, pp. 249-261, 1965

15. Hill, R. L., Swenson, R. T., and Schwartz, H. C., J. Biol. Chem., Vol. 235, pp. 3182-3187, 1960

MYOGLOBIN - WHALE

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 V L S E G E W Q L V L H V W A K V E A D V A G H G Q D I L I
 31 R L F K S H P E T L E K F D R F K H L K T E A E M K A S E D
 61 L K K H G V T V L T A L G A I L K K K G H H E A E L K P L A
 91 Q S H A T K H K I P I K O L E F I S E A I I H V L H S R H P
121 G N F G A D A Q G A M N K A L E L F R K D I A A K O K E L G
151 O Q G *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 VAL LEU SER GLU GLY GLU TRP GLN LEU VAL LEU HIS VAL TRP ALA
    LYS VAL GLU ALA ASP VAL ALA GLY HIS GLY GLN ASP ILU LEU ILU
 31 ARG LEU PHE LYS SER HIS PRO GLU THR LEU GLU LYS PHE ASP ARG
    PHE LYS HIS LEU LYS THR GLU ALA GLU MET LYS ALA SER GLU ASP
 61 LEU LYS LYS HIS GLY VAL THR VAL LEU THR ALA LEU GLY ALA ILU
    LEU LYS LYS LYS GLY HIS HIS GLU ALA GLU LEU LYS PRO LEU ALA
 91 GLN SER HIS ALA THR LYS HIS LYS ILU PRO ILU LYS TYR LEU GLU
    PHE ILU SER GLU ALA ILU ILU HIS VAL LEU HIS SER ARG HIS PRO
121 GLY ASN PHE GLY ALA ASP ALA GLN GLY ALA MET ASN LYS ALA LEU
    GLU LEU PHE ARG LYS ASP ILU ALA ALA LYS TYR LYS GLU LEU GLY
151 TYR GLN GLY ***
```

COMPOSITION

17	ALA	A	5	GLN	Q	18	LEU	L	6	SER	S
4	ARG	R	14	GLU	E	19	LYS	K	5	THR	T
2	ASN	N	11	GLY	G	2	MET	M	2	TRP	W
6	ASP	D	12	HIS	H	6	PHE	F	3	TYR	O
0	CYS	C	9	ILU	I	4	PRO	P	8	VAL	V

TOTAL NO. OF ACIDS = 153

* EDMUNDSON, A. B., NATURE, VOL.205, NO.4974, PP.883-887, FEBRUARY 27, 1965

DIHEME PEPTIDE - CHROMATIUM

THE PEPTIDE CONTAINS TWO HEME GROUPS. THE FIRST IS COVALENTLY BONDED TO CYSTEINES 5 AND 8. THERE IS ONLY ONE OTHER CYSTEINE AVAILABLE FOR THE OBSERVED COVALENT BONDING OF THE SECOND HEME.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7
/ F A G K C S Q C H T L V A D E G S A K C H T F D E G S /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
/// PHE ALA GLY LYS CYS SER GLN CYS HIS THR LEU VAL ALA ASP GLU
  GLY SER ALA LYS CYS HIS THR PHE ASP GLU GLY SER ///
```

COMPOSITION

3	ALA	A	1	GLN	Q	1	LEU	L	3	SER	S
0	ARG	R	2	GLU	E	2	LYS	K	2	THR	T
0	ASN	N	3	GLY	G	0	MET	M	0	TRP	W
2	ASP	D	2	HIS	H	2	PHE	F	0	TYR	O
3	CYS	C	0	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS IN FRAGMENT = 27

* DUS,K., BARTSCH, R.G., AND KAMEN,M.D., J. BIOL. CHEM., VOL.237, NO.10, PP.3083-3093, OCT., 1962

FERREDOXIN - CLOSTRIDIUM PASTEURIANUM

THE PROTEIN CONTAINS 7 SULPHIDE AND 7 IRON ATOMS PER MOLECULE.
IT DOES NOT CONTAIN HEME.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 A O K I A D S C V S C/G A C/A S E C P V N A I S Q G D S I F/
31 V I D A D T C I D C G N C A N V C P V G A P V Q E *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 ALA TYR LYS ILU ALA ASP SER CYS VAL SER CYS/GLY ALA CYS/ALA
   SER GLU CYS PRO VAL ASN ALA ILU SER GLN GLY ASP SER ILU PHE/
31 VAL ILU ASP ALA ASP THR CYS ILU ASP CYS GLY ASN CYS ALA ASN
   VAL CYS PRO VAL GLY ALA PRO VAL GLN GLU ***
```

COMPOSITION

8	ALA	A	2	GLN	Q	0	LEU	L	5	SER	S
0	ARG	R	2	GLU	E	1	LYS	K	1	THR	T
3	ASN	N	4	GLY	G	0	MET	M	0	TRP	W
5	ASP	D	0	HIS	H	1	PHE	F	1	TYR	O
8	CYS	C	5	ILU	I	3	PRO	P	6	VAL	V

TOTAL NO. OF ACIDS = 55

* TANAKA,M.,NAKASHIMA,T.,BENSON,A.,MOWER,H.F.,AND YASUNOBU,K.T.,
BIOCHEM. BIOPHYS. RES. COMMUN., VOL. 16, NO.5,
PP.422-427, 1964

AZURIN - PSEUDOMONAS FLUORESCENS

THE BLUE PROTEIN CONTAINS ONE COPPER ATOM PER MOLECULE.

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 A E C S V D I Q G N D Q M Q F N T N A I T V D K S C K Q F T
 31 V N L S H P G N L P K N V M G H N W V L S T A A D M Q G V V
 61 T D G M A S G L D K D O L K P D D S R V I A H T K L I G S G
 91 E K D S V T F D V S K L K E G E Q O M F F C T F P G H S A L
121 M K G T L T L K *
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 ALA GLU CYS SER VAL ASP ILU GLN GLY ASN ASP GLN MET GLN PHE
    ASN THR ASN ALA ILU THR VAL ASP LYS SER CYS LYS GLN PHE THR
 31 VAL ASN LEU SER HIS PRO GLY ASN LEU PRO LYS ASN VAL MET GLY
    HIS ASN TRP VAL LEU SER THR ALA ALA ASP MET GLN GLY VAL VAL
 61 THR ASP GLY MET ALA SER GLY LEU ASP LYS ASP TYR LEU LYS PRO
    ASP ASP SER ARG VAL ILU ALA HIS THR LYS LEU ILU GLY SER GLY
 91 GLU LYS ASP SER VAL THR PHE ASP VAL SER LYS LEU LYS GLU GLY
    GLU GLN TYR MET PHE PHE CYS THR PHE PRO GLY HIS SER ALA LEU
121 MET LYS GLY THR LEU THR LEU LYS ***
```

COMPOSITION

7	ALA	A	6	GLN	Q	10	LEU	L	10	SER	S
1	ARG	R	4	GLU	E	11	LYS	K	10	THR	T
7	ASN	N	11	GLY	G	6	MET	M	1	TRP	W
11	ASP	D	4	HIS	H	6	PHE	F	2	TYR	O
3	CYS	C	4	ILU	I	4	PRO	P	10	VAL	V

TOTAL NO. OF ACIDS = 128

* AMBLER,R.P.,AND BROWN,L.H., J. MOL. BIOL., VOL. 9, NO.3, PP. 825-828, SEPT., 1964

RIBONUCLEASE - BOVINE

DISULPHIDE BONDS ARE FORMED BETWEEN CYSTEINES AT POSITIONS 26 AND 84, 40 AND 95, 58 AND 110, AND 65 AND 72.

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 K E T A A A K F E R Q H M D S S T S A A S S S N D C N Q M M
 31 K S R N L T K D R C K P V N T F V H E S L A D V Q A V C S Q
 61 K N V A C K N G Q T N C D Q S D S T M S I T D C R E T G S S
 91 K D P N C A D K T T Q A N K H I I V A C E G N P D V P V H F
121 D A S V *
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 LYS GLU THR ALA ALA ALA LYS PHE GLU ARG GLN HIS MET ASP SER
    SER THR SER ALA ALA SER SER SER ASN TYR CYS ASN GLN MET MET
 31 LYS SER ARG ASN LEU THR LYS ASP ARG CYS LYS PRO VAL ASN THR
    PHE VAL HIS GLU SER LEU ALA ASP VAL GLN ALA VAL CYS SER GLN
 61 LYS ASN VAL ALA CYS LYS ASN GLY GLN THR ASN CYS TYR GLN SER
    TYR SER THR MET SER ILU THR ASP CYS ARG GLU THR GLY SER SER
 91 LYS TYR PRO ASN CYS ALA TYR LYS THR THR GLN ALA ASN LYS HIS
    ILU ILU VAL ALA CYS GLU GLY ASN PRO TYR VAL PRO VAL HIS PHE
121 ASP ALA SER VAL ***
```

COMPOSITION

12	ALA	A	7	GLN	Q	2	LEU	L	15	SER	S
4	ARG	R	5	GLU	E	10	LYS	K	10	THR	T
10	ASN	N	3	GLY	G	4	MET	M	0	TRP	W
5	ASP	D	4	HIS	H	3	PHE	F	6	TYR	O
8	CYS	C	3	ILU	I	4	PRO	P	9	VAL	V

TOTAL NO. OF ACIDS = 124

* SMYTH, D.G., STEIN,W.H. AND MOORE, S., J. BIOL. CHEM., VOL.238, NO.1, PP.227-234, JAN., 1963

TRYPSIN INHIBITOR - BOVINE

DISULPHIDE BONDS ARE FORMED BETWEEN CYSTEINES AT POSITIONS 5-55, 14-38, AND 30-51.

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 R P D F C L E P P O T G P C K A R I I R O F O N A K A G L C
31 Q T F V O G G C R A K R N N F K S A E D C M R T C G G A *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 ARG PRO ASP PHE CYS LEU GLU PRO PRO TYR THR GLY PRO CYS LYS
   ALA ARG ILU ILU ARG TYR PHE TYR ASN ALA LYS ALA GLY LEU CYS
31 GLN THR PHE VAL TYR GLY GLY CYS ARG ALA LYS ARG ASN ASN PHE
   LYS SER ALA GLU ASP CYS MET ARG THR CYS GLY GLY ALA ***
```

COMPOSITION

6	ALA	A	1	GLN	Q	2	LEU	L	1	SER	S
6	ARG	R	2	GLU	E	4	LYS	K	3	THR	T
3	ASN	N	6	GLY	G	1	MET	M	0	TRP	W
2	ASP	D	0	HIS	H	4	PHE	F	4	TYR	O
6	CYS	C	2	ILU	I	4	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 58

- KASSELL, B., RADICEVIC, M., ANSFIELD,M. J., AND LASKOWSKI, M., BIOCHEM. BIOPHYS. RES. COMMUN., VOL. 18, NO,2, PP.255-258, 1965

DLOUHA, V., POSPISILOVA, D., MELOUN, B. AND SORM, F., COLLECTION CZECH. CHEM. COMMUN., VOL. 30, PP.1311-1325, 1965

THE SEQUENCE REPORTED HERE DIFFERS FROM THE ABOVE IN HAVING THE ILU (I) DELETED AT POSITION 19.

CHAUVET, J., NOUVEL, G., AND ACHER, R., BIOCHIM. BIOPHYS. ACTA , VOL. 92, PP. 200-201, 1964

THE SEQUENCE REPORTED HERE DIFFERS FROM THE ABOVE IN THE FOLLOWING RESPECTS.
THE ARG (R) FROM POSITION 42 HAS BEEN REMOVED AND INSERTED BETWEEN POSITIONS 20 AND 21. THE GLN (Q) AT POSITION 31 HAS BEEN DELETED AND A GLU (E) ADDED BETWEEN POSITIONS 32 AND 33.

KASSELL, B., AND LASKOWSKI, M., BIOCHEM. BIOPHYS. RES. COMMUN. VOL 20, NO.4, PP.463-468, 1965

TOBACCO MOSAIC VIRUS

ACETYL - AT AMINO END

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 S O S I T T P S Q F V F L S S A W A D P I E L I N L C T N A
 31 L G N Q F Q T Q Q A R T V Q V R Q F S Q V W K P S P Q V T V
 61 R F P D S D F K V O R O N A V L D P L V T A L L G A F D T R
 91 N R I I Q V Q D Q A N P T T A Q T L D A T R R V D D A T V A
121 I R S A D I N L I V E L I R G T G S Q N R S S F E S S S G L
151 V W T S G P A T *
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 SER TYR SER ILU THR THR PRO SER GLN PHE VAL PHE LEU SER SER
    ALA TRP ALA ASP PRO ILU GLU LEU ILU ASN LEU CYS THR ASN ALA
 31 LEU GLY ASN GLN PHE GLN THR GLN GLN ALA ARG THR VAL GLN VAL
    ARG GLN PHE SER GLN VAL TRP LYS PRO SER PRO GLN VAL THR VAL
 61 ARG PHE PRO ASP SER ASP PHE LYS VAL TYR ARG TYR ASN ALA VAL
    LEU ASP PRO LEU VAL THR ALA LEU LEU GLY ALA PHE ASP THR ARG
 91 ASN ARG ILU ILU GLN VAL GLN ASP GLN ALA ASN PRO THR THR ALA
    GLN THR LEU ASP ALA THR ARG ARG VAL ASP ASP ALA THR VAL ALA
121 ILU ARG SER ALA ASP ILU ASN LEU ILU VAL GLU LEU ILU ARG GLY
    THR GLY SER TYR ASN ARG SER SER PHE GLU SER SER SER GLY LEU
151 VAL TRP THR SER GLY PRO ALA THR ***
```

COMPOSITION

14	ALA	A	13	GLN	Q	12	LEU	L	16	SER	S
11	ARG	R	3	GLU	E	2	LYS	K	16	THR	T
8	ASN	N	6	GLY	G	0	MET	M	3	TRP	W
10	ASP	D	0	HIS	H	8	PHE	F	4	TYR	O
1	CYS	C	9	ILU	I	8	PRO	P	14	VAL	V

TOTAL NO. OF ACIDS = 158

• ANDERER, F.A., Z. NATURFORSCH., VOL. 17, PP.526-543, 1962

STRUCTURE REVISIONS AND CONFIRMATIONS.

ANDERER, F.A., UHLIG, H., WEBER, E., AND SCHRAMM, G., NATURE, VOL. 186, NO.4729, PP.922-925, JUNE 18, 1960

FUNATSU, G., TSUGITA, A., AND FRAENKEL-CONRAT, H., ARCH. BIOCHEM. BIOPHYS., VOL. 105, NO.1, PP.25-41, APR. 1964

TOBACCO MOSAIC VIRUS STRAIN DAHLMENSE

ACETYL- AT AMINO END

```
     1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
   1 S O S I T S P S Q F V F L S S V W A D P I E L L N V C T S S
  31 L G N Q F Q T Q Q A R T T Q V Q Q F S E V W K P F P Q S T V
  61 R F P G D V O K V O R O N A V L D P L I T A L L G T F D T R
  91 N R I I E V E N Q Q S P T T A E T L D A T R R V D D A T V A
 121 I R S A N I N L V N E L V R G T G L O N Q N T F E S M S G L
 151 V W T S A P A S •
```

```
      1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
   1 SER TYR SER ILU THR SER PRO SER GLN PHE VAL PHE LEU SER SER
     VAL TRP ALA ASP PRO ILU GLU LEU LEU ASN VAL CYS THR SER SER
  31 LEU GLY ASN GLN PHE GLN THR GLN GLN ALA ARG THR THR GLN VAL
     GLN GLN PHE SER GLU VAL TRP LYS PRO PHE PRO GLN SER THR VAL
  61 ARG PHE PRO GLY ASP VAL TYR LYS VAL TYR ARG TYR ASN ALA VAL
     LEU ASP PRO LEU ILU THR ALA LEU LEU GLY THR PHE ASP THR ARG
  91 ASN ARG ILU ILU GLU VAL GLU ASN GLN GLN SER PRO THR THR ALA
     GLU THR LEU ASP ALA THR ARG ARG VAL ASP ASP ALA THR VAL ALA
 121 ILU ARG SER ALA ASN ILU ASN LEU VAL ASN GLU LEU VAL ARG GLY
     THR GLY LEU TYR ASN GLN ASN THR PHE GLU SER MET SER GLY LEU
 151 VAL TRP THR SER ALA PRO ALA SER ***
```

COMPOSITION

11	ALA	A	12	GLN	Q	13	LEU	L	16	SER	S
9	ARG	R	7	GLU	E	2	LYS	K	17	THR	T
10	ASN	N	6	GLY	G	1	MET	M	3	TRP	W
7	ASP	D	0	HIS	H	8	PHE	F	5	TYR	O
1	CYS	C	7	ILU	I	8	PRO	P	15	VAL	V

TOTAL NO. OF ACIDS = 158

• WITTMANN-LIEBOLD,B. AND WITTMANN, H. G., Z. VERERBUNGS.,
VOL. 94, PP. 427-435, 1963

CHYMOTRYPSINOGEN-A - BOVINE

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 C G V P A I Q P V L S G L S R I V G D E E A V P G S W P W Q
 31 V S L Q D K T G F H F C G G S L I N E N W V V T A A H C G V
 61 T T S D V V V A G E F D Q G S S S E K I Q K L K I A K V F K
 91 N S K O N S L T I N N N I T L L K L S T A A S F S Q T V S A
121 V C L P S A S D D F A A G T T C V T T G W G L T R O T N A N
151 T P D R L Q Q A S L P L L S N T N C K K O W G T K I K D A M
181 I C A G A S G V S S C M G D S G G P L V C K K N G A W T L V
211 G I V S W G S S T C S T S T P G V O A R V T A L V N W V Q Q
241 T L A A N •
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 CYS GLY VAL PRO ALA ILU GLN PRO VAL LEU SER GLY LEU SER ARG
    ILU VAL GLY ASP GLU GLU ALA VAL PRO GLY SER TRP PRO TRP GLN
 31 VAL SER LEU GLN ASP LYS THR GLY PHE HIS PHE CYS GLY GLY SER
    LEU ILU ASN GLU ASN TRP VAL VAL THR ALA ALA HIS CYS GLY VAL
 61 THR THR SER ASP VAL VAL VAL ALA GLY GLU PHE ASP GLN GLY SER
    SER SER GLU LYS ILU GLN LYS LEU LYS ILU ALA LYS VAL PHE LYS
 91 ASN SER LYS TYR ASN SER LEU THR ILU ASN ASN ASN ILU THR LEU
    LEU LYS LEU SER THR ALA ALA SER PHE SER GLN THR VAL SER ALA
121 VAL CYS LEU PRO SER ALA SER ASP ASP PHE ALA ALA GLY THR THR
    CYS VAL THR THR GLY TRP GLY LEU THR ARG TYR THR ASN ALA ASN
151 THR PRO ASP ARG LEU GLN GLN ALA SER LEU PRO LEU LEU SER ASN
    THR ASN CYS LYS LYS TYR TRP GLY THR LYS ILU LYS ASP ALA MET
181 ILU CYS ALA GLY ALA SER GLY VAL SER SER CYS MET GLY ASP SER
    GLY GLY PRO LEU VAL CYS LYS LYS ASN GLY ALA TRP THR LEU VAL
211 GLY ILU VAL SER TRP GLY SER SER THR CYS SER THR SER THR PRO
    GLY VAL TYR ALA ARG VAL THR ALA LEU VAL ASN TRP VAL GLN GLN
241 THR LEU ALA ALA ASN ***
```

COMPOSITION

22	ALA	A	10	GLN	Q	19	LEU	L	28	SER	S
4	ARG	R	5	GLU	E	14	LYS	K	23	THR	T
14	ASN	N	23	GLY	G	2	MET	M	8	TRP	W
9	ASP	D	2	HIS	H	6	PHE	F	4	TYR	O
10	CYS	C	10	ILU	I	9	PRO	P	23	VAL	V

TOTAL NO. OF ACIDS = 245

• HARTLEY, B.S., BROWN, J.R., KAUFFMAN, D.L., AND SMILLIE,L.B., NATURE, VOL.207, NO.5002, PP.1157-1159, SEPT.11, 1965

THIS SEQUENCE HAS BEEN CORRECTED BY DELETING SER (S) WHICH WAS AT POSITION 215.

BROWN, J.R., AND HARTLEY, B. S., BIOCHEM J., VOL. 89, 59P, 1963

THE ACTIVE SITE SERINE IS AT POSITION 195

KEIL, B., PRUSIK, Z., AND SORM, F., BIOCHIM. BIOPHYS. ACTA, VOL. 78, P. 559-561, 1963

DISULPHIDE BRIDGES LINK POSITIONS 1-122, 42-58, 136-201, 168-182 AND 191-220.

KOSTKA, V., MELOUN, B., AND SORM,F., COLLECTION CZECH. CHEM. COMMUN., VOL. 28, PP.2779-2805, 1963

HARTLEY,B.S., NATURE, VOL. 201, NO. 4962, PP.1284-1287, MARCH 28,1964

TRYPSINOGEN

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 V D D D D K I V G G O T C G A N T V P O Q V S L N S G O H F
 31 C G G S L I N S Q W V V S A A H C O K S G I Q V R L G E D N
 61 I N V V E G D E Q F I S A S K S I V H P S O N(P,L,T,N)N N D
 91 I M L I K L K S A A S L N S R V A S I S L P T S C A S A G T
121 Q C L I S G W G N T K S S G T S O P D V L K C L K A P I L S
151 D S S C K S A O P G Q I T S N M F C A G O L E G G K N S C Q
181 G D S G G P V V C S G K L Q G I V S W G S G C A Q K N K P G
211 V O T K V C N O V S W I K Q T I A S N *
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 VAL ASP ASP ASP ASP LYS ILU VAL GLY GLY TYR THR CYS GLY ALA
    ASN THR VAL PRO TYR GLN VAL SER LEU ASN SER GLY TYR HIS PHE
 31 CYS GLY GLY SER LEU ILU ASN SER GLN TRP VAL VAL SER ALA ALA
    HIS CYS TYR LYS SER GLY ILU GLN VAL ARG LEU GLY GLU ASP ASN
 61 ILU ASN VAL VAL GLU GLY ASP GLU GLN PHE ILU SER ALA SER LYS
    SER ILU VAL HIS PRO SER TYR ASN(PRO,LEU,THR,ASN)ASN ASN ASP
 91 ILU MET LEU ILU LYS LEU LYS SER ALA ALA SER LEU ASN SER ARG
    VAL ALA SER ILU SER LEU PRO THR SER CYS ALA SER ALA GLY THR
121 GLN CYS LEU ILU SER GLY TRP GLY ASN THR LYS SER SER GLY THR
    SER TYR PRO ASP VAL LEU LYS CYS LEU LYS ALA PRO ILU LEU SER
151 ASP SER SER CYS LYS SER ALA TYR PRO GLY GLN ILU THR SER ASN
    MET PHE CYS ALA GLY TYR LEU GLU GLY GLY LYS ASN SER CYS GLN
181 GLY ASP SER GLY GLY PRO VAL VAL CYS SER GLY LYS LEU GLN GLY
    ILU VAL SER TRP GLY SER GLY CYS ALA GLN LYS ASN LYS PRO GLY
211 VAL TYR THR LYS VAL CYS ASN TYR VAL SER TRP ILU LYS GLN THR
    ILU ALA SER ASN ***
```

COMPOSITION

14	ALA	A	10	GLN	Q	14	LEU	L	33	SER	S
2	ARG	R	4	GLU	E	15	LYS	K	10	THR	T
16	ASN	N	25	GLY	G	2	MET	M	4	TRP	W
10	ASP	D	3	HIS	H	3	PHE	F	10	TYR	O
12	CYS	C	15	ILU	I	9	PRO	P	18	VAL	V

TOTAL NO. OF ACIDS = 229

• WALSH, K., AND NEURATH, H., PROC. NATL. ACAD. SCI. U.S., VOL. 52, NO.4, PP.884-889, 1964

KAUFFMAN, D. L., J. MOL. BIOL., VOL.12, PP.929-932, 1965

DISULPHIDE BRIDGES WERE FOUND BETWEEN LINKS 13-143, 31-47
115-216, 122-189, 154-168, AND 179-203.
THE ACTIVE SERINE IS AT LINK 183.

PAPAIN

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1/I P E O V D W R Q K G A V T P V K N Q G S C G S C W/A F/I I/
 31 R N T P O O E G V Q R O C R S R E K G P O A A K T D G V R Q
 61 V Q P O N Q G A L L O S I A N Q P S V V L Q A A G K D F Q L
 91 O R G G I F V G P C G N K V D H A V A A V G O N P G O I L I
121 K N S W G T G W G E N G O I R I K T G N L N Q O S E Q E L L
151 D C D R R S O G C O P G D G W/S A L/V A Q O G I H O R G T G
181 N S O G V C G L O T S S F O P V K N •
```

```
     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1/ILU PRO GLU TYR VAL ASP TRP ARG GLN LYS GLY ALA VAL THR PRO
    VAL LYS ASN GLN GLY SER CYS GLY SER CYS TRP/ALA PHE/ILU ILU/
 31 ARG ASN THR PRO TYR TYR GLU GLY VAL GLN ARG TYR CYS ARG SER
    ARG GLU LYS GLY PRO TYR ALA ALA LYS THR ASP GLY VAL ARG GLN
 61 VAL GLN PRO TYR ASN GLN GLY ALA LEU LEU TYR SER ILU ALA ASN
    GLN PRO SER VAL VAL LEU GLN ALA ALA GLY LYS ASP PHE GLN LEU
 91 TYR ARG GLY GLY ILU PHE VAL GLY PRO CYS GLY ASN LYS VAL ASP
    HIS ALA VAL ALA ALA VAL GLY TYR ASN PRO GLY TYR ILU LEU ILU
121 LYS ASN SER TRP GLY THR GLY TRP GLY GLU ASN GLY TYR ILU ARG
    ILU LYS THR GLY ASN LEU ASN GLN TYR SER GLU GLN GLU LEU LEU
151 ASP CYS ASP ARG ARG SER TYR GLY CYS TYR PRO GLY ASP GLY TRP/
    SER ALA LEU/VAL ALA GLN TYR GLY ILU HIS TYR ARG GLY THR GLY
181 ASN SER TYR GLY VAL CYS GLY LEU TYR THR SER SER PHE TYR PRO
    VAL LYS ASN ***
```

COMPOSITION

13	ALA	A	12	GLN	Q	10	LEU	L	12	SER	S
11	ARG	R	6	GLU	E	9	LYS	K	7	THR	T
12	ASN	N	27	GLY	G	0	MET	M	5	TRP	W
7	ASP	D	2	HIS	H	4	PHE	F	19	TYR	O
7	CYS	C	10	ILU	I	10	PRO	P	15	VAL	V

TOTAL NO. OF ACIDS = 198

• LIGHT, A., FRATER, R., KIMMEL, J., AND SMITH, E.L., PROC. NATL. ACAD. SCI. U.S., VOL.52, NO.5, PP.1276-1283, NOV. 1964

DISULPHIDE BRIDGES ARE FORMED BETWEEN CYSTEINES AT POSITIONS 43 AND 152, 100 AND 186, AND 22 AND 159.

THE ACTIVE SULFHYDRYL GROUP IS AT POSITION 25.

LYSOZYME - CHICKEN

LYSOZYME HAS A BETA (1-4) GLUCOSAMINIDASE ACTIVITY WITH THE ABILITY TO HYDROLYSE A MUCOPOLYSACCHARIDE COMPONENT OF SOME BACTERIAL CELL WALLS RELEASING N-ACETYL AMINO SUGARS DERIVED FROM GLUCOSAMINE AND MURAMIC ACID.

	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	6	7	8	9	0
1	K	V	F	G	R	C	E	L	A	A	A	M	K	R	H	G	L	D	N	O	R	G	O	S	L	G	N	W	V	C
31	A	A	K	F	E	S	N	F	N	T	Q	A	T	N	R	N	T	D	G	S	T	D	O	G	I	L	Q	I	N	S
61	R	W	W	C	N	D	G	R	T	P	G	S	R	N	L	C	N	I	P	C	S	A	L	L	S	S	D	I	T	A
91	S	V	N	C	A	K	K	I	V	S	D	G	D	G	M	N	A	W	V	A	W	R	N	R	C	K	G	T	D	V
121	Q	A	W	I	R	G	C	R	L	*																				

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	LYS	VAL	PHE	GLY	ARG	CYS	GLU	LEU	ALA	ALA	ALA	MET	LYS	ARG	HIS
	GLY	LEU	ASP	ASN	TYR	ARG	GLY	TYR	SER	LEU	GLY	ASN	TRP	VAL	CYS
31	ALA	ALA	LYS	PHE	GLU	SER	ASN	PHE	ASN	THR	GLN	ALA	THR	ASN	ARG
	ASN	THR	ASP	GLY	SER	THR	ASP	TYR	GLY	ILU	LEU	GLN	ILU	ASN	SER
61	ARG	TRP	TRP	CYS	ASN	ASP	GLY	ARG	THR	PRO	GLY	SER	ARG	ASN	LEU
	CYS	ASN	ILU	PRO	CYS	SER	ALA	LEU	LEU	SER	SER	ASP	ILU	THR	ALA
91	SER	VAL	ASN	CYS	ALA	LYS	LYS	ILU	VAL	SER	ASP	GLY	ASP	GLY	MET
	ASN	ALA	TRP	VAL	ALA	TRP	ARG	ASN	ARG	CYS	LYS	GLY	THR	ASP	VAL
121	GLN	ALA	TRP	ILU	ARG	GLY	CYS	ARG	LEU	***					

COMPOSITION

12	ALA	A	3	GLN	Q	8	LEU	L	10	SER	S
11	ARG	R	2	GLU	E	6	LYS	K	7	THR	T
13	ASN	N	12	GLY	G	2	MET	M	6	TRP	W
8	ASP	D	1	HIS	H	3	PHE	F	3	TYR	O
8	CYS	C	6	ILU	I	2	PRO	P	6	VAL	V

TOTAL NO. OF ACIDS = 129

• CANFIELD, R., J. BIOL. CHEM., VOL.238, NO.8, PP.2698-2707, AUG., 1963

CANFIELD, R., LIU,A.K., J. BIOL. CHEM., VOL.240, NO.5, PP. 1997-2002, MAY 1965

ABOVE SEQUENCE CONFIRMED IN THIS WORK.
DISULPHIDE BONDS ARE FOUND BETWEEN 6 AND 127, 30 AND 115, 64 AND 80, AND 76 AND 94.

JOLLES, J., JAUREGUI-ADELL, J., BERNIER, I., AND JOLLES, P., BIOCHIM. BIOPHYS. ACTA, VOL.78, PP.668-689, 1963

THIS SEQUENCE DIFFERS FROM THE ABOVE AS FOLLOWS, 40-GLN, 41-ALA, 42-THR, 43-THR, 46-ASP, 58-ASN, 59-ILU, 92-ASN, AND 93-VAL.

BLAKE, C.C.F., KOENING, D.F., MAIR, G.A., NORTH, A.C.T., PHILLIPS, D.C., AND SARMA, V.R., NATURE, NO. 4986, PP. 757-761, MAY 22, 1965

A 2 ANGSTROM RESOLUTION FOURIER SYNTHESIS HAS BEEN PERFORMED BY X-RAY CRYSTALLOGRAPHIC METHODS. THE LOCATION OF THE FOUR DISULPHIDE BRIDGES HAS BEEN CONFIRMED. THE BINDING SITE OF THE INHIBITOR N-ACETYL-GLUCOSAMINE AND ITS DIMER HAS BEEN FOUND TO BE VERY EXTENSIVE INVOLVING RESIDUES AT POSITIONS 44, 46, 47, 48, 50, 52, 57, 59, 61-63, 72, 73, 97, 99-101, 103, 107-110, 113, AND 114.

GLUCAGON - BOVINE

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 H S Q G T F T S D O S K O L D S R R A Q D F V Q W L M N T *
```

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	HIS	SER	GLN	GLY	THR	PHE	THR	SER	ASP	TYR	SER	LYS	TYR	LEU	ASP
	SER	ARG	ARG	ALA	GLN	ASP	PHE	VAL	GLN	TRP	LEU	MET	ASN	THR	***

COMPOSITION

1	ALA	A	3	GLN	Q	2	LEU	L	4	SER	S
2	ARG	R	0	GLU	E	1	LYS	K	3	THR	T
1	ASN	N	1	GLY	G	1	MET	M	1	TRP	W
3	ASP	D	1	HIS	H	2	PHE	F	2	TYR	O
0	CYS	C	0	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 29

• BROMER, W.W., SINN, L.G., AND BEHRENS, O.K., J. AM. CHEM. SOC., VOL. 79, PP. 2807-2810, JUNE 5, 1957

ARGININE VASOPRESSIN - BOVINE

THE C TERMINAL GLYCINE IS PRESENT AS THE AMIDE.
THE TWO CYSTEINES ARE LINKED BY A DISULPHIDE BOND.

```
  1 2 3 4 5 6 7 8 9
1 C O F Q N C P R G *

  1   2   3   4   5   6   7   8   9
1 CYS TYR PHE GLN ASN CYS PRO ARG GLY ***
```

COMPOSITION

0	ALA	A	1	GLN	Q	0	LEU	L	0	SER	S
1	ARG	R	0	GLU	E	0	LYS	K	0	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	1	PHE	F	1	TYR	O
2	CYS	C	0	ILU	I	1	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 9

* DU VIGNEAUD, V., LAWLER, H. C., AND POPENOE, E. A., J. AM. CHEM. SOC., VOL.75, PP.4880-4881, OCT. 5, 1953

ACHER, R., AND CHAUVET, J., BIOCHIM. BIOPHYS. ACTA, VOL. 12, PP.487-488, 1953

THIS WORK CONFIRMED THE SEQUENCE ABOVE, HOWEVER GLU (E) AND ASP (D) WERE NOT DISTINGUISHED FROM GLN (Q) AND ASN (N).

LYSINE VASOPRESSIN - PIG

THE C TERMINAL GLYCINE IS PRESENT AS THE AMIDE.
THE TWO CYSTEINES ARE LINKED BY A DISULPHIDE BOND.

```
  1 2 3 4 5 6 7 8 9
1 C.O.F.Q.N.C.P.K.G.*

   1   2   3   4   5   6   7   8   9
1 CYS.TYR.PHE.GLN.ASN.CYS.PRO.LYS.GLY ***
```

COMPOSITION

0	ALA	A	1	GLN	Q	0	LEU	L	0	SER	S
0	ARG	R	0	GLU	E	1	LYS	K	0	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	1	PHE	F	1	TYR	O
2	CYS	C	0	ILU	I	1	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 9

* POPENOE, E. A., LAWLER, H. C., AND DU VIGNEAUD, V.,
J. AM. CHEM. SOC., VOL.74, P.3713, JULY 20, 1952

OXYTOCIN - BOVINE

THE C TERMINAL GLYCINE IS PRESENT AS THE AMIDE.
THE TWO CYSTEINES ARE LINKED BY A DISULPHIDE BOND.
OXYTOCIN IS THE PRINCIPAL UTERINE CONTRACTING AND MILK EJECTING HORMONE OF THE POSTERIOR PITUITARY.

```
  1 2 3 4 5 6 7 8 9
1 C O I Q N C P L G *

   1   2   3   4   5   6   7   8   9
1 CYS TYR ILU GLN ASN CYS PRO LEU GLY ***
```

COMPOSITION

0	ALA	A	1	GLN	Q	1	LEU	L	0	SER	S
0	ARG	R	0	GLU	E	0	LYS	K	0	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	1	TYR	O
2	CYS	C	1	ILU	I	1	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 9

* DU VIGNEAUD, V., RESSLER, C., TRIPPETT, S., J. BIOL. CHEM., VOL.205, PP.949-957, 1953

TUPPY, H. AND MICHL, H., MONATSH. CHEM., VOL.84, PP.1011-1020, 1953

HYPERTENSIN - BOVINE

```
  1 2 3 4 5 6 7 8 9 0
1 D R V O V H P F H L *
```

```
   1   2   3   4   5   6   7   8   9  10
1 ASP ARG VAL TYR VAL HIS PRO PHE HIS LEU ***
```

COMPOSITION

0	ALA	A	0	GLN	Q	1	LEU	L	0	SER	S
1	ARG	R	0	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	0	GLY	G	0	MET	M	0	TRP	W
1	ASP	D	2	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 10

* ELLIOT, D. F., AND PEART, W. S., BIOCHEM. J., VOL.65, PP.246-254, 1957

ALPHA MELANOCYTE-STIMULATING HORMONE - BOVINE, PIG, AND HORSE

ACETYL AT AMINO END.
C-TERMINAL VALINE IS AMINATED.

1	2	3	4	5	6	7	8	9	0	1	2	3
S	O	S	M	E	H	F	R	W	G	K	P	V *

1	2	3	4	5	6	7	8	9	10	11	12	13
SER	TYR	SER	MET	GLU	HIS	PHE	ARG	TRP	GLY	LYS	PRO	VAL ***

COMPOSITION

0	ALA	A	0	GLN	Q	0	LEU	L	2	SER	S
1	ARG	R	1	GLU	E	1	LYS	K	0	THR	T
0	ASN	N	1	GLY	G	1	MET	M	1	TRP	W
0	ASP	D	1	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 13

* HARRIS, J. I. AND LERNER, A. B., NATURE, VOL.179, NO.4574, PP.1346-1347, JUNE 29, 1957 (PIG)

LI, C. H., LABORATORY INVESTIGATION, VOL. 8, NO.2, PP.574-587, 1959 (BOVINE)

DIXON, J. S. AND LI, C. H., J. AM. CHEM. SOC., VOL.82, PP.4568-4572, SEPT. 5, 1960 (HORSE)

BETA MELANOCYTE-STIMULATING HORMONE - BOVINE

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8
1 D S G P O K M E H F R W G S P P K D *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ASP SER GLY PRO TYR LYS MET GLU HIS PHE ARG TRP GLY SER PRO
  PRO LYS ASP ***
```

COMPOSITION

0	ALA	A	0	GLN	Q	0	LEU	L	2	SER	S
1	ARG	R	1	GLU	E	2	LYS	K	0	THR	T
0	ASN	N	2	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	3	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 18

* GESCHWIND, I.I., LI, C. H., AND BARNAFI, L., J. AM. CHEM. SOC., VOL. 79, PP.1003-1004, FEB. 20, 1957

BETA MELANOCYTE-STIMULATING HORMONE - PIG

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8
1 D E G P O K M E H F R W G S P P K D *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ASP GLU GLY PRO TYR LYS MET GLU HIS PHE ARG TRP GLY SER PRO
  PRO LYS ASP ***
```

COMPOSITION

0	ALA	A	0	GLN	Q	0	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	2	LYS	K	0	THR	T
0	ASN	N	2	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	3	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 18

* HARRIS, J. I. AND ROOS, P., NATURE, VOL.178, NO.4524, P. 90, JULY 14, 1956

GESCHWIND, I.I., LI, C. H., AND BARNAFI, L., J. AM. CHEM. SOC., VOL. 79, PP.620-625, FEB. 5, 1957

BETA MELANOCYTE-STIMULATING HORMONE - HORSE

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8
1 D E G P O K M E H F R W G S P R K D *


   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ASP GLU GLY PRO TYR LYS MET GLU HIS PHE ARG TRP GLY SER PRO
  ARG LYS ASP ***
```

COMPOSITION

0	ALA	A	0	GLN	Q	0	LEU	L	1	SER	S
2	ARG	R	2	GLU	E	2	LYS	K	0	THR	T
0	ASN	N	2	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	2	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 18

* DIXON, J. S. AND LI, C. H., GEN. COMP. ENDOCRINOL., VOL.1, PP.161-169, 1961

BETA MELANOCYTE-STIMULATING HORMONE - HUMAN

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2
1 A E K K D E G P O R M E H F R W G S P P K D *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA GLU LYS LYS ASP GLU GLY PRO TYR ARG MET GLU HIS PHE ARG
  TRP GLY SER PRO PRO LYS ASP ***
```

COMPOSITION

1	ALA	A	0	GLN	Q	0	LEU	L	1	SER	S
2	ARG	R	3	GLU	E	3	LYS	K	0	THR	T
0	ASN	N	2	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	3	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 22

• HARRIS, J. I., NATURE, VOL. 184, NO. 4681, PP.167-169, JULY 18, 1959

BETA CORTICOTROPIN - PIG

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 S O S M E H F R W G K P V G K K R R P V K V O P G A E D D Q
31 L A E A F P L E F •
```

```
    1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 SER TYR SER MET GLU HIS PHE ARG TRP GLY LYS PRO VAL GLY LYS
   LYS ARG ARG PRO VAL LYS VAL TYR PRO GLY ALA GLU ASP ASP GLN
31 LEU ALA GLU ALA PHE PRO LEU GLU PHE ***
```

COMPOSITION

3	ALA	A	1	GLN	Q	2	LEU	L	2	SER	S
3	ARG	R	4	GLU	E	4	LYS	K	0	THR	T
0	ASN	N	3	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	3	PHE	F	2	TYR	O
0	CYS	C	0	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 39

• WHITE, W. F., AND LANDMANN, W. A., J. AM. CHEM. SOC., VOL. 77, PP.1711-1712, MARCH 20, 1955

HOWARD, K. S., SHEPHERD, R. G., EIGNER, E. A., DAVIS, D. S., AND BELL, P. H., J. AM. CHEM. SOC., VOL.77, PP.3419-3420, JUNE 20, 1955

BELL, P. H., J. AM. CHEM. SOC., VOL.76, PP.5565-5567, NOV. 1954

THIS SEQUENCE DIFFERS FROM THAT SHOWN ABOVE BY REMOVING THE ASP (D) FROM POSITION 29 AND INSERTING IT BETWEEN POSITIONS 24 AND 25.

ALPHA CORTICOTROPIN - SHEEP AND BOVINE

```
   1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
 1 S O S M E H F R W G K P V G K K R R P V K V O P D G E A E D
31 S A Q A F P L E F •
```

```
   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
 1 SER TYR SER MET GLU HIS PHE ARG TRP GLY LYS PRO VAL GLY LYS
   LYS ARG ARG PRO VAL LYS VAL TYR PRO ASP GLY GLU ALA GLU ASP
31 SER ALA GLN ALA PHE PRO LEU GLU PHE ***
```

COMPOSITION

3	ALA	A	1	GLN	Q	1	LEU	L	3	SER	S
3	ARG	R	4	GLU	E	4	LYS	K	0	THR	T
0	ASN	N	3	GLY	G	1	MET	M	1	TRP	W
2	ASP	D	1	HIS	H	3	PHE	F	2	TYR	O
0	CYS	C	0	ILU	I	4	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 39

• LI, C.H., GESCHWIND, I. I., COLE, D., RAACK, I. D., HARRIS, J.I., AND DIXON, J. S., NATURE, VOL.176, NO.4484, PP.687-689, OCT. 8, 1955 (SHEEP)

LI, C. H., DIXON, J. S., AND CHUNG, D., J. AM. CHEM. SOC., VOL. 80, P.2587, 1958 (BOVINE)

INSULIN A - BOVINE

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I V E Q C C A S V C S L O Q L E N O C N *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU VAL GLU GLN CYS CYS ALA SER VAL CYS SER LEU TYR GLN
  LEU GLU ASN TYR CYS ASN ***
```

COMPOSITION

1	ALA	A	2	GLN	Q	2	LEU	L	2	SER	S
0	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
2	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
4	CYS	C	1	ILU	I	0	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 21

* SANGER, F. AND THOMPSON, E. O. P., BIOCHEM J., VOL.53, PP. 353-374, 1953

THE AMIDE GROUPS WERE SUBSEQUENTLY DETERMINED.

RYLE, A. P., SANGER, F., SMITH, L.F., AND KITAI, R., BIOCHEM. J., VOL. 60, PP. 541-556, 1955

INSULIN A - BONITO

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I(H,E,E,C(C,K,P,H)C,D,L)F E L E D D C N *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU(HIS,GLU,GLU,CYS(CYS,LYS,PRO,HIS)CYS,ASP,LEU)PHE GLU
  LEU GLU ASP TYR CYS ASN ***
```

COMPOSITION

0	ALA	A	0	GLN	Q	2	LEU	L	0	SER	S
0	ARG	R	4	GLU	E	1	LYS	K	0	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
2	ASP	D	2	HIS	H	1	PHE	F	1	TYR	O
4	CYS	C	1	ILU	I	1	PRO	P	0	VAL	V

TOTAL NO. OF ACIDS = 21

* KOTAKI, A., J. BIOCHEM. (TOKYO), VOL.53, NO.1, PP.61-70, 1963

INSULIN A - HORSE

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I V E Q C C T G I C S L O Q L E N O C N *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU VAL GLU GLN CYS CYS THR GLY ILU CYS SER LEU TYR GLN
  LEU GLU ASN TYR CYS ASN ***
```

COMPOSITION

0	ALA	A	2	GLN	Q	2	LEU	L	1	SER	S
0	ARG	R	2	GLU	E	0	LYS	K	1	THR	T
2	ASN	N	2	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
4	CYS	C	2	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 21

* HARRIS, J. I., SANGER, F., AND NAUGHTON, M. A., ARCH. BIOCHEM. BIOPHYS., VOL.65, PP.427-438, 1956

SOME EVIDENCE FOR THE SEQUENCE WAS DERIVED FROM HOMOLOGY WITH BOVINE INSULIN.

INSULIN A - SHEEP

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I V E Q C C A G V C S L O Q L E N O C N *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU VAL GLU GLN CYS CYS ALA GLY VAL CYS SER LEU TYR GLN
  LEU GLU ASN TYR CYS ASN ***
```

COMPOSITION

1	ALA	A	2	GLN	Q	2	LEU	L	1	SER	S
0	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
2	ASN	N	2	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
4	CYS	C	1	ILU	I	0	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 21

* BROWN, H., SANGER, F., AND KITAI, R., BIOCHEM. J., VOL.60, PP.556-565, 1955

SOME EVIDENCE FOR THE SEQUENCE WAS DERIVED FROM HOMOLOGY WITH BOVINE INSULIN.

INSULIN A - SPERM WHALE, FIN-WHALE, PIG, AND HUMAN

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I V E Q C C T S I C S L O Q L E N O C N *

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU VAL GLU GLN CYS CYS THR SER ILU CYS SER LEU TYR GLN
  LEU GLU ASN TYR CYS ASN ***
```

COMPOSITION

0	ALA	A	2	GLN	Q	2	LEU	L	2	SER	S
0	ARG	R	2	GLU	E	0	LYS	K	1	THR	T
2	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
4	CYS	C	2	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 21

* BROWN, H., SANGER, F., AND KITAI, R., BIOCHEM. J., VOL.60, PP.556-565, 1955 (PIG)

SOME EVIDENCE FOR THE SEQUENCE WAS DERIVED FROM HOMOLOGY WITH BOVINE INSULIN.

HARRIS, J. I., SANGER, F., AND NAUGHTON, M. A., ARCH. BIOCHEM. BIOPHYS., VOL.65, PP.427-438, 1956 (SPERM WHALE)

HAMA, H., TITANI, K., SAKAKI, S., AND NARITA, K., J. BIOCHEM. (TOKYO), VOL.56, NO.3, PP.285-293, 1964 (FIN-WHALE)

THIS WORK CONFIRMED THE SEQUENCE ABOVE, EXCEPT GLU (E) AND GLN (Q) WERE INTERCHANGED AT POSITIONS 15 AND 17.

NICOL, D. S. H. AND SMITH, L. F., NATURE, VOL.187, NO.4736, PP.483-485, AUG. 6, 1960 (HUMAN)

INSULIN A - SEI-WHALE

```
      |
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1
1 G I V E Q C C A S T C S L O Q L E N O C N *
```

```
   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY ILU VAL GLU GLN CYS CYS ALA SER THR CYS SER LEU TYR GLN
  LEU GLU ASN TYR CYS ASN ***
```

COMPOSITION

1	ALA	A	2	GLN	Q	2	LEU	L	2	SER	S
0	ARG	R	2	GLU	E	0	LYS	K	1	THR	T
2	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
4	CYS	C	1	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 21

ISHIHARA, Y., SAITO, T., ITO, Y., AND FUJINO, M., NATURE, VOL.181, NO.4621, PP.1461-1469, MAY 24, 1958 (SEI-WHALE)

INSULIN B - BOVINE, SHEEP, HORSE, HUMAN, PIG, AND SPERM WHALE

TWO DISULPHIDE BONDS CONNECT THE A AND B CHAINS.
A7 IS BONDED TO B7 AND A20 IS BONDED TO B19. IN ADDITION THERE IS A BOND FROM A6 TO A11.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0

1 F V N Q H L C G S H L V E A L O L V C G E R G F F O T P K A

•

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15

1 PHE VAL ASN GLN HIS LEU CYS GLY SER HIS LEU VAL GLU ALA LEU

  TYR LEU VAL CYS GLY GLU ARG GLY PHE PHE TYR THR PRO LYS ALA

  ***
```

COMPOSITION

2	ALA	A	1	GLN	Q	4	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	1	LYS	K	1	THR	T
1	ASN	N	3	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	2	HIS	H	3	PHE	F	2	TYR	O
2	CYS	C	0	ILU	I	1	PRO	P	3	VAL	V

TOTAL NO. OF ACIDS = 30

• RYLE, A. P., SANGER, F., SMITH, L.F., AND KITAI, R., BIOCHEM. J., VOL. 60, PP.541-556, 1955 (BOVINE, SHEEP, AND PIG)

SANGER, F. AND TUPPY, H., BIOCHEM. J., VOL.49, PP.481-490, 1951 (BOVINE)
THE AMIDE GROUPS WERE SUBSEQUENTLY DETERMINED.

HARRIS, J. I., SANGER, F., AND NAUGHTON, M. A., ARCH. BIOCHEM. BIOPHYS., VOL.65, PP.427-438, 1956 (SPERM WHALE AND HORSE)

ISHIHARA, Y., SAITO, T., ITO, Y., AND FUJINO, M., NATURE, VOL.181, NO.4621, PP.1461-1469, MAY 24,1958 (SPERM AND SEI-WHALE)

NICOL, D. S. H. AND SMITH, L. F., NATURE, VOL.187, NO.4736, PP.483-485, AUG. 6, 1960 (HUMAN)

HUMAN INSULIN B CHAIN IS IDENTICAL WITH ABOVE EXCEPT THAT POSITION 30 IS THR (T).

INSULIN B - BONITO

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
1 A A N(P,H,L)C(G,S,H,L,V,E,A,L)O L(V,C,G,E)R G F F O Q P K •

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ALA ASN(PRO,HIS,LEU)CYS(GLY,SER,HIS,LEU,VAL,GLU,ALA,LEU)
  TYR LEU(VAL,CYS,GLY,GLU)ARG GLY PHE PHE TYR GLN PRO LYS ***
```

COMPOSITION

3	ALA	A	1	GLN	Q	4	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	1	LYS	K	0	THR	T
1	ASN	N	3	GLY	G	0	MET	M	0	TRP	W
0	ASP	D	2	HIS	H	2	PHE	F	2	TYR	O
2	CYS	C	0	ILU	I	2	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 29

• KOTAKI, A., J. BIOCHEM.(TOKYO), VOL.51, NO.4, PP.301-309, 1962

FIBRINOPEPTIDE A - BOVINE

FIBRINOPEPTIDES ARE THOSE PORTIONS OF VERTEBRATE FIBRINOGEN MOLECULES WHICH ARE PROTEOLYTICALLY REMOVED BY THE ENZYME THROMBIN. THEIR REMOVAL PERMITS SPONTANEOUS POLYMERIZATION OF THE PARENT MOLECULES TO FORM AN INSOLUBLE FIBRINOGEL. SINCE THE FUNCTION OF THE FIBRINOPEPTIDES IS RATHER NON-SPECIFIC, LARGE SEQUENCE CHANGES ARE OBSERVED AMONG CLOSELY RELATED SPECIES.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 E D G S D P P S G D F L T E G G G V R /
```

```
   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLU ASP GLY SER ASP PRO PRO SER GLY ASP PHE LEU THR GLU GLY
  GLY GLY VAL ARG ///
```

COMPOSITION

0	ALA	A	0	GLN	Q	1	LEU	L	2	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	1	THR	T
0	ASN	N	5	GLY	G	0	MET	M	0	TRP	W
3	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	2	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 19

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE A - SHEEP

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 A D D S D P V G G E F L A E G G G V R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ASP ASP SER ASP PRO VAL GLY GLY GLU PHE LEU ALA GLU GLY
  GLY GLY VAL ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	1	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	5	GLY	G	0	MET	M	0	TRP	W
3	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 19

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE A - GOAT

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 A D D S D P V G G E F L A E G G G V R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ASP ASP SER ASP PRO VAL GLY GLY GLU PHE LEU ALA GLU GLY
  GLY GLY VAL ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	1	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	5	GLY	G	0	MET	M	0	TRP	W
3	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 19

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE A - REINDEER

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 A D G S D P A G G E F(L,A,E,G,G,G,V)R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ASP GLY SER ASP PRO ALA GLY GLY GLU PHE(LEU,ALA,GLU,GLY,
  GLY,GLY,VAL)ARG ///
```

COMPOSITION

3	ALA	A	0	GLN	Q	1	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	6	GLY	G	0	MET	M	0	TRP	W
2	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 19

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE A - PIG

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7
1 A E V Q D K G E F L A E G G G V R /
```

```
   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA GLU VAL GLN ASP LYS GLY GLU PHE LEU ALA GLU GLY GLY GLY
  VAL ARG ///
```

COMPOSITION

2	ALA	A	1	GLN	Q	1	LEU	L	0	SER	S
1	ARG	R	3	GLU	E	1	LYS	K	0	THR	T
0	ASN	N	4	GLY	G	0	MET	M	0	TRP	W
1	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	0	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 17

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE A - HUMAN

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6
1 A D S G E G D F L A E G G G V R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ASP SER GLY GLU GLY ASP PHE LEU ALA GLU GLY GLY GLY VAL
  ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	1	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	5	GLY	G	0	MET	M	0	TRP	W
2	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 16

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE,* VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

PHOSPHO-SERINE OCCURS AT POSITION 3 IN ABOUT HALF THE MOLECULES. A MINOR COMPONENT FRAGMENT, WITH THE N TERMINAL ALANINE MISSING, HAS BEEN DETECTED IN ALL INDIVIDUALS.

FIBRINOPEPTIDE A - RABBIT

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6
1 V D P G E T S F L(T,E,G,G)D A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 VAL ASP PRO GLY GLU THR SER PHE LEU(THR,GLU,GLY,GLY)ASP ALA
  ARG ///
```

COMPOSITION

1	ALA	A	0	GLN	Q	1	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	2	THR	T
0	ASN	N	3	GLY	G	0	MET	M	0	TRP	W
2	ASP	D	0	HIS	H	1	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 16

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - BOVINE

PYRROLIDONE CARBOXYLIC ACID - AT AMINO END
SO4 ATTACHED TO TYROSINE AT POSITION 5

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
1 F P T D O D E G Q D D R P K V G L G A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 PHE PRO THR ASP TYR ASP GLU GLY GLN ASP ASP ARG PRO LYS VAL
  GLY LEU GLY ALA ARG ///
```

COMPOSITION

1	ALA	A	1	GLN	Q	1	LEU	L	0	SER	S
2	ARG	R	1	GLU	E	1	LYS	K	1	THR	T
0	ASN	N	3	GLY	G	0	MET	M	0	TRP	W
4	ASP	D	0	HIS	H	1	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	2	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 20

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202,
NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - SHEEP

SO4 ATTACHED TO TYROSINE AT POSITION 5

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
1 G O L D O D E V D D N R A K L P L D A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY TYR LEU ASP TYR ASP GLU VAL ASP ASP ASN ARG ALA LYS LEU
  PRO LEU ASP ALA ARG ///
```

COMPOSITION

2 ALA A	0 GLN Q	3 LEU L	0 SER S
2 ARG R	1 GLU E	1 LYS K	0 THR T
1 ASN N	1 GLY G	0 MET M	0 TRP W
5 ASP D	0 HIS H	0 PHE F	2 TYR O
0 CYS C	0 ILU I	1 PRO P	1 VAL V

TOTAL NO. OF ACIDS = 20

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - GOAT

SO4 ATTACHED TO TYROSINE AT POSITION 5

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
1 G O L D O D E V D D N R A K L P L D A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 GLY TYR LEU ASP TYR ASP GLU VAL ASP ASP ASN ARG ALA LYS LEU
  PRO LEU ASP ALA ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	3	LEU	L	0	SER	S
2	ARG	R	1	GLU	E	1	LYS	K	0	THR	T
1	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
5	ASP	D	0	HIS	H	0	PHE	F	2	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 20

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP.147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - REINDEER

PYRROLIDONE CARBOXYLIC ACID - AT AMINO END
SO4 ATTACHED TO TYROSINE AT POSITION 4
A MUTANT HAS BEEN FOUND WHERE GLYCINE REPLACES HISTIDINE
IN POSITION 9.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9

1 L A D O D E V(E,H,D)R A K L H L D A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15

1 LEU ALA ASP TYR ASP GLU VAL(GLU,HIS,ASP)ARG ALA LYS LEU HIS

  LEU ASP ALA ARG ///
```

COMPOSITION

3	ALA	A	0	GLN	Q	3	LEU	L	0	SER	S
2	ARG	R	2	GLU	E	1	LYS	K	0	THR	T
0	ASN	N	0	GLY	G	0	MET	M	0	TRP	W
4	ASP	D	2	HIS	H	0	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 19

* DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202,
NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - PIG

SO4 ATTACHED TO TYROSINE AT POSITION 4

```
  1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9
1 A I D O D E D E D G R P K V H V D A R /
```

```
   1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
1 ALA ILU ASP TYR ASP GLU ASP GLU ASP GLY ARG PRO LYS VAL HIS
  VAL ASP ALA ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	0	LEU	L	0	SER	S
2	ARG	R	2	GLU	E	1	LYS	K	0	THR	T
0	ASN	N	1	GLY	G	0	MET	M	0	TRP	W
5	ASP	D	1	HIS	H	0	PHE	F	1	TYR	O
0	CYS	C	1	ILU	I	1	PRO	P	2	VAL	V

TOTAL NO. OF ACIDS = 19

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - HUMAN

PYRROLIDONE CARBOXYLIC ACID - AT AMINO END
PHOSPHO-SERINE OCCURS IN POSITION 11.

```
  1 2 3 4 5 6 7 8 9 0 1 2 3
1 G V N D N E E G F F S A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13
1 GLY VAL ASN ASP ASN GLU GLU GLY PHE PHE SER ALA ARG ///
```

COMPOSITION

1	ALA	A	0	GLN	Q	0	LEU	L	1	SER	S
1	ARG	R	2	GLU	E	0	LYS	K	0	THR	T
2	ASN	N	2	GLY	G	0	MET	M	0	TRP	W
1	ASP	D	0	HIS	H	2	PHE	F	0	TYR	O
0	CYS	C	0	ILU	I	0	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 13

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

FIBRINOPEPTIDE B - RABBIT

SO4 ATTACHED TO TYROSINE AT POSITION 4

```
  1 2 3 4 5 6 7 8 9 0 1 2 3
1 A D D O(D,E,P,L,D,V)D A R /

   1   2   3   4   5   6   7   8   9  10  11  12  13
1 ALA ASP ASP TYR(ASP,GLU,PRO,LEU,ASP,VAL)ASP ALA ARG ///
```

COMPOSITION

2	ALA	A	0	GLN	Q	1	LEU	L	0	SER	S
1	ARG	R	1	GLU	E	0	LYS	K	0	THR	T
0	ASN	N	0	GLY	G	0	MET	M	0	TRP	W
5	ASP	D	0	HIS	H	0	PHE	F	1	TYR	O
0	CYS	C	0	ILU	I	1	PRO	P	1	VAL	V

TOTAL NO. OF ACIDS = 13

• DOOLITTLE, R. F. AND BLOMBACK, B., NATURE, VOL. 202, NO. 4928, PP. 147-152, APRIL 11, 1964

Immunoglobulins are serum proteins distinguishable by electrophoretic mobilities, sedimentation coefficients and differential solubilities in variable ethanol-salt solutions. Of these, the gamma globulins are associated with normal antibody function. A proposal for the structure of gamma globulin has been made by Porter (1959) and Fleishman et al, (1963).



Gamma globulin is thought to be a tetramer consisting of two pairs of identical polypeptide chains held in a particular configuration by disulfide bonds. There are two L (m.w. 20-25,000 each) and two H chains (m.w. 50,000-55,000 each). Because of the chemical problems associated with elucidation of gamma globulin structure, attention has turned to the abundantly produced, structurally similar globulins found in multiple myeloma.

Bence-Jones proteins are found exclusively in the urine of all multiple myeloma patients, and probably represent abnormal protein synthesized by the multiple myeloma tumor cell. They are thought to be made exclusively of L chains, related to gamma globulins, (Edelman and Gally, 1962, S. Cohen, 1963, Putnam 1962). It is thought that determination of the amino acid sequence of a particular individual's Bence-Jones protein would reflect a homologous sequence in that individual's antibody structure, thereby partially elucidating the structure of gamma globulin.

Cohen, S. Biochem. J. Vol. 89, p. 334 (1963)
Edelman, G. M. and Gally, J. A. J. Exp. Med. Vol. 116, p. 207 (1962)
Fleishman, J. B. et al. Biochem. J. Vol. 88, p. 220 (1963)
Porter, R. R. Biochem. J. Vol. 73, p. 119 (1959)
Putnam, F. W. Biochim. Biophys. Acta. Vol. 63, p. 539 (1962)

BENCE-JONES PROTEIN

```
    1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0
  1 D(T,S,S,S,E,E,P,M,I)L S(S,G,A,V)D R(D,T,T,S,S,E,E,A,V,I,I,I,
 31 F,C)L(O,D,W,E,E,P,G)K K A P K L L I O D A S K L E(S,P,G,A,V)
 61 R F S(D,T,T,S,G,G,G)F T(D,S,S,E,E,P,I,L)I A T O(D,D,T,E,E,P,
 91 L,L,C,O,F,F)G(T,G,G)K V D F K R T(S,P,A,A,V)V F I(D,S,E,E,P,
121 P,F)L K S(T,S,G,A)V(V,C)L L D(D,P,F)O R E A K V E W K V(D,D,
151 D,S,S,E,E,G,A,L)E S(D,T,S,E,E,V)K D(T,S)O S S S T L L T L S
181 K A D O E K H K L O A C E V(T,E,G,H)L S(T,S,P,V)K S F D R G
211 E C *

     1   2   3   4   5   6   7   8   9  10  11  12  13  14  15
  1 ASP(THR,SER,SER,SER,GLU,GLU,PRO,MET,ILU)LEU SER(SER,GLY,ALA,
    VAL)ASP ARG(ASP,THR,THR,SER,SER,GLU,GLU,ALA,VAL,ILU,ILU,ILU,
 31 PHE,CYS)LEU(TYR,ASP,TRP,GLU,GLU,PRO,GLY)LYS LYS ALA PRO LYS
    LEU LEU ILU TYR ASP ALA SER LYS LEU GLU(SER,PRO,GLY,ALA,VAL)
 61 ARG PHE SER(ASP,THR,THR,SER,GLY,GLY,GLY)PHE THR(ASP,SER,SER,
    GLU,GLU,PRO,ILU,LEU)ILU ALA THR TYR(ASP,ASP,THR,GLU,GLU,PRO,
 91 LEU,LEU,CYS,TYR,PHE,PHE)GLY(THR,GLY,GLY)LYS VAL ASP PHE LYS
    ARG THR(SER,PRO,ALA,ALA,VAL)VAL PHE ILU(ASP,SER,GLU,GLU,PRO,
121 PRO,PHE)LEU LYS SER(THR,SER,GLY,ALA)VAL(VAL,CYS)LEU LEU ASP
   (ASP,PRO,PHE)TYR ARG GLU ALA LYS VAL GLU TRP LYS VAL(ASP,ASP,
151 ASP,SER,SER,GLU,GLU,GLY,ALA,LEU)GLU SER(ASP,THR,SER,GLU,GLU,
    VAL)LYS ASP(THR,SER)TYR SER SER SER THR LEU LEU THR LEU SER
181 LYS ALA ASP TYR GLU LYS HIS LYS LEU TYR ALA CYS GLU VAL(THR,
    GLU,GLY,HIS)LEU SER(THR,SER,PRO,VAL)LYS SER PHE ASP ARG GLY
211 GLU CYS ***
```

COMPOSITION

13	ALA	A	0	GLN	Q	17	LEU	L	29	SER	S
5	ARG	R	24	GLU	E	14	LYS	K	17	THR	T
0	ASN	N	13	GLY	G	1	MET	M	2	TRP	W
20	ASP	D	2	HIS	H	10	PHE	F	8	TYR	O
5	CYS	C	8	ILU	I	11	PRO	P	13	VAL	V

TOTAL NO. OF ACIDS = 212

• HILSCHMANN, N. AND CRAIG, L.C., PROC. NATL. ACAD. SCI. U.S., VOL.53, NO.6, PP.1403-1409, 1965

AUTHOR INDEX

			PAGE
ACHER, R.	TI	BOPA	5.001
	PR	BOAR	8.101
AMBLER, R. P.	CY	PS	1.006
	AZ	PS	3.003
ANDERER, F. A.	TM	TM	6.001
ANSFIELD, M. J.	TI	BOPA	5.001
BAHL, O. P.	CY	RS	1.009
BAGLIONI, C.	GL	HUH	2.020
BARNAFI, L.	TN	BOBM	8.102
	TN	PGBM	8.203
BARTSCH, R. G.	DH	CH	3.001
BEALE, D.	GL	HUH	2.020
BEHRENS, O. K.	GN	BO	8.001
BELL, P. H.	TN	PGAC	8.206
BENSON, A.	CY	PG	1.005
	FE	CP	3.002
BERNIER, I.	LS	CH	7.201
BLAKE, C. C. F.	LS	CH	7.201
BLOMBACK, B.	FB	BOA	9.001
	FB	SHA	9.002
	FB	GTA	9.003
	FB	RDA	9.004
	FB	PGA	9.005
	FB	HUA	9.006
	FB	RTA	9.007
	FB	BOB	9.101
	FB	SHB	9.102
	FB	GTB	9.103
	FB	RDB	9.104
	FB	PGB	9.105
	FB	HUB	9.106
	FB	RTB	9.107
BRAUNITZER, G.	GL	HUHA	2.001
	GL	HUHB	2.002
	GL	HOHA	2.006
BROMER, W. W.	GN	BO	8.001
BROWN, H.	IS	SHA	8.304
	IS	WPA	8.305
BROWN, J. R.	TR	BOCH	7.001
BROWN, L. H.	AZ	PS	3.003
BUETTNER-JANUSCH, J.	GL	LEHB	2.007
CANFIELD, R.	LS	CH	7.201
CHAUVET, J.	TI	BOPA	5.001
	PR	BOAR	8.101
CHUNG, D.	TN	SBAC	8.207
COHEN, S.			10.000
COLE, D.	TN	SBAC	8.207
CORMICK, J.	GL	HUHA	2.001
	GL	HUHG	2.003
CRAIG, L. C.	BJ	HU	10.001
DAVIS, D. S.	TN	PGAC	8.206
DIXON, J. S.	TN	BPAM	8.201
	TN	HOBM	8.204
	TN	SBAC	8.207
DLOUHA, V.	TI	BOPA	5.001

DOOLITTLE, R. F.	FB	BOA	9.001
	FB	SHA	9.002
	FB	GTA	9.003
	FB	RDA	9.004
	FB	PGA	9.005
	FB	HUA	9.006
	FB	RTA	9.007
	FB	BOB	9.101
	FB	SHB	9.102
	FB	GTB	9.103
	FB	RDB	9.104
	FB	PGB	9.105
	FB	HUB	9.106
	FB	RTB	9.107
DUS, K.	DH	CH	3.001
DU VIGNEAUD, V.	PR	BOAR	8.101
	PR	PGLS	8.102
	PR	BOOX	8.103
EDELMAN, G. M.			10.000
EDMUNDSON, A. B.	GL	WHMY	2.101
EFRON, M. L.	GL	HUH	2.020
EIGNER, E. A.	TN	PGAC	8.206
ELLIOT, D. F.	PR	BOHY	8.104
FLEISHMAN, J. B.			10.000
FRAENKEL-CONRAT, H.	TM	TM	6.001
FRATER, R.	PA	PA	7.101
FUJINO, M.	IS	WHA	8.306
	IS	BOB	8.321
FUNATSU, G.	TM	TM	6.001
GALLY, J. A.			10.000
GEHRING-MULLER, R.	GL	HUHA	2.001
	GL	HUHB	2.002
GERALD, P. S.	GL	HUH	2.020
GESCHWIND, I. I.	TN	BOBM	8.202
	TN	PGBM	8.203
	TN	SBAC	8.207
HAMA, H.	IS	WPA	8.305
HARRIS, J. I.	TN	BPAM	8.201
	TN	PGBM	8.203
	TN	HUBM	8.205
	TN	SBAC	8.207
	IS	HOA	8.303
	IS	WPA	8.305
	IS	BOB	8.321
HARTLEY, B. S.	TR	BOCH	7.001
HIGA, H.	CY	PG	1.005
HILL, R. J.	GL	HUHA	2.001
HILL, R. L.	GL	LEHB	2.007
	GL	HUH	2.020
HILSCHMANN, N.	GL	HUHA	2.001
	GL	HUHB	2.002
	BJ	HU	10.001
HILSE, K.	GL	HUHA	2.001
	GL	HUHB	2.002

HOBOM, G.	GL	HUHA	2.001
	GL	HUHB	2.002
HOWARD, K. S.	TN	PGAC	8.206
HUNT, J. A.	GL	HUH	2.020
INGRAM, V. M.	GL	HUH	2.020
ISHIHARA, Y.	IS	WHA	8.306
	IS	BOB	8.321
ITO, Y.	IS	WHA	8.306
	IS	BOB	8.321
JAUREGUI-ADELL, J.	LS	CH	7.201
JOLLES, J.	LS	CH	7.201
JOLLES, P.	LS	CH	7.201
JONES, R. T.	GL	HUHG	2.003
KAMEN, M. D.	DH	CH	3.001
KASSELL, B.	TI	BOPA	5.001
KAUFFMAN, D. L.	TR	BOCH	7.001
	TR	BOTR	7.002
KEIL, B.	TR	BOCH	7.001
KIMMEL, J.	PA	PA	7.101
KINGMA, S.	GL	HUH	2.020
KITAI, R.	IS	BOA	8.301
	IS	SHA	8.304
	IS	WPA	8.305
	IS	BOB	8.321
KOENING, D. F.	LS	CH	7.201
KONIGSBERG, W.	GL	HUHA	2.001
KOSTKA, V.	TR	BOCH	7.001
KOTAKI, A.	IS	BNA	8.302
	IS	BNB	8.322
KREIL, G.	CY	HO	1.003
	CY	TF	1.007
LANDMANN, W. A.	TN	PGAC	8.206
LASKOWSKI, M.	TI	BOPA	5.001
LAWLER, H. C.	PR	BOAR	8.101
	PR	PGLS	8.102
LEHMANN, H.	GL	HUH	2.020
LERNER, A. B.	TN	BPAM	8.201
LI, C. H.	TN	BPAM	8.201
	TN	BOBM	8.202
	TN	PGBM	8.203
	TN	HOBM	8.204
	TN	SBAC	8.207
LIGHT, A.	PA	PA	7.101
LIU, A. K.	LS	CH	7.201
MAIR, G. A.	LS	CH	7.201
MARGOLIASH, E.	CY	CH	1.002
	CY	HO	1.003
	CY	PG	1.005
MARTIN, N.	GL	HUH	2.020
MATSUBARA, H.	CY	HU	1.004
	CY	PG	1.005
MATSUDA, G.	GL	HOHA	2.006
MELOUN, B.	TI	BOPA	5.001
	TR	BOCH	7.001
MICHL, H.	PR	BOOX	8.103

MOORE, S.	RN	BO	4.001
MOWER, H. F.	FE	CP	3.002
MULLER, C. J.	GL	HUH	2.020
MURAKAMI, H.	CY	BY	1.001
MURAYAMA, M.	GL	HUH	2.020
NAKASHIMA, T.	CY	PG	1.005
	FE	CP	3.002
NARITA, K.	CY	BY	1.001
	IS	WPA	8.305
NAUGHTON, M. A.	IS	HOA	8.303
	IS	WPA	8.305
	IS	BOB	8.321
NEEDLEMAN, S. B.	CY	CH	1.002
	CY	PG	1.005
NEURATH, H.	TR	BOTR	7.002
NICOL, D. S. H.	IS	WPA	8.305
	IS	BOB	8.321
NORTH, A. C. T.	LS	CH	7.201
NOUVEL, G.	TI	BOPA	5.001
PALEUS, S.	CY	PG	1.005
	CY	RR	1.010
	CY	SM	1.011
PAULING, L.	GL	HUH	2.020
PHILLIPS, D. C.	LS	CH	7.201
PEART, W. S.	PR	BOHY	8.104
PORTER, R. R.			10.000
POPENOE, E. A.	PR	BOAR	8.101
	PR	PGLS	8.102
POSPISILOVA, D.	TI	BOPA	5.001
PRUSIK, Z.	TR	BOCH	7.001
PUTNAM F. W.			10.000
RAACK, I. D.	TN	SBAC	8.207
RADICEVIC, M.	TI	BOPA	5.001
RESSLER, C.	PR	BOOX	8.103
RHINESMITH, H. W.	GL	HUH	2.020
ROOS, P.	TN	PGBM	8.203
RUDLOFF, V.	GL	HUHA	2.001
	GL	HUHB	2.002
RYLE, A. P.	IS	BOA	8.301
	IS	BOB	8.321
SAITO, T.	IS	WHA	8.306
	IS	BOB	8.321
SAKAKI, S.	IS	WPA	8.305
SANGER, F.	IS	BOA	8.301
	IS	HOA	8.303
	IS	SHA	8.304
	IS	WPA	8.305
	IS	BOB	8.321
SARMA, V. R.	LS	CH	7.201
SCHNEIDER, R. G.	GL	HUH	2.020
SCHRAMM, G.	TM	TM	6.001
SCHROEDER, W. A.	GL	HUHA	2.001
	GL	HUHG	2.003
	GL	HUH	2.020
SCHWARTZ, H. C.	GL	HUH	2.020

SHELTON, J. B.	GL	HUHA	2.001
	GL	HUHG	2.003
SHELTON, J. R.	GL	HUHA	2.001
	GL	HUHG	2.003
SHEPHERD, R. G.	TN	PGAC	8.206
SINN, L. G.	GN	BO	8.001
SMILLIE, L. B.	TR	BOCH	7.001
SMITH, D. B.	GL	HOHB	2.005
SMITH, E. L.	CY	HO	1.003
	CY	HU	1.004
	CY	RS	1.009
	PA	PA	7.101
SMITH, L. F.	IS	BOA	8.301
	IS	WPA	8.305
	IS	BOB	8.321
SMYTH, D. G.	RN	BO	4.001
SORM, F.	TI	BOPA	5.001
	TR	BOCH	7.001
STEIN, W. H.	RN	BO	4.001
STEWART, J. W.	CY	CH	1.002
	CY	PG	1.005
SWENSON, R. T.	GL	HUH	2.020
TANAKA, M.	FE	CP	3.002
THOMPSON, E. O. P.	IS	BOA	8.301
TITANI, K.	CY	BY	1.001
	IS	WPA	8.305
TRIPPETT, S.	PR	BOOX	8.103
TSUGITA, A.	TM	TM	6.001
TUPPY, H.	CY	HO	1.003
	CY	PG	1.005
	CY	SW	1.008
	CY	RR	1.010
	CY	SM	1.011
	PR	BOOX	8.103
	IS	BOB	8.321
UHLIG, H.	TM	TM	6.001
WALSH, K.	TR	BOTR	7.002
WATSON-WILLIAMS, E. J.	GL	HUH	2.020
WEBER, E.	TM	TM	6.001
WHITE, W. F.	TN	PGAC	8.206
WITTMANN, H. G.	TM	TMD	6.002
WITTMANN-LIEBOLD, B.	GL	HUHA	2.001
	GL	HUHB	2.002
	TM	TMD	6.002
YAOI, Y.	CY	BY	1.001
YASUNOBU, K. T.	CY	PG	1.005
	FE	CP	3.002
ZUCKERKANDL, E.	GL	GOHB	2.004