


## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

E 7.5-10402  
2R-143395  
RECEIVED  
NASA STI FACILITY  
ACQ. BR.  
SEP 19 1975

QUARTERLY REPORT TO NASA ON  
LANDSAT 2 PROJECT NO.29020  
HYDROLOGICAL INVESTIGATIONS IN NORWAY



FROM JOHNNY SKORVE, CO-INVESTIGATOR  
UNIVERSITY OF OSLO

"Made available under NASA sponsorship  
in the interest of early and wide dis-  
semination of Earth Resources Survey  
Program information and without liability  
for any use made thereof."

INTRODUCTION

LANDSAT 2 images of Southern Norway from February and  
March 1975 have been studied. The scenes show a winter  
situation, and at that time of the year, the snowmelting  
is neglectable.

DESCRIPTION AND PRELIMINARY RESULTS

Except for a rim along the southeastern coast and the  
outer Oslofjord lowland, the area is snowcovered. These  
February and March images are the first LANDSAT winter  
pictures of the Oslo region. The winter scenes reflect  
interesting features.

Most of Norway experienced an extremely mild winter in  
1975, and the LANDSAT 2 images do show the resulting  
extraordinary icesituation. Fjords, lakes and rivers that  
during winter time nearly always are completely icecovered,  
were this year icefree or only partly icecovered.

Changes that have taken place during two LANDSAT cycles,  
have been determined by comparing the 14. February and  
21. March images.

The snow on the LANDSAT 2 images do strongly enhance  
features like roads and urban settlements. One illustrating  
example is seen on the flat, open terrain east of lake Mjøsa.  
Here small villages and narrow roads are clearly seen, while  
they are completely invisible on snowfree images.

N75-33457

Unclas  
00402


G3/43

(E75-10402) HYDROLOGICAL INVESTIGATIONS IN  
NORWAY Quarterly Report (Oslo Univ.) 2 P  
CSCL 05B  
HC \$3.25

A study of scene no.2024-10031 and 2024-10035 of 15.February proves to be of geological interest. This is because of the enhancement introduced through the combination of low solar angle and snowcover. These LANDSAT images should be very valuable as supplementary material for the structure and lineament mapping already done on the basis of LANDSAT 1 images of Norway from 1972 and 73.

I find the quality of LANDSAT 2 imagery superior to that of number one.

The study of spring and summer images will now be initiated for mapping the decrease in areal extent of the snowcover and determination of snowlines. The LANDSAT 2 data will be compared with oblique angle aerial photographs of selected areas taken simultaneously with LANDSAT 2 passes over Norway. Ground truth measurements and data will be used in the further interpretation of the LANDSAT 2 images.



Johnny Skørve

Oslo, 28.August 1975