

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASA CR  
147748

**FINAL REPORT**

**Contract NAS 9-13647**

**Support of In-Flight Experiments**

**May 31, 1976**

(NASA-CR-147748) SUPPORT OF IN-FLIGHT EXPERIMENTS FINAL REPORT (TEXAS A&M UNIV.)  
175 P HC \$6.75 CSCL 06F  
N76-25758  
UNCLAS  
G3/51 28294

**Submitted to**

**National Aeronautics and Space Administration**

**by**

**Dr. Karl P. Kuchnow  
Biology Department  
Texas A&M University**



FINAL REPORT

Contract NAS 9-13647

Support of In-Flight Experiments

May 31, 1976

Submitted to

National Aeronautics and Space Administration

by

Dr. Karl P. Kuchnow

Biology Department

Texas A&M University

## TABLE OF CONTENTS

	page
I. BIBLIOGRAPHY OF OVA AND SPERM PRESERVATION TECHNIQUES.	1
A. Vertebrate Embryology and Gamete Preservation	1
B. Fish Embryology and Reproduction	4
C. <u>Fundulus</u> Embryology and Reproduction	22
II. OVA PRESERVATION TESTS	35
A. Ficoll-Dextran Solutions	35
B. Holtfreter's Solution	40
C. Sucrose	40
D. Dry Preservation	41
E. Salinity	42
F. pH	48
G. Analysis of Ovarian Fluid	70
III. SPERM PRESERVATION TESTS	72
A. Ficoll-Dextran Solutions	72
B. Preliminary Tests on Sperm, Temperature, pH, Salinity Interactions	74
C. Effect of Buffer, ATP, Oxygen and Salinity of Post-Preser- vation Mixing Medium	76
D. Effect of ATP Concentration	77
E. Temperature	78
F. Salinity	79

	ii
G. pH	91
H. Whole Testis	91
IV. DISCUSSION OF OVA AND SPERM PRESERVATION TESTS.	107
A. Ova Preservation	107
B. Sperm Preservation	108
V. EMBRYO PRESERVATION TESTS	110
VI. ASTP VISUAL ORIENTATION TESTS	118
A. Bibliography of Pertinent Fish Vision Literature	119
B. Protocol of Visual Acclimation Procedures	125
C. Protocol of ASTP Film Analysis	127
D. Results of ASTP Film Analysis	129
VII. ELECTRON MICROSCOPY	160

## INTRODUCTION

The specimens of Fundulus heteroclitus used in this study were obtained from stock maintained at the National Aeronautics and Space Administration laboratories in Houston, Texas. These estuarine minnows were originally collected from the near shore water around Beaufort, North Carolina. In the laboratory fish were held in four LS-700 Living Stream Tanks.<sup>1</sup> The tanks were filled with approximately 150 gal. of 21<sup>0</sup>/<sub>00</sub> synthetic sea water.<sup>2</sup>

The water quality was monitored regularly for uniformity of salinity, temperature, pH, and ammonia concentration. For this study it was necessary to maintain the fishes in a state of continuous gamete production. This was accomplished by simulating the natural physical conditions which prevail in the Beaufort, North Carolina area during the spring and early summer breeding season. The 21<sup>0</sup>/<sub>00</sub> sea water was held at 20 ± 2C, and a light-dark cycle of 16 hours light and 8 hours dark was enforced. The fish were fed once daily (mid afternoon) with "Oregon Moist Pellets".

The fresh ova were stripped by holding the female in a damp paper towel with the ventral surface up and gently exerting pressure to force the ova out the ovipositor. Because of the relatively small quantity of sperm obtainable from males, it proved advantageous to

---

<sup>1</sup> Frigid Units, Inc., Toledo, Ohio.

<sup>2</sup> Instant Ocean, Aquarium Systems, Inc., Eastlake, Ohio.

collect the sperm in a pipette. Males were held ventral surface up and the urogenital aperture was wiped clean. Pressure was applied to the sides of the abdomen and, using a Pasteur pipette (equipped with a small rubber suction bulb) placed over the spermatid opening, the sperm was drawn up as it exited. This collection method made it easy to harvest the small quantities of sperm and also prevented fecal matter and urine from contaminating the sperm sample.

The sections that follow outline the various techniques used and the results obtained of attempts to achieve satisfactory preservation of ova and sperm of Fundulus heteroclitus, in terms of the greatest amount of time that fertility could be retained, and also the retention of maximum fertility. Also included in this report are the results of tests on delayed embryogenesis, should the preservation of individual gametes not prove feasible, as well as preliminary treatment of data on the orientation of ASTP juveniles.

## I. BIBLIOGRAPHY OF OVA AND SPERM PRESERVATION TECHNIQUES

## A. Vertebrate Embryology and Gamete Preservation.

1. Beatty, R.A., S. Gluecksohn-Waelsch. 1972. Genetics of the Spermatozoan. Proc. Int. Symp. Edinburgh.
2. Doebbzzer, G.F. 1966. Cryoprotective agents; a review of structure and function. Cryobiology 3: 2-11.
3. Flipse, R.J. and J.O. Almquist. 1956. Diluters for bovine semen. J. Dairy Sci. 39: 1690-1696.
4. Graham, E.F., Crabo, B.G., Brown, K.L. 1972. Effects of some zwitterion buffers on the freezing and storage of spermatozoa. J. Dairy Sci. 55(3): 372-378.
5. Gray, J., 1928. The effect of dilution on the activity of spermatozoa. Brit. J. Exp. Biol. 5: 362-365.
6. Harvey, E.B. 1930. The effect of lack of O<sub>2</sub> on the sperm and unfertilized eggs of Arabacia punctulata, and on fertilization. Biol. Bull. 89: 288-292.
7. Hyman, Libbie H. 1921. The metabolic gradients of vertebrate embryos. Biol. Bull. Vol. 40: 32-72.
8. Jones, R.C. 1965. The use of dimethyl sulphoxide, glycerol, and reconstituted skim milk for preservation of ram spermatozoa. Australian J. Biol. Sci. 18: 887-900.

9. Martin, I.C.A. 1965. The effects of rate of cooling to 5 C, storage at 5 C, glycerol concentration, NaCl, fructose and glycerine on revival of deep frozen bull sperm.  
J. Agr. Sci. 64: 425-432.
10. Mohri, H., and I. Yasumaso. 1963. Studies on the respiration of sea urchin spermatozoa; the effect of  $P_{CO_2}$ .  
J. Exp. Biol. 40: 573-586.
11. Monroy, A., and B. Baccetti. 1975. Morphological changes in the surface of the egg of Xenopus laevis in the course of development. I. Fertilization and early cleavage.  
J. Ultr. Res. 50: 131-142.
12. Proudfoot, F.G., 1973. Effects of high pressure gases on the motility and fertilizing capacity of avian spermatozoa stored in vitro. J. Reprod. Fertil. 31(3): 367-371.
13. Segal, J.J., R. Crozier, P.A. Corfman, P.G. Condliffe. 1973. The regulation of mammalian reproduction. C.C. Thomas, Springfield, Ill.
14. Shaver, J.R., S.H. Barch, L.C. Attinasi. 1973. Effects of immune globulins on the fertilizing capacity of frog spermatozoa. Biol. Reprod. 8: 112-118.
15. Shivers, G.A., and A.B. Dudkiewicz. 1972. Inhibition of sperm-egg interaction by specific antibody. Science 178(4066): 1211-1213.

16. Smith, K.D., and E. Steinberger. 1973. Survival of spermatozoa in a human sperm bank. J. Amer. Med. Assoc. 223: 774-777.
17. Southwick, W.E. 1939. Activity-preventing and egg-sea water neutralizing substances from spermatozoa of Echinometra subangularis. Biol. Bull. 77: 147-156.
18. Whittingham, D.G., S.P. Leibo, P. Mazur. 1972. Survival of mouse embryos frozen to  $-196^{\circ}$  and  $-269^{\circ}$ C. Science 178: 412-414.
19. Yoshizak, N., and C. Katagiri. 1975. Cellular basis for the production and secretion of the hatching enzyme by frog embryos. J. Exp. Zool. 192: 203-236.
20. Ziomek, C.A., and D. Epel. 1975. Polyspermy block of Spisula eggs is prevented by cytochalasin-B. Science 189: 139-141.

## B. Fish Embryology and Reproduction

1. Afzelius, B.A., and A. Murray. 1957. The acrosomal reaction of spermatozoa during fertilization or treatment with egg water. Exp. Cell. Res. 12: 325-337.
2. Aravindan, C.M., and K.G. Padmanabhan. 1972. Formation of the micropyle in Tilapia mossambica Peters and Stigmato-  
gobius jowonicus. Acta. Zool. 53: 45-47.
3. Arcarese, G., G. Ravagnan, P. Ghittino. 1972. First positive results in the artificial propagation of sea bass (Dicentrarchus labrax) on a vast scale. Riv. Ital. Piscic. Ittiop. 7: 27-33.
4. Arndt, E.A. 1960. Untersuchungen über die Eihüllen von Cyprinidien. Z. Zellforsch. 52: 315-327.
5. Auld, A.H., and J.R. Schubel. 1974. Preliminary observations on the efficiency of a commercially available fungal inhibitor and its toxicity to fish eggs. Chesapeake Science 15: 115-116.
6. Azevedo, E. 1975. Evolution of the envelopes of the oocytes in Xiphophorus helleri (Teleost, viviparous). J. de Microsc. 21: 21-43-54.
7. Bailey, W., and R.L. Boyd. 1971. A preliminary report on spawning and rearing of grass carp. (Ctenopharyngodon idello) in Arkansas. Proc. 24th Ann. Conf. Southeast Assoc. Game and Fish Commissioners. 1970: 560-569.

8. Barrett, I. 1951. Fertility of salmonid eggs and sperm after storage. J. Fish. Res. Bd. Can. 8(3): 125-133.
9. Becher, H. 1928. Beitrag zur feineren Struktur der Zona radiata des Knochenfischeies und über ein durch die Struktur der Eihülle bedingtes optisches Phänomen. Z. Mikroskop. Anat. Forsch. 13: 591-624.
10. Bell, G.R., G.E. Hoskins, and J.W. Bagshaw. 1969. On the structure and enzymatic degradation of the external membrane of the salmon egg. Canad. J. Zool. 47: 146-148.
11. Blaxter, J.H.S. 1953. Sperm storage and cross fertilization of spring and autumn spawning herring. Nature 172: 1189.
12. Blaxter, J.H.S. 1955. Herring rearing. I. The storage of herring gametes. Marine Res. Scotland. 3: 1-12.
13. Blaylock, B.G., and N.A. Griffith. 1971. A laboratory technique for spawning carp. Prog. Fish. Cult. 33: 48-50.
14. Bowden, B.S. 1969. A new method for obtaining precisely timed inseminations in viviparous fishes. Prog. Fish. Cult. 31: 229-230.
15. Brown, C.H. 1955. Egg-capsule proteins of selachians and trout. Quart. J. Microsc. Sci. 96: 483-488.
16. Bushuyev, V.P. 1971. Interspecific differences in the heat resistance of the sperm of some Pacific Salmon (Genus Oncorhynchus). J. Ichthyol. 11: 382-386.

17. Butcher, A.D. 1944. Preliminary observations on storage of the milt of trout. Austr. J. Sci. 7: 23-25.
18. Chaudhry, H.S. 1956. The origin and structure of the zona pellucida in the ovarian eggs of teleosts. Z. Zellforsch. 43: 478-485.
19. DeCiechowski, J. 1965. The development of fish embryos in a non-aqueous medium. Acta. Embryol. Morphol. Exptl. 8: 183-188.
20. Demski, L.S., D.H. Bauer, J.W. Gerald. 1975. Sperm release evoked by electrical stimulation of the fish brain: a functional anatomical study. J. Exp. Zool. 191: 215-278.
21. Dettlaff, T.A. 1962. Cortical changes in acipenserid eggs during fertilization and artificial activation. J. Embryol. Exp. Morph. 10: 1-26.
22. de Vlaming, V.L. 1972. Environmental controls of teleost reproductive cycles: a brief review. J. Fish Biol. 4: 131-140.
23. Donaldson, F.M., J.D. Funk, F.C. Withler, R.B. Morley. 1972. Fertilization of pink salmon (Oncorhynchus gorbuscha) ova by spermatozoa from gonadotropin-injected juveniles. J. Fish. Res. Bd. Can. 29: 13-18.
24. Eddy, F.B. 1974. Osmotic properties of the perivitelline fluid and some properties of the chorion of Atlantic

- Salmon eggs (Salmo salar) J. Zool. 174: 237-243.
25. Eggert, B. 1929. Entwicklung und Bau der Eier von Salaria  
flavoumbrinus Rupp. Zool. Anz. 83: 241-253.
26. Eigenmann, C.H. 1890. On the egg-membranes and micropyle of  
some osseous fishes. Bull. Mus. Comp. Zool. Harvard Coll.  
19: 129-154.
27. Ellis, W.G., and J.W. Jones, 1939. The activity of the spermato-  
zoa of Salmo salar in relation to osmotic pressure.  
J. Exp. Biol. 16: 530-534.
28. Fish, F.F., and R.E. Burrows. 1939. Notes on the effect of low  
temperatures on eyed eggs. Prog. Fish. Cult. 45: 28-31.
29. Fisher, H., O. Hug, W. Lippert. 1952. EM Studien an Forellen-  
spermatozoen und ihren Zellkernen. Chromosoma 5: 69-80.
30. Flügel, H. 1967. Licht-elektronenmikroskopische Untersuchungen  
an Oozyten und Eiern einiger Knochenfische. Z. Zellforsch.  
Mikrosk. Anat. 83: 82-116.
31. Foerster, R.E. 1965. Effect of retention of sperm and ova of  
sockeye salmon Oncorhynchus nerka in water and without  
addition of water on fertility. J. Fish. Res. Bd. Can.  
22(6): 1503-1521.
32. Ginzburg, A.S. 1961. The block to polyspermy in sturgeon and  
trout with special reference to the role of cortical  
granules (alveoli). J. Embryol. Exp. Morph. 9: 173-190.

33. Ginzburg, A.S. 1963. Sperm-egg association and its relationship to the activation of the egg in salmonid fishes. J. Embryol. Exp. Morph. 11: 13-33.
34. Ginzburg, A.S. 1972. Fertilization in fishes and the problem of polyspermy. Israel Program for Scientific Translations Jerusalem 1972. Keter Press - Jerusalem.
35. Götting, K.J. 1965. Die Feinstruktur der Hüllschichten reifender Oocyten von Agonus cataphractus L. (Teleostii, Agonidae). Z. Zellforsch. mikrosk. Anat. 66: 405-414.
36. Götting, K.J. 1966. Zur Feinstruktur der Oocyten mariner Teleosteer. Helgoländer Wiss. Meeresunters. 13: 118-170.
37. Gray, J. 1920. The relation of the animal cell to electrolytes. I. A physiological study of the egg of the trout. J. Physiol. 53: 308-
38. Graybill, J.R., and H.F. Horton. 1969. Limited fertilization of steelhead trout eggs with cryo-preserved sperm. J. Fish. Res. Bd. Can. 26: 1400-1404.
39. Hagenmaier, H.E. 1973. The hatching process in fish embryos. III. The structure, polysaccharide and protein cytochemistry of the chorion of the trout egg, Salmo gairdneri (Rich). Acta Histochem. 47(1): 61-69.
40. Hagenmaier, H.E., and R. Wilhelm. 1972. Der Aufbau der Eihülle und ihre Veränderungen während der Keimesentwicklung bei der Forelle (Salmo trutta fario L.) Experientia 28: 605-607.

41. Hagström, B.F., and S. Lönning. 1968. EM studies of unfertilized and fertilized eggs from marine teleosts. Sarsia 33: 73-80.
42. Hart, N.H., and M. Messina. 1972. Artificial insemination of ripe eggs in the Zebra fish, Brachydanio rerio. Copeia 1972. 2: 302-305.
43. Hayes, F.R. 1949. The growth, general chemistry and temperature relations of salmonid eggs. Quart. Rev. Biol. 24: 281-308.
44. Hepworth, W.G. 1972. Study of the preservation of fish sperm by freezing. Wyoming Fish and Game Comm. Completion Rept. FW-3-R-9.
45. Hettler, W.F., Jr. 1970. Effect of paternal death on sperm visibility in the orangethroat darter. Prog. Fish Cult. 32: 209-211.
46. Hines, R., and A. Yashuov. 1971. Some environmental factors influencing the activity of spermatozoa of Mugil capito, a grey mullet. J. Fish Biol. 3: 123-128.
47. His, W. 1873. Untersuchungen über das Ei und die Eientwicklung bei Knochenfischen. Leipzig, F.C.W. Vogel.
48. Hisaoka, K.K. 1958. Microscopic studies of the teleost chorion. Trans. Am. Microsc. Soc. 77: 240-
49. Hodgins, H.F., and G.J. Ridgeway. 1964. Recovery of viable salmon spermatozoa after fast freezing. Prog. Fish Cult. 25: 95.

50. Hoyle, R.J., and Idler, D.P. 1968. Preliminary results in the fertilization of eggs with frozen sperm of Atlantic salmon (Salmo salar). J. Fish. Res. Bd. Can. 25: 1295-1297.
51. Huet, M., and J.A. Timmermans. 1973. Textbook of fish culture: Breeding and cultivation of fish. Fishing News, London U.K. 436 pp.
52. Huver, C.W. 1963. A chemical technique for dechorionating teleost eggs. Copeia 3: 591-592.
53. Ivankov, V.N., and V.P. Kurdyayeva. 1973. Systematic differences and the ecological importance of the membranes in fish eggs. J. Ichthyol. 13: 864-873.
54. Iwamatsu, T. 1965. A further study on fertilizability of fish oocytes. Annotnes Zool. Jap. 35: 190-197.
55. Iwamatsu, T. 1962. Structural changes in the egg surface after fertilization in the fish Oryzias latipes. Annotnes Zool. Jap. 41: 145-153.
56. Iwamatsu, T. 1965. Effect of acetone on the cortical changes at fertilization of the eggs of the medaka, Oryzias latipes. Embryologia 9: 1-12.
57. Kalber, F.A. 1973. Techniques for measuring osmotic and ionic concentrations in fish eggs and larva. Proc. of workshop on egg, larval and juvenile stages of fish in Atlantic Coast Estuaries. NOAA Tech. Pub. #1: p. 36.

58. Kinne, O., and Kinne E.M. 1962. Rates of development in embryos of a cyprinodont fish exposed to different temperature-salinity-oxygen combinations. Can. J. Zool. 40: 231-253.
59. Kokhonoskaya, E.M., V.A. Teodorovich, and I.A. Sadov, 1970. Bactericidal water sterilizer for fish roe incubators. Comm. Fish. Abst. 24: 33.
60. Kossman, H. 1973. Study of the preservation of carp sperm (Cyprinus carpio). Arch. Fischereiwiss. 24: 125-128.
61. Kubota, M., and K. Hagiya, and S. Kimura. 1971. Amino acid composition and solubilization of fish egg shell membrane. J. Tokyo Univ. Fish. 57: 87-94.
62. Kucherova, F.N. 1972. Cation composition of the sperm cells and seminal fluid of Vimba vimba L. J. Ichthyol. 12: 666-670.
63. Kudo, S. 1967. EM observations on the cortical changes in the egg of Carassius carassius. I. The release of granules. Sci. Rep. Tohoku Univ. Ser. 4 33: 185-195.
64. Kusa, M. 1958. Explosion of isolated cortical alveoli of the stickleback egg. Annot. Zool. Jap. 31: 212-215.
65. Kravilashvili, I.S., and V.P. Bozhkova, and L.M. Chailakhyan. 1971. Periodic changes in resistance and membrane potential accompanying cleavage in egg of the loach Misgurnus fossilis. Sov. J. Dev. Biol. 2: 344-348.

66. Kwain, W. 1975. Embryonic development, early growth, and meristic variation in rainbow trout (Salmo gairdneri) exposed to combinations of light intensity and temperature. J. Fish. Res. Bd. Can. 32: 347-402.
67. Lasker, R., and R.H. Tenaza, and L.C. Chamberlain. 1972. The response of Salton Sea fish eggs and larvae to salinity stress. Calif. Fish and Game 58: 58-66.
68. Loeb, J. 1915. On the nature of the conditions which determine or prevent the entrance of the spermatozoa into the egg. Am. Nat. 49: 257.
69. Loeffler, C.A., and S. Løvtrup. 1970. Water balance in the salmon egg. J. Exp. Biol. 52: 291-298.
70. Lønning, S. 1972. Comparative electron microscopic studies of teleostean eggs with special reference to the chorion. Sarsia 49: 41-48.
71. Lowman, F.G. 1953. Electron microscope studies of silver salmon spermatozoa (Oncorhynchus kisutch Walbaum) Exp. Cell. Res. 5: 335-360.
72. Maddock, B.G. 1974. A technique to prolong the incubation period of brown trout ova. Prog. Fish. Cult. 36: 219-222.
73. Maitland, P.S. 1967. The artificial fertilization and rearing of the eggs of Coregonus clupeoides Lacepede. Proc. Roy. Soc. Edinb. (B) 70: 82-106.

74. Makeyeva, A.P., and B.V. Verigin. 1971. Use of the method of pituitary injections in the propagation of silver carp and grass carp. J. Ichthyol. 11: 174-185.
75. Masamichi, Y., and K. Yamagami. 1975. Electron microscopic studies on choriolysis by the hatching enzyme of the teleost, Oryzias latipes. Dev. Biol. 43: 313-321.
76. May, R.C. 1975. Effects of temperature and salinity on fertilization, embryonic development and hatching in Bairdiella icistia (Pisces: Sciaenidae), and the effect of parental salinity acclimation on embryonic and larval salinity tolerance. Fishery Bull. 73: 1-22.
77. Middaugh, D.P., and R.L. Yoakum. 1974. The use of chorionic gonadotropin to induce laboratory spawning of the Atlantic Croaker, Micropogon undulatus, with notes on subsequent embryonic development. Chesapeake Science 15: 110-114.
78. Miller, R.W. 1972. Three methods for determining dissolved oxygen concentrations near fish embryos. Prog. Fish Cult. 34: 39-42.
79. Mitchum, R.W. 1963. Study of the preservation of fish sperm by freezing. Wyoming Game Fish Completion Rept. FW-3-R-10 33-37.
80. Mounib, M.S., and J.S. Eisan. 1969. Alanine and aspartate aminotransferases in eggs and sperm of fish. Life Sci. 8: 531-534.

81. Mounib, M.S., and P.C. Huang, and D.R. Idler. 1968. Cryogenic preservation of Atlantic cod. (Gadus morhua) sperm. J. Fish. Res. Bd. Can. 25: 2623-2632.
82. Müller, H. 1961. EM Untersuchungen zur Bildung der Eihüllen bei Zahnkarpfen. Verh. Dtsch. Ges. 294-306.
83. Müller, H., and G. Sterba. 1963. EM Untersuchungen über Bildung und Struktur der Eihüllen bei Knochenfischen. II. Die Eihüllen jüngerer und älterer Oozyten von Cynolebias belotti Steindachner (Cyprinodontidae). Zool. Jb. Anat. 80: 469-488.
84. Müller, J. 1854. Ueber zahlreiche Porenkanäle in der Eikapsel der Fische. Müller Arch. Anat. Physiol. 1: 186-
85. Nakano, E. 1956. Changes in the egg membrane of the fish egg during fertilization. Embryologia 3: 89-103.
86. Newman, H.H. 1905. On some factors governing the permeability of the egg membrane by the sperm. Biol. Bull. 9: 378.
87. Ogino, C., and S. Tasada. 1962. Changes in inorganic constituents of developing rainbow trout eggs. Bull. Jap. Soc. Scient. Fish. 28: 788-791.
88. Ohtsuka, E. 1960. On the hardening of the chorion of the fish egg after fertilization. The mechanism of chorion hardening in Ozyzias latipes. Biol. Bull. 118: 120-128.

89. Ott, A.G., and H.F. Horton, 1971. Fertilization of chinook and coho salmon egg with cryo-preserved sperm. J. Fish. Res. Bd. Can. 28: 745-748.
90. Pandey, S., and W.S. Hoar. 1972. Induction of ovulation in goldfish by clomiphene citrate. Can. J. Zool. 50: 1679-1680.
91. Plosila, D.S., and W.T. Keller. 1974. Effects of quantity of stored sperm and water on fertilization of brook trout eggs. Prog. Fish Cult. 36: 42-45.
92. Plosila, D.S., and W.T. Keller, and T.J. McCartney. 1972. Effects of sperm storage and dilution on fertilization of brook trout eggs. Prog. Fish Cult. 34: 179-181.
93. Poon, D.C., and A.K. Johnson. 1970. The effect of delayed fertilization on transported salmon eggs. (Oncorhynchus gorbusha). Prog. Fish Cult. 32: 81-84.
94. Post, G., D.V. Power, and T.M. Kloppel. 1974. Survival of rainbow trout eggs often receiving physical shocks of known magnitude. Trans. Am. Fish. Soc. 103: 711-716.
95. Price, J.W. 1943. A device for observing living fish embryos at controlled temperatures. Ohio J. Sci. 43(2): 83-85.
96. Reichert, K.B. 1856. Ueber die Micropyle der Fischeier und über einen bisher unbekanntem, eigentümlichen Bau des Nahrungsdotters reifer und befruchteter Fischeier (Hecht). Müllers Arch. Anat. Physiol. 83-124.

97. Retzins, G. 1912. Zur Kenntnis der Hüllen und besonders des Follikelepitels an den Eieren der Wirbeltiere.  
Biol. Untersuch. Neue Folg. 17: 1-52.
98. Rothschild, L. 1958. Fertilization in fish and lampreys.  
Biol. Rev. Camb. Phil. Soc. 33: 372-392.
99. Rucker, R.R., and J.F. Conrad, and C.W. Dickerson. 1960.  
Ovarian fluid: its role in fertilization. Prog. Fish Cult. 22: 77-78.
100. Sadov, I.A. 1963. Structure and formation of egg membranes in Acipenseridae and in some representatives of Teleostei.  
Trud. Inst. Morf. Zhiv. 38: 110-188.
101. Sadov, I.A. 1958. Concerning the development of the oocyte covering in the Russian, starred and small sturgeon.  
Biol. Sci. 120: 361-364.
102. Salisbury, G.W., and R.G. Hart. 1975. Functional integrity of spermatozoa after storage. Bioscience 25(3): 159-165.
103. Siegel, G. 1958. Zur Morphologie der Eihüllen südamerikanischer Zahnkarpfen. Wiss. Z. Fr. Schiller Univ. Jena, Nat. wiss. R. Heft 2/3: 229-231.
104. Sneed, K.E., and H.P. Clemens. 1956. Survival of fish sperm after freezing and storage at low temperatures. Prog. Fish Cult. 18: 99-103.
105. Stahl, A., and C. Leroy. 1961. L'ovogénèse chez les poissons teleostéens. I. Origine et signification de la zona

- radiata et de ses annexes. Arch. Anat. Micr. Morph. Exp. 50: 251-268.
106. Sterba, G. 1957. Zur Differenzierung der Eihüllen bei Knochenfischen. Z. Zellforsch. 46: 717-728.
107. Sterba, G. 1958. Die Eihüllen des Schmerlen-Eies (Nemachilus barbatula L.) Z. Mikr. Anat. Forsch. 63: 581-588.
108. Sterba, G. Ueber die Struktur der Eihüllen bei einigen Knochenfischen. Naturwiss. 44: 187-
109. Sterba, G., and H. Franke. 1959. Zur EM Struktur der Corticalmembrane der Knochenfischeier. Naturwiss. 46: 93.
110. Sterba, G., and H. Müller. 1962. EM Untersuchungen über Bildung und Struktur der Eihüllen bei Knochenfischen. I Die Hüllen junger Oocyten von Cynolebius belotti Steindachner (Cyprinodontidae). Zool. Jb. Anat. Ontog. 80: 65-80.
111. Strawn, K., and C. Hubbs. 1956. Observations on stripping small fishes for experimental purposes. Copeia 2: 114-116.
112. Szöllösi, D., and R. Billard. 1974. The micropyle of trout eggs and its reaction to different incubation media. J. de Microsc. 21: 55-62.
113. Tegner, M.J., and D. Epel. 1973. Sea urchin sperm-egg interactions studied with the scanning EM. Science 179: 685-688.
114. Thomas, A.E. 1975. Effect of egg concentration in an incubation channel on survival of chinook salmon fry. Trans. Am. Fish. Soc. 104: 335-337.

115. Timoshina, L.A. 1972. Embryonic development of the rainbow trout (Salmo gairdneri irideus Gibb) at different temperatures. J. Ichthyol. 12: 425-432.
116. Truscott, B., and D.R. Idler, and R.J. Hoyle, and H.C. Freeman. 1968. Subzero preservation of Atlantic salmon sperm. J. Fish. Res. Bd. Can. 25: 363-372.
117. Tsukahara, J. 1971. Ultrastructural study on the attaching filaments and villi of the oocyte of Oryzias latipes during oogenesis. Dev. Growth Differ. 13: 173-180.
118. Turdakov, A.F. 1971. The effect of temperature conditions on the speed and fertilizing capacity of the spermatozoa of some ISSYK-KUL fishes. J. Ichthyol. 11: 206-215.
119. Turdakov, A.F. 1971. Rate of sperm motion in body fluid diluted with a varying amount of water observed for three species of Teleostei fish. Sov. J. Dev. Biol. 2: 230-235.
120. Turdakov, A.F., and N.A. Aminova. 1973. A study of the heat resistance of the spermatozoa in a number of species of teleost fish. J. Ichthyol. 13: 198-204.
121. Uwa, H. 1967. A study on the relationship between sperm penetration and egg activation in the Medaka, Oryzias latipes. J. Fac. Sci. Shinushu Univ. 2: 87-94.
122. von Riehl, R., and K.J. Götting. 1974. Structure and occurrence of the micropyle in egg cells and eggs of teleost fishes. Arch. Hydrobiol. 74: 393-402.

123. von Westernhagen, H. 1968. Experiments in rearing the eggs of the haddock (Melanogrammus aeglefinus L.) under combined salinity and temperature conditions. Ber. Dent. Wiss. Komm. Meeresforsch. 19: 270-287.
124. Weisel, G.F. 1948. Relation of salinity to the activity of the spermatozoa of Gillichthys, a marine teleost. Physiol. Zool. 21: 40-48.
125. Werner, W.H.R. 1934. A note on the use of physiological saline as defined herein as an aid in the artificial spawning of speckled trout. Trans. Am. Fish. Soc. 64th Ann. Mtg., Montreal. 346-350.
126. Withler, F.C., and R.M. Humphreys. 1967. Duration of fertility of ova and sperm of sockeye (Oncorhynchus nerka) and pink O. gorbuscha salmon. J. Fish. Res. Bd. Can. 24: 1573-1578.
127. Withler, F.C., and R.B. Morley. 1968. Effects of chilled storage on viability of stored ova and sperm of sockeye and pink salmon. J. Fish. Res. Bd. Can. 25: 2695-2699.
128. Yamamoto, K., and Y. Nagahama. 1966. A method to induce artificial spawning of goldfish all through the year. Bull. Jap. Soc. Sci. Fish. 32: 977-983.
129. Yamamoto, T. 1954. Cortical changes in eggs of the goldfish (Carassius auratus) and the pond smelt (Hypomesus olidus) at the time of fertilization and activation. Jap. J. Ichthyol. 3: 162-170.

130. Yamamoto, T.S. 1957. Some experiments in the chemical changes in the membrane of salmon eggs occurring at the time of activation. Jap. J. Ichth. 6: 54-58.
131. Yamamoto, T.S. 1957. Some morphological and physiological aspects of the eggs of teleostean fishes. J. Fac. Sci. Hokkaido Univ. 13: 484-488.
132. Yamamoto, T.S. 1958. Biochemical property of the membrane of the herring egg, with special reference to the role of the micropyle in fertilization. J. Fac. Sci. Hokkaido Univ. Ser. 6, Vol. 14: 9-16.
133. Yamamoto, T.S. 1961. Physiology of fertilization in fish eggs. Intl. Rev. Cytol. 12: 361-405.
134. Yawagimachi, R. 1957. Studies of fertilization in Clupea pailasi. I. Extension of fertilizable life of the unfertilized eggs by means of isotonic Ringer's solution. Zool. Mag. 66: 218-22.
135. Yoshida, T., and M. Nomura. 1972. A substance enhancing sperm motility in the ovarian fluid of rainbow trout. Bull. Jap. Soc. Sci. Fish. 38: 1073.
136. Young, E.G., and W.R. Inman. 1938. The protein of the casing of salmon eggs. J. Biol. Chem. 124: 189-193.
137. Zirges, M.H., and L.D. Curtis. 1972. Viability of fall chinook salmon eggs spawned and fertilized 24 hours after death of the female. Prog. Fish Cult. 34(4): 190.

138. Zotin, A.I. 1953. Changes occurring in egg membranes of Acipenseridae during development. C.R. Acad. Sci. URSS 92: 443-446.
139. Zotin, A.I. 1953. Initial stages of hardening in egg membranes of salmonid fishes. C.R. Acad. Sci. URSS 89: 573-576.
140. Zotin, A.I. 1958. The mechanism of hardening of the salmonid egg membrane after fertilization or spontaneous activation. J. Embryol. Exp. Morphol. 6: 546-568.

C. Fundulus Embryology and Reproduction

1. Anderson, E. 1970. Cortical alveoli formation and vitellogenesis during oocyte differentiation in the pipe fish Syngnathus fuscus and the killifish, Fundulus heteroclitus. J. Morph. 125: 23-59.
2. Armstrong, P.B. 1928. The antagonism between acetic acid and the chlorides of sodium, potassium, and calcium as manifested in developing Fundulus embryos. J. Gen. Physiol. 11: 515.
3. Armstrong, P.B. 1936. Mechanism of hatching in Fundulus heteroclitus. Biol. Bull. 71: 407.
4. Armstrong, P.B., and J.S. Child. 1965. Stages in the normal development of Fundulus heteroclitus. Biol. Bull. 128: 143-168.
5. Armstrong, P.B. 1931. Functional reactions in the embryonic heart accompanying the ingrowth and development at the vagus innervation. J. Exp. Zool. 58: 43-67.
6. Barnes, L.J. 1953. A further study of centrifugation and low temperature on the development of Fundulus heteroclitus. Biol. Bull. 105: 370.
7. Bennett, M.V.L., and M.E. Spira, and G.D. Pappas. 1972. Properties of electrotonic junctions between embryonic cells of Fundulus. Dev. Biol. 29: 419-435.

8. Boyd, J.F., and R.C. Simmonds. 1974. Continuous laboratory production of fertile Fundulus heteroclitus (Walbaum) eggs lacking chorionic fibrils. J. Fish. Biol. 6: 389-394.
9. Boyd, M. 1928. A comparison of the oxygen consumption of unfertilized and fertilized eggs of Fundulus heteroclitus. Biol. Bull. 55: 92.
10. Brown, O.H. 1905. The permeability of the membranes of the egg of Fundulus heteroclitus. Am. J. Physiol. 14: 354.
11. Brummett, A.R. 1954. The relationships of the germ ring to the formation of the tail bud in Fundulus as demonstrated by the carbon marking technique. J. Exp. Zool. 125: 447-485.
12. Brummett, A.R. 1966. Observations on the eggs and breeding season of Fundulus heteroclitus at Beaufort, North Carolina. Copeia 1966, 616-620.
13. Brummett, A.R. 1968. Deletion-Transplantation experiments on embryos of Fundulus heteroclitus. I. The posterior embryonic shield. J. Exp. Zool. 169: 315-333.
14. Brummett, A.R. 1969. Deletion-Transplantation experiments on embryos of Fundulus heteroclitus. J. Exp. Zool. 172: 443-463.
15. Buno, W., and R. Chambers. 1947. Direct determinations of internal hydrostatic pressure of Fundulus egg. Biol. Bull. 93: 190.

16. Burger, J.W. 1939. Some experiments on the relation of the external environment to the spermatogenic cycle in Fundulus heteroclitus. Biol. Bull. 77: 96-103.
17. Chambers, R., and E.L. Chambers, and C.T. Kao. 1951. The internal hydrostatic pressure of the unfertilized activated Fundulus egg exposed to various experimental conditions. Biol. Bull. 101: 206-207.
18. Chen, T.R. 1970. Fish chromosomes prep.: air dried displays of cultured ovarian cells in 2 species of killifishes (Fundulus) J. Fish. Res. Bd. Can. 27: 158-161.
19. Chen, T.R., and F.H. Ruddle. 1970. A chromosome study of four species and a hybrid of the killifish genus Fundulus (Cyprinodontidae). Chromosoma (Berl.) 29: 255-267.
20. Coghill, G.E. 1933. Somatic myogenic action in embryos of Fundulus heteroclitus. Proc. Soc. Exp. Biol. Med., 31: 62-64.
21. Costello, D.P., M.E. Davidson, A. Eggers, M.H. Fox, and C. Hanley. 1957. Methods for obtaining and handling marine eggs and embryos. Marine Biological Laboratory, Woods Hole, Mass.
22. Crawford, R.B., and C.E. Wilde. 1970. Temporal relationships of protein synthesis to morphogenesis in the Fundulus embryo. Bull. Mt. Desert Isl. Biol. Lab. 1970.
23. Crawford, R.B., F.J. Hendler, and C.E. Wilde. 1967. Protein synthesis in embryos of Fundulus heteroclitus. Bull. Mt. Desert Isl. Biol. Lab. 7: 14-15.

24. Crawford, R.B., C.E. Witte Jr., M.H. Heinemann, and J.J. Hendler.  
Morphogenetic disturbances from timed inhibitions of  
protein synthesis in Fundulus. J. Emb. and Exp. Morph.  
29: 363-382.
25. Crawford, R.B., M.H. Heinemann, and C.E. Wilde, Jr. 1968.  
Conditions of inhibition of morphogenesis and macromole-  
cular syntheses in Fundulus heteroclitus embryos. Bull. Mt.  
Desert Isl. Bio. Lab. 8: 13-15.
26. Eisler, R., and G. LaRoche. 1972. Elemental composition of the  
estuarine teleost Fundulus heteroclitus (L.) J. Exp. Mar.  
Biol. Ecol. 9: 29-42.
27. Forsthoefel, P. 1951. The occurrence of twining in Fundulus  
heteroclitus after centrifugation. Biol. Bull. 101:  
221-222.
28. Gabriel, M.L. 1942. The effect of temperature on vertebral numbers  
in Fundulus. The collecting Net 17(5): 85-86.
29. Gabriel, M.L. 1944. Factors affecting the number and form of  
vertebra in Fundulus heteroclitus. J. Exp. Zool. 95(1):  
105-147.
30. Garside, E.T., and C.M. Jordan. 1968. Upper lethal temperatures  
at various levels of salinity in the euryhaline cyprinodonts  
Fundulus heteroclitus and F. diaphanus after isosmotic  
acclimation. J. Fish. Res. Bd. Can. 25: 2717-2720.
31. Gee, W.P. 1916. Effects of acute alcoholization on the germ  
cells of Fundulus heteroclitus. Biol. Bull. 31: 379-406.

32. Glaser, O. 1929. Temperature and heart rate in Fundulus embryos. Brit. J. Exp. Biol. 6: 325-339.
33. Hinrichs, A.M. 1925. Modification of development on the basis of differential susceptibility to radiation. I. Fundulus and U.V. radiation. J. Morph. 41: 239-265.
34. Hoadley, L. 1928. On the localization of developmental potencies in the embryo of Fundulus heteroclitus. J. Exp. Zool. 52: 7-44.
35. Huver, Charles W. 1960. The stage at fertilization of the egg of Fundulus heteroclitus. Biol. Bull. 119: 320.
36. Joseph, E.B., and V.P. Saksena. 1966. Determination of salinity tolerances in mummichog (Fundulus heteroclitus) larvae obtained from hormone-induced spawning. Chesapeake Sci. 7: 193-197.
37. Kagan, Benjamin M. 1935. The fertilizable period of the eggs of Fundulus heteroclitus and some associated phenomena. Biol. Bull. 69: 185-201.
38. Kaighn, M.E. 1964. A biochemical study of the hatching process in Fundulus heteroclitus. Dev. Biol. 9: 56-80.
39. Kao, C.T., and R. Chambers. 1954. The internal hydrostatic pressure of the Fundulus egg. I. The activated egg. J. Exp. Biol. 31: 139-149.
40. Kao, C.T. et. al. 1954. II. Permeability of the chorion. J. Cell Comp. Physiol. 44: 447-461.

41. Kellicott, W.E. 1916. The effect of low temperature upon the development of Fundulus. Am. J. Anat. 20: 449.
42. Kemp, N.E., and M.D. Allen. 1956. Electron microscopic observations on changes in the cortical cytoplasm after fertilization of Fundulus eggs. Biol. Bull. 111: 305.
43. Kemp, N.E. 1956. EM observations on the development of the chorion of Fundulus. Biol. Bull. 111: 293.
44. Kemp, N.E., and M.D. Allen. 1956. Electron microscopy of growing oocytes of Fundulus. Anat. Rec. 124: 460-461.
45. Kemp, N.E., and E. Hibbard. 1957. Protoplasmic bridges between follicle cells and developing oocytes of Fundulus heteroclitus. Biol. Bull. 113: 329.
46. Kessel, R.G. 1960. The role of cell division in gastrulation of Fundulus heteroclitus. Exptl. Cell. Res. 20: 277-282.
47. Lentz, T.L., and J.P. Trinkaus. 1967. A fine structural study of cytodifferentiation during cleavage, blastula and gastrula stages of Fundulus heteroclitus. J. Cell Biol. 32: 127-138.
48. Lewis, W.H. 1912. Experiments on localization and regeneration in the embryonic shield and germ ring of a teleost fish. (F. heteroclitus). Anat. Rec. 6: 325-333.
49. Lufts, B., G.E. Pickford, and J.W. Atz. 1968. The effects of low temperature, and cortisol on testicular regression in the hypophsectomized cyprinodont fish Fundulus heteroclitus. Biol. Bull. 134: 74-86.

50. Marza, V.D., and E.V. Marza, and M.J. Guthrie. 1937. Histochemistry of the ovary of Fundulus heteroclitus with special reference to the differentiating oocytes. Biol. Bull. 73: 67.
51. Matthews, S.A. 1938. The seasonal cycle in the gonads of Fundulus. Biol. Bull. 75: 66-74.
52. Matthews, S.A. 1939. The relationship between the pituitary gland and the gonads in Fundulus. Biol. Bull. 76: 241.
53. Matthews, S.A. 1939. The effects of light and temperature on the male sexual cycle in Fundulus. Biol. Bull. 77: 92-95.
54. Milkman, R. 1954. Controlled observation of hatching in F. heteroclitus. Biol. Bull. 107: 300.
55. Mizejewski, G.J., and G.M. Ramm. 1968. Carbon particle uptake by macrophages in the ovary and thyroid of Fundulus during various seasonal cycles. Chesapeake Sci. 9: 233-237.
56. Moore, C.R. 1917. On the capacity for fertilization after the initiation of development. Biol. Bull. 33: 258.
57. Newman, H.H. 1907. Spawning behavior and sexual dimorphism in Fundulus heteroclitus and allied fish. Biol. Bull. 12: 314.
58. Newman, H.H. 1908. Process of heredity as exhibited by development of Fundulus hybrids. J. Exp. Zool. 5: 503-561.

59. Nicholas, J.S. 1927. The application of experimental methods to the study of developing Fundulus embryos. Proc. Nat. Acad. Sci. 13: 695-698.
60. Oppenheimer, J.M. 1934. Experiments on early developing stages of Fundulus. Proc. Nat. Acad. Sci. 20: 536-538.
61. Oppenheimer, J.M. 1936. Processes of localization in developing Fundulus. J. Exp. Zool. 73: 405-444.
62. Oppenheimer, J.M. 1936. Transplantation experiments on developing teleosts (Fundulus and Perca). J. Exp. Zool. 72: 409-437.
63. Oppenheimer, J.M. 1936. Localized vital staining of teleostean embryos. Science 82: 598.
64. Oppenheimer, J.M. 1936. The development of isolated blastoderms of Fundulus heteroclitus. J. Exp. Zool. 72: 247-269.
65. Oppenheimer, J.M. 1941. The anatomical relationships of abnormally located Mauthner's cells in Fundulus embryos. J. Comp. Neurol. 74: 131-167.
66. Oppenheimer, J.M. 1946. A case of a typical twinning in Fundulus heteroclitus. Anat. Rec. 95: 67-71.
67. Oppenheimer, J.M. 1947. Organization of the teleost blastoderm. Quart. Rev. Biol. 22: 105.
68. Oppenheimer, J.M. 1950. Functional regulation in Fundulus heteroclitus embryos with abnormal central nervous systems. J. Exp. Zool. 115: 471-492.

69. Oppenheimer, J.M. 1955. The differentiation of derivatives of the lower germ layers in Fundulus following the implantation of shield grafts. J. Exp. Zool. 128: 525-560.
70. Oppenheimer, J.M. 1955. Ectopic differentiation of ears in secondary embryos of Fundulus. Proc. Nat. Acad. Sci. 41: 680-684.
71. Oppenheimer, J.M. 1959. Extraembryonic transplantation of sections of the Fundulus embryonic shield. J. Exp. Zool. 140: 247-267.
72. Oppenheimer, J.M. 1964. The development of isolated Fundulus embryonic shields in salt solution. Acta. Embryol. Morph. Exp. 7: 143-154.
73. Pickford, G.E., et al. 1969. Studies on the blood serum of the euryhaline cyprinodont fish, Fundulus heteroclitus, adapted to fresh or to salt water. Trans. Conn. Acad. Arts. Sci. 43: 25-70.
74. Pickford, G.E., B. Lofts, G. Bara, and J.W. Atz. 1972. Testis stimulation in hypophysectomized male killifish, Fundulus heteroclitus, treated with mammalian growth hormone and/or luteinizing hormone. Biol. Reprod. 7: 370-386.
75. Potts, W.T.W., and D.H. Evans. 1967. Sodium and chloride balance in the killifish, Fundulus heteroclitus. Biol. Bull. 113: 411-475.

76. Richards, A., and R.P. Porter. 1935. II. The mitotic index in preneuronal tube stages of Fundulus heteroclitus. Am. J. Anat. 56: 365-393.
77. Richards, S.W. 1966. Comparison of post-larvae and juveniles of Fundulus heteroclitus and F. majoris. Trans. Am. Fish. Soc. 95: 218-226.
78. Rugh, R. 1954. The effect of various levels of x-irradiation on the gametes and early embryos of Fundulus heteroclitus. Biol. Bull. 107: 319-320.
79. Schwartz, R.J., and C.E. Wilde, Jr. 1973. Changes in protein synthesis in the morphogenesis of Fundulus heteroclitus. Nature: 245: 376-379.
80. Scott, G.G., and W.E. Kellicott. 1916. The consumption of oxygen during the development of Fundulus heteroclitus. Anat. Rec. 11: 531.
81. Shanklin, P.F. 1954. Evidence for active transport in Fundulus embryos. Biol. Bull. 107: 320.
82. Shanklin, D.R. 1959. Studies on the Fundulus chorion. J. Cell Comp. Physiol. 53: 1-12.
83. Shanklin, P.R., and P.B. Armstrong. 1952. The osmotic behavior and anatomy of the Fundulus chorion. Biol. Bull., 103: 295.
84. Shaver, J.R. 1951. The occurrence of twinning in Fundulus heteroclitus after centrifugation. Biol. Bull. 101: 201.

85. Shaver, J.R., and S. Ito. 1952. The effect of centrifugation and low temperature on development of Fundulus heteroclitus. Biol. Bull. 103: 309.
86. Shaver, J.R. 1953. The development of isolated blastomeres of Fundulus heteroclitus. Biol. Bull. 105: 383.
87. Slicher, Anna M. 1961. Endocrinological and hematological studies in Fundulus heteroclitus (Linn.). Bull. of Bingham Oceanographic Collection. 17(3): 55.
88. Solberg, A.N. 1938. The susceptibility of Fundulus heteroclitus embryos to x-irradiation. J. Exp. Zool. 78: 441-469.
89. Spitz, L.M., and T.B. Burnett. The tyrosinases of Fundulus heteroclitus at different stages of embryonic development. J. Embryol. Exp. Morph. 19: 1-8.
90. Stockard, C.R. 1906. The development of Fundulus heteroclitus in solutions of lithium chloride, with appendix on its development in fresh water. J. Exp. Zool. 3: 99.
91. Stockard, C.R. 1907. The influence of external factors, chemical and physical, on the development of Fundulus heteroclitus. J. Exp. Zool. 4: 165.
92. Trinkaus, J.P. 1948. Properties of the surface coat in embryos of Fundulus heteroclitus. Biol. Bull. 95: 271.
93. Trinkaus, J.P. 1949. The surface gel layer of Fundulus eggs in relation to epiboly. Proc. Nat. Acad. Sci. U.S.A. 35: 218.

94. Trinkaus, J.P. 1951. A study of the mechanism of epiboly in the eggs of Fundulus heteroclitus. J. Exp. Zool. 118: 269-319.
95. Trinkaus, J.P. 1963. The cellular basis of Fundulus epiboly. Adhesivity of blastula and gastrula cells in culture. Dev. Biol. 7: 513-532.
96. Trinkaus, J.P. 1967. Fundulus. In Methods in Developmental Biology. pp. 113-122. F.H. Wilt, N.K. Wessells, eds. T.Y. Crowell, N.Y.
97. Trinkaus, J.P., and J.W. Drake. 1956. Exogenous control of morphogenesis in isolated Fundulus blastomeres by nutrient chemical factors. J. Exp. Zool. 132: 311.
98. Trinkaus, J.P. 1973. Role of the periblast in Fundulus epiboly. Sov. J. Dev. Biol. 2: 326-328.
99. Umminger, Bruce L. 1970. Physiological studies on supercooled killifish (Fundulus heteroclitus). II. Serum organic constituents and the problem of supercooling. J. Exp. Zool.
100. Wessells, Norman K., and F.J. Swartz. 1953. Relation of the micropyle to cortical changes at fertilization in the egg of Fundulus heteroclitus. Anat. Rec. 117: 557.
101. Wilhelmi, A.E., G.E. Pickford, and W.H. Sawyer. 1955. Initiation of the spawning reflex response in Fundulus by the administration of fish and mammalian neurohypophysial preparations and synthetic oxytocin. Endocrinology 57: 243-252.

102. Williams, G. Jr., R.C. Simmonds, and J.F. Boyd. 1975.  
Effects of the antimicrobial furanace on Fundulus hetero-  
clitus embryos. J. Fish. Res. Bd. Can. 32: 69-71.
103. Wyman, L.C. 1924. The reactions of the melanophores of  
embryonic and larval Fundulus heteroclitus to certain  
chemical substances. J. Exp. Zool. 40: 161-180.

## II. OVA PRESERVATION TESTS

## A. Ficoll-Dextran Solution.

Methods.

Stock solutions of Ficoll<sup>1</sup> (2.5 gm/100 ml) and Dextran<sup>2</sup> (5.0 gm/100 ml) were made up in 10<sup>0</sup>/00, 20<sup>0</sup>/00, and 35<sup>0</sup>/00 Instant Ocean.

These stock solutions were used to prepare 5 ml Ficoll-Dextran test solutions with the following composition.

- a) Full strength Dextran
- b) F:D = 1:1
- c) F:D = 2:1
- d) F:D = 5:1
- e) F:D = 10:1
- f) full strength Ficoll

Six batches of the above test solutions were made up from each of the stock solutions at 10<sup>0</sup>/00, 20<sup>0</sup>/00, 35<sup>0</sup>/00, for 24, 48, and 72 hr tests to be run at both 4C and 20C. All control tests were carried out at the appropriate salinity.

Ova to be tested were stripped, pooled, and divided into four roughly equal lots and placed in 10 ml plastic petri dishes. One batch was used as a control group and was fertilized immediately. 5 ml quantities of one of the F:D test solutions were added to the other 3 dishes. These were placed in a temperature chamber at 20C and one of the dishes removed at 24, 48, and 72 hr intervals and the ova tested with fresh sperm. The above procedure was repeated at 4C.

Results.

The results of the tests are shown in Tables 1, 2, and 3. In no case tested did the presence of Ficoll or Dextran extend the fertilizable period of the ova.

Table 1: Effect of F:D solutions in  $10^0/00$  sea water on the fertilizability of Fundulus heteroclitus ova.

F:D Ratio	Time	20C		4C	
		# ova	# fertilized	# ova	# fertilized
F:D = 0:1	C	9	9	16	15
	24	14	0	10	0
	48	14	0	17	0
	72	9	0	13	0
F:D = 1:1	C	6	6	19	18
	24	15	0	20	0
	48	7	0	8	0
	72	9	1	14	0
F:D = 2:1	C	19	13	12	10
	24	9	0	24	0
	48	14	0	17	0
	72	42	0	17	0
F:D = 5:1	C	12	11	19	19
	24	13	0	16	0
	48	21	0	10	0
	72	10	0	12	0
F:D = 10:1	C	29	29	9	9
	24	28	0	8	0
	48	31	0	6	0
	72	16	0	6	0
F:D = 1:0	C	39	38	11	11
	24	21	0	18	0
	48	16	0	16	0
	72	17	0	20	0

Table 2: Effect of F:D solutions in 20<sup>0</sup>/100 sea water on the fertilizability of Fundulus heteroclitus ova.

F:D Ratio	Time	20C		4C	
		# ova	# fertilized	# ova	# fertilized
F:D = 0:1	C	12	11	12	11
	24	9	0	37	0
	48	10	0	22	0
	72	14	0	22	0
F:D = 1:1	C	16	16	16	16
	24	12	0	22	0
	48	18	0	23	0
	72	23	0	33	0
F:D = 2:1	C	16	16	16	16
	24	22	0	37	0
	48	24	0	38	0
	72	26	0	47	0
F:D = 5:1	C	16	16	16	16
	24	10	0	10	0
	48	12	0	8	0
	72	18	0	11	0
F:D = 10:1	C	13	13	13	13
	24	12	0	6	0
	48	10	0	6	0
	72	11	0	5	0
F:D = 1:0	C	17	17	17	17
	24	7	0	15	0
	48	11	0	12	0
	72	16	0	12	0

Table 3: Effect of F:D solutions in 35<sup>0</sup>/<sub>00</sub> sea water on the fertilizability of Fundulus heteroclitus ova.

F:D Ratio	Time	20C		4C	
		# ova	# fertilized	# ova	# fertilized
F:D = 0:1	C	18	9	6	6
	24	13	0	13	0
	48	26	0	9	0
	72	28	0	10	0
F:D = 1:1	C	11	8	24	22
	24	11	0	25	0
	48	10	0	23	0
	72	10	0	26	0
F:D = 2:1	C	9	1	11	10
	24	6	0	10	0
	48	5	0	10	0
	72	5	0	10	0
F:D = 5:1	C	19	7	21	19
	24	23	0	18	0
	48	21	0	16	0
	72	18	0	15	0
F:D = 10:1	C	28	20	23	19
	24	25	0	19	0
	48	25	0	20	0
	72	29	0	19	0
F:D = 1:0	C	19	19	14	14
	24	26	0	14	0
	48	25	0	15	0
	72	28	0	9	0

### B. Holtfreter's Solution.

#### Methods.

Stock solutions of 100<sup>0</sup>/<sub>0</sub>, 50<sup>0</sup>/<sub>0</sub>, 25<sup>0</sup>/<sub>0</sub>, and 10<sup>0</sup>/<sub>0</sub> Holtfreter's solution were prepared by dilution with distilled water. Freshly stripped ova were divided into 5 lots, one to act as control which was immediately fertilized, the other 4 lots placed in finger bowls with the test Holtfreter's solutions. The test lots were incubated at 22C for 48 hr and then tested with fresh sperm.

#### Results.

	# ova	# fertilized
Control	25	23
100 <sup>0</sup> / <sub>0</sub> Holtfreter's	22	0
50 <sup>0</sup> / <sub>0</sub> "	27	0
25 <sup>0</sup> / <sub>0</sub> "	24	0
10 <sup>0</sup> / <sub>0</sub> "	20	0

### C. Sucrose Solution.

#### Methods.

A 2M sucrose solution was prepared in distilled water. Freshly stripped ova were placed in 5 ml aliquots of this solution and

stored at 20C and 4C. At 24, 48, 72 hr intervals batches of ova were tested for fertilizability with fresh sperm.

<u>Results.</u>			
	Time	# ova	# fertilized
20C	24	26	0
	48	11	0
	72	-	-
4C	24	9	0
	48	15	0
	72	14	0

#### D. Dry Preservation.

##### Methods.

Freshly stripped ova were placed in small plastic capsules (Beem, Inc.) capable of holding approx. 150 ova. All tests were conducted with the ova in a dry state. No additional supportive media were added other than the ovarian fluid which accompanies the ova when stripped from the female.

The capsules were capped and placed in a Lauda-Brinkman K-2/R circulating water bath. The temperature range of 0-30C was tested at two degree intervals. Ova were tested for fertilizability at intervals of 6, 12, 24, 48, 72, 96 hr. at each temperature. For

testing, ova were removed from the water bath, placed in a Petri dish and fresh sperm was expressed onto them, after which they were covered with 20<sup>0</sup>/<sub>00</sub> sea water and allowed to incubate with the sperm for 5-10 minutes. The ova were then rinsed and checked 2-3 hr. later for fertilization, at which time they were at the 4-8 cell stage. Each batch was subsequently monitored until hatching, with abortions and abnormalities recorded. Each test was replicated at least 4 times.

#### Results.

Percent fertilization (with standard deviation) at the times and temperatures tested are shown in Table 4 and summarized in Fig. 1. Percent abnormality and abortion are given in Table 5.

#### E. Salinity.

##### Methods.

Stock solutions of Instant Ocean ranging from 0-35<sup>0</sup>/<sub>00</sub> at 5<sup>0</sup>/<sub>00</sub> increments were prepared. Ova were stripped into 100 ml Petri dishes containing about 20 ml of the test sea water. Batches were kept at 22C and tested at 1, 3, 6, 9, 12 and 15 hr intervals with fresh sperm.

Similar tests were also run at 12C. Fresh ova were placed in small plastic capsules which were then filled with water of a test salinity and stored in a water bath. Ova were checked for fertilizability at intervals of 6, 12, 24, 48, 72 hrs.

Table 4. Percent fertilization of temperature-treated ova after varying exposure times over the range of 0-30°C. The values represent the means.

Temp. °C	Exposure Time (Hours)							
	6	12	24	48	72	96	120	168
0	0							
2	0							
4	42.9 <sup>±</sup> 4 (a)	0	0	0	0	0		
6	65.9 <sup>±</sup> 7	55.1 <sup>±</sup> 11	26.7 <sup>±</sup> 14	0	0	0		
8	90.6 <sup>±</sup> 2	82.6 <sup>±</sup> 4	44.5 <sup>±</sup> 9	24.1 <sup>±</sup>	8.4 <sup>±</sup> 3	0		
10	95.0 <sup>±</sup> 4	94.7 <sup>±</sup> 5	94.3 <sup>±</sup> 2	76.4 <sup>±</sup> 13	67.4 <sup>±</sup> 13	20.3 <sup>±</sup> 7	0	0
12	98.0 <sup>±</sup> 3	96.7 <sup>±</sup> 3	95.9 <sup>±</sup> 5	78.0 <sup>±</sup> 4	64.1 <sup>±</sup> 8	46.7 <sup>±</sup> 2	42.2 <sup>±</sup> 8	14.7 <sup>±</sup> 9
14	96.3 <sup>±</sup> 4	92.7 <sup>±</sup> 5	88.7 <sup>±</sup> 10	84.9 <sup>±</sup> 11	7.8 <sup>±</sup> 2	0		
16	98.6 <sup>±</sup> 2	95.9 <sup>±</sup> 4	25.3 <sup>±</sup> 7	6.2 <sup>±</sup> 4	0	0		
18	90.5 <sup>±</sup> 3	84.3 <sup>±</sup> 4	21.7 <sup>±</sup> 12	1.2 <sup>±</sup> 4	0	0		
20	94.6 <sup>±</sup> 9	91.0 <sup>±</sup> 3	24.3 <sup>±</sup> 8	5.8 <sup>±</sup> 3	0	0		
22	72.8 <sup>±</sup> 4	69.0 <sup>±</sup> 4	12.0 <sup>±</sup> 8	0	0	0		
24	67.0 <sup>±</sup> 5	40.0 <sup>±</sup> 8	2.7 <sup>±</sup> 3	0	0	0		

Table 4 (continued)

Temp. °C	6	12	24	48	72	96	120	168
26	55.7 <sup>±</sup> 6	46.0 <sup>±</sup> 8	3.6 <sup>±</sup> 5	0	0	0		
28	31.5 <sup>±</sup> 6	25.4 <sup>±</sup> 7	1.3 <sup>±</sup> 2	0	0	0		
30	22.0 <sup>±</sup> 3	16.9 <sup>±</sup> 5	0	0	0	0		

(a) Standard deviations have been rounded to the nearest whole number.

FIGURE 1. The effect of temperature on the fertilizability of ova exposed over the range of 0-30C. Each fertilization percent represents the mean of four replicate tests.




Table 5 Percentage abortion of embryos from temperature-treated ova.

Temp. °C	Exposure Time (Hcurs)					
	6	12	24	48	72	96
4	87.2(a) - (b)	-	-	-	-	-
6	65.0	90.0	56.0	-	-	-
8	20.4	42.7	77.5	71.4	87.0	-
10	12.0	23.0	19.6	50.4	63.0	74.0
12	10.0	7.3	8.0	37.7	22.8	50.0
14	6.8	4.1	14.0	22.0	22.2	-
16	5.0	3.0	26.6	42.8	-	-
18	14.8	12.0	7.6	0	-	-
20	0	2.0	57.9	40.0	-	-
22	8.3	11.1	46.1	-	-	-
24	37.5	44.2	100.0	-	-	-
26	60.0	54.5	77.0	-	-	-
28	10.3	38.0	65.0	-	-	-
30	33.3	50.0	-	-	-	-

(a) Each percent represents mortalities and those embryos which exhibited structural deformities.

(b) Fertilization did not occur.

Percent fertility, abnormality and abortion were monitored as in Section D.

#### Results.

Percent fertilization at the salinities, times, and temperatures tested are shown in Tables 6, 7 and summarized in Figs. 2, 3.

Abnormality and abortion rates are shown in Tables 8, 9.

#### F. pH

##### Methods.

Batches of ova were treated at 22C in sea water of two salinities, 10<sup>0</sup>/00 and 22<sup>0</sup>/00, the pH of which was adjusted to some value between 6.0 and 9.4. Standard buffers of pH 4.01 and pH 10.0 were used to adjust the pH of the test medium to the desired value. Ova were stripped into 100 ml Petri dishes containing 20 ml of the test solution, and tested at intervals of 1, 3, 6, 9, 12, 15 hr. exposure. Ova were also treated at 12C, in 10<sup>0</sup>/00 salinity at various pH values.

##### Results.

Percent fertilization is shown in Tables 10-13 and summarized in Figs. 4-7 . Abnormality and abortion rates are shown in Tables 14-15.

Table 6. Percent fertilization of salinity-treated ova after varying exposure times over the range of 0-35‰/oo at ambient temperature (22-25°C).

		Exposure Time (Hours)					
		1	3	6	9	12	15
SALINITY ‰	0	0	0	0	0	0	0
	5	78.2 <sup>±</sup> 8(a)	67.3 <sup>±</sup> 15	5.2 <sup>±</sup> 2	1.2 <sup>±</sup> 2	1.5 <sup>±</sup> 2	0
	10	96.0 <sup>±</sup> 4	88.5 <sup>±</sup> 4	64.9 <sup>±</sup> 9	25.6 <sup>±</sup> 8	14.0 <sup>±</sup> 6	4.6 <sup>±</sup> 1
	15	49.7 <sup>±</sup> 7	24.6 <sup>±</sup> 6	25.0 <sup>±</sup> 6	9.7 <sup>±</sup> 2	3.6 <sup>±</sup> 4	5.0 <sup>±</sup> 1
	20	35.7 <sup>±</sup> 15	23.3 <sup>±</sup> 12	5.1 <sup>±</sup> 2	1.7 <sup>±</sup> 2	1.2 <sup>±</sup> 2	0
	25	10.2 <sup>±</sup> 3	4.5 <sup>±</sup> 6	2.1 <sup>±</sup> 2	0	0	0
	30	2.3 <sup>±</sup> 2	0	0	0	0	0
	35	0	0	0	0	0	0

(a) Values represent the means. Standard deviation have been rounded to the nearest whole number.

Table 7. Fertilization percentages of ova exposed to a 0-35‰ salinity range at 12°C over varying time periods. The pH of sea water was adjusted to 8.0-8.4 with pH 10.00 buffer.

		Exposure Time (Hours)				
		6	12	24	48	72
SALINITY (‰)	0	0	0	0	0	0
	5	88.7 <sup>±</sup> 7(a)	70.5 <sup>±</sup> 7	38.6 <sup>±</sup> 9	4.1 <sup>±</sup> 5	0
	10	95.3 <sup>±</sup> 2	88.4 <sup>±</sup> 7	66.5 <sup>±</sup> 7	10.4 <sup>±</sup> 3	0
	15	24.6 <sup>±</sup> 6	20.0 <sup>±</sup> 5	8.1 <sup>±</sup> 4	1.0 <sup>±</sup> 1	0
	20	12.5 <sup>±</sup> 3	1.7 <sup>±</sup> 2	0	0	0
	25	15.7 <sup>±</sup> 5	0	0	0	0
	30	0	0	0	0	0
35	0	0	0	0	0	

(a) Values represent the means. Standard deviations have been rounded to the nearest whole number.

FIGURE 2. The effect of salinity on the fertilizability of ova exposed over the range of 0-35<sup>0</sup>/<sub>00</sub> at ambient temperature (22-25C). Each fertilization percent represents the mean of four replicate tests.




FIGURE 3. Percent fertilization of ova exposed to the salinity range of 0-25<sup>0</sup>/<sub>00</sub> at 12C. pH of artificial sea water was adjusted to 8.0-8.4.




Table 8 Percentage abortion of embryos resulting from salinity exposed ova maintained at ambient temperature, 22-25°C.

		Exposure Time (Minutes)					
		1	3	6	9	12	15
SALINITY ‰	5	0	4.2(a)	0	0	0	-(b)
	10	0	0	8.7	4.7	0	0
	15	2.7	0	3.8	14.0	0	0
	20	2.0	0	12.4	0	0	-
	25	3.4	0	0	-	-	-
	30	0	0	-	-	-	-

Table 9 Percentage abortion of embryos resulting from salinity exposed ova maintained at 12°C.

		Exposure Time (Hours)			
		6	12	24	48
SALINITY ‰	5	8.3(a)	7	16	100
	10	5.0	3.1	20.4	84
	15	2.1	6.3	34	100
	20	20.4	0	0	0
	25	10	0	0	0

(a) Values represent total mortalities and structurally deformed embryos.

(b) Fertilization did not occur.

Table 10 Percentage fertilization of ova exposed to 10<sup>0</sup>/oo sea water adjusted to the pH range 6.0-9.5. Tests were conducted at ambient temperature (22-25°C). Values represent the mean.

	Exposure Time (Hours)					
	3	6	9	12	15	24
6.0	28.5 <sup>±</sup> 6(a)	1.25 <sup>±</sup> 2	0	0	0	0
6.5	23.8 <sup>±</sup> 4	2.4 <sup>±</sup> 2	0	0	0	0
6.8	42.7 <sup>±</sup> 17	15.9 <sup>±</sup> 6	5.5 <sup>±</sup> 4	2.3 <sup>±</sup> 3	0	0
7.0	61.4 <sup>±</sup> 25	19.2 <sup>±</sup> 8	4.6 <sup>±</sup> 3	0	0	0
7.4	96.5 <sup>±</sup> 3	76.0 <sup>±</sup> 6	29.7 <sup>±</sup> 10	11.3 <sup>±</sup> 4	0	0
pH 7.8	96.5 <sup>±</sup> 4	73.9 <sup>±</sup> 3	27.4 <sup>±</sup> 8	7.9 <sup>±</sup> 3	3.7 <sup>±</sup> 3	0
8.2	97.5 <sup>±</sup> 5	92.1 <sup>±</sup> 6	50.8 <sup>±</sup> 16	42.6 <sup>±</sup> 15	8.3 <sup>±</sup> 3	0
8.6	96.9 <sup>±</sup> 4	83.6 <sup>±</sup> 11	46.6 <sup>±</sup> 11	12.5 <sup>±</sup> 3	0	1.4 <sup>±</sup> 2
9.0	94.1 <sup>±</sup> 4	53.9 <sup>±</sup> 14	37.3 <sup>±</sup> 7	0	0	0
9.2	92.6 <sup>±</sup> 6	45.4 <sup>±</sup> 7	25.1 <sup>±</sup> 6	0	0	0
9.5	0	0	0	0	0	0

(a) Standard deviations have been rounded to the nearest whole number.

Table (11) Percent Fertilization of ova exposed to a salinity of 10‰ at 12°C over the pH range 6.0-9.5. Values represent the means.

pH	Exposure Time (Hours)		
	24	48	72
6.0	10.6 <sup>±</sup> 2(a)	0	0
6.5	32.0 <sup>±</sup> 4	11.8 <sup>±</sup> 8	0
6.8	31.3 <sup>±</sup> 6	6.6 <sup>±</sup> 2	0
7.0	47.5 <sup>±</sup> 7	11.3 <sup>±</sup> 8	0
7.4	72.0 <sup>±</sup> 6	9.8 <sup>±</sup> 5	0
7.8	66.7 <sup>±</sup> 6	10.7 <sup>±</sup> 4	0
8.2	51.5 <sup>±</sup> 8	12.2 <sup>±</sup> 9	1.6 <sup>±</sup> 1
8.6	39.1 <sup>±</sup> 13	5.4 <sup>±</sup> 4	1.1 <sup>±</sup> 1
9.0	57.8 <sup>±</sup> 15	9.8 <sup>±</sup> 3	0
9.2	54.1 <sup>±</sup> 9	11.5 <sup>±</sup> 9	0
9.5	0		

(a) Standard deviations have been rounded to nearest whole number.

Table 12 Percent fertilization of ova exposed to 22<sup>o</sup>/oo sea water at 12<sup>o</sup>C over the pH range 6.5-9.5. Values represent the means.

	Exposure Time (Hours)		
	6	12	24
6.5	0	0	0
6.8	0	0	0
7.0	3.3 <sup>±</sup> 1 (a)	0	0
pH 7.4	9.7 <sup>±</sup> 2	8.9 <sup>±</sup> 4	0
7.8	11.9 <sup>±</sup> 2	7.8 <sup>±</sup> 3	0
8.2	13.1 <sup>±</sup> 3	7.7 <sup>±</sup> 3	0
8.6	8.1 <sup>±</sup> 2	5.0 <sup>±</sup> 2	0
9.0	8.2 <sup>±</sup> 2	2.6 <sup>±</sup> 2	0
9.2	3.1 <sup>±</sup> 2	0	0
9.5	0	0	0

(a) Standard deviations have been rounded to nearest whole number.

Table (13) Percentage fertilization of ova exposed to 22<sup>0</sup>/oo sea water adjusted to the pH range 6.0-9.5. Tests were conducted at ambient temperature (22-25<sup>0</sup>C). Values represent the mean.

		Exposure Time (Hours)			
		1	3	6	9
pH	6.0	0	0		0
	6.5	3.5 <sup>±</sup> 1(a)	0	0	0
	6.8	0	0	0	0
	7.0	5.9 <sup>±</sup> 1	1.7 <sup>±</sup> 2	0	0
	7.4	16.2 <sup>±</sup> 5	3.9 <sup>±</sup> 2	0.6 <sup>±</sup> 1	0
	7.8	21.3 <sup>±</sup> 1	8.0 <sup>±</sup> 2	0	0
	8.2	24.3 <sup>±</sup> 4	16.6 <sup>±</sup> 2	2.0 <sup>±</sup> 1	0
	8.6	17.5 <sup>±</sup> 3	3.5 <sup>±</sup> 2	0	0
	9.0	10.6 <sup>±</sup> 2	8.6 <sup>±</sup> 4	0	0
	9.2	2.6 <sup>±</sup> 1	0	0	0
	9.5	0	0	0	0

(a) Standard deviations have been rounded to the nearest whole number.

FIGURE 4. Percent fertilization of ova treated at  $10^0/00$  over the pH range of 6.0-9.5 at 22-25C.




FIGURE 5. Percent fertilization of ova treated at 22<sup>0</sup>/00 over the pH range 6.0-9.4, at 22-25C.




FIGURE 6. Percent fertilization of ova exposed to  $10^0/00$  at 12C over the pH range of 6.0-9.5.




FIGURE 7. Percent fertilization of ova exposed to 22<sup>0</sup>/<sub>00</sub> at 12C  
over the pH range 6.4-9.4.




Table 14 Percent abortion of embryos from pH-treated ova.  
 A. Ova exposed to 10<sup>0</sup>/oo sea water at 22-25<sup>0</sup>C.  
 B. Ova exposed to 22<sup>0</sup>/oo sea water at 22-25<sup>0</sup>C.

		Exposure Time (Hours)				
A.		3	6	9	12	15
pH	6.0	21.8(a)	100	-(b)	-	-
	6.5	38.4	40	-	-	-
	6.8	9.2	19.0	25	50	-
	7.0	10.5	21.0	45.4	-	-
	7.4	4.3	3.5	9.0	21	33
	7.8	4.0	3.5	4.3	14.2	-
	8.2	3.2	3.3	15.5	8.1	50
	8.6	6.1	5.7	10.4	11.1	33
	9.0	7.3	10.4	18.1	-	-
	9.2	14.5	20.0	10.0	-	-

		Exposure Time (Hours)		
B.		1	3	6
pH	6.5	100(a)	-(b)	-
	7.0	37	50	100
	7.4	31.3	100	-
	7.8	27	66	-
	8.2	24	81.2	66.6
	8.6	26	81.8	-
	9.0	85.7	33.3	-

(a) Values represent total mortalities and embryos exhibiting structural deformities.

(b) Fertilization did not occur.

Table 15 Percentage abortion of embryos from ova exposed to two salinities at the same temperature and pH range.

A. Ova exposed to 10<sup>0</sup>/oo at 12C.

B. Ova exposed to 22<sup>0</sup>/oo at 12C.

		Exposure Time (Hours)		
		24	48	72
A.	pH 6.0	70.5 (a)	- (b)	-
	6.5	30.4	75.0	-
	6.8	24.0	56.0	-
	7.0	27.7	30.5	-
	7.4	9.7	50.0	-
	7.8	12.5	22.7	-
	8.2	21.8	16.6	66.0
	8.6	17.0	28.5	0
	9.0	27.0	49.0	-
	9.2	48.0	42.0	-
		<hr/>		
B.	pH 7.4	10.0 (a)	30.7	- (b)
	7.8	0	30.0	-
	8.2	9.3	11.1	-
	8.6	21.0	28.5	-
	9.0	29.0	66.0	-
	9.2	39.0	-	-

(a) Values represent total abnormalities and embryos which exhibited structural deformities.

(b) No fertilization occurred.

## G. Analysis of Ovarian Fluids

### 1. pH.

#### Methods.

- a) Ovarian fluid accompanying stripped ova was drawn into 50 $\mu$ l capillary tubes.
- b) The tubes were sealed at both ends to prevent further changes in pH due to absorbance of CO<sub>2</sub> from the air.
- c) The fluid was analyzed using a BMS3 MK2 Blood Microsystem Apparatus.
- d) The fluid from 10 females was measured and a mean value obtained.

#### Results.

- a) The ovarian fluid averaged pH  $8.0 \pm 0.4$ .

### 2. Elemental Composition.

#### Methods.

- a) Ovarian fluid accompanying stripped ova was pooled and stored in a frozen state until about 0.5 ml had accumulated.
- b) Blood was also collected from the caudal artery for comparison with ovarian fluid.

Results.

Table 16.

µgm/gm sample

	<u>Na</u>	<u>Cl</u>	<u>K</u>	<u>Ca</u>	<u>Mg</u>
Blood	2480 (3.2%)*	3800 (6.1%)	592 (4%)	821 (25%)	200 (10%)
Ovarian Fluid	3860 (3.0%)	5350 (6.0%)	374 (11%)	217 (25%)	116 (10%)

\* Values in parentheses indicate the estimated error within the measurement. Na, Cl, K analyses were performed by instrumental neutron activation analysis (i.e. reactor neutron irradiation followed by gamma ray spectrometry). Ca, Mg results obtained by atomic absorption flame photometry.

### III. SPERM PRESERVATION TESTS

#### A. Ficoll<sup>1</sup>-Dextran<sup>2</sup> Solution

##### Methods.

- 1) Stock solutions of Ficoll and Dextran were prepared as in II-A.
- 2) The same F:D ratios were used as in II-A.
- 3) Quantities of pooled sperm were incubated with the various F/D solutions at both 4C and 20C for 48 and 72 hr and then tested with fresh ova.

##### Results.

Table 17 : Fertility of Fundulus heteroclitus sperm incubated in various Ficoll-Dextran solutions in 35<sup>0</sup>/<sub>00</sub> sea water, at 4C for 48 hr.

F:D Ratio	# ova	# fertilized
0:1	16	0
1:1	16	0
2:1	19	0
5:1	17	0
10:1	13	0
1:0	10	0

<sup>1</sup>Ficoll: Pharmacia Chemicals - Lot No. 2300

<sup>2</sup>Dextran: General Biochemicals - Lot No. 41923B

Table 18: Fertility of Fundulus heteroclitus sperm incubated in various Ficoll-Dextran solutions in 20<sup>0</sup>/00 sea water, at 4C for 72 hr.

F:D Ratio	# ova	# fertilized
0:1	21	0
1:1	27	0
2:1	21	0
5:1	18	0
10:1	24	0
1:0	19	0

Table 19: Fertility of Fundulus heteroclitus sperm incubated in various Ficoll-Dextran solutions in 35<sup>0</sup>/00 sea water at 20C for 48 hr.

F:D Ratio	# ova	# fertilized
0:1	6	0
1:1	4	0
2:1	5	0
5:1	5	0
10:1	17	0
1:0	17	0

B. Preliminary Tests on Sperm, Temperature, pH, Salinity, Extender  
Solution Interactions.

Methods.

A large number of single tests were performed to investigate the role of various parameters on sperm preservation in an attempt to find possible favorable areas for more definitive experimentation.

Results.

Table 20: Role of Various Parameters on Preservation of Sperm  
Fertility.

	Conditions	Time (hr)	Temp. (°C)	# ova	# fert.	% fert.
1)	dry-mixed w/20 <sup>0</sup> / <sub>00</sub> SW prior to exposing to ova	24	4	28	0	0
2)	dry-mixed w/stock pH7 buffer prior to exposing to ova	24	4	33	7	21
3).	dry-mixed w/pH7 buffer diluted 10 X prior to exposing to ova	24	4	49	19	39
4)	as in (3) - diluted 100 X	24	4	39	6	15
5)	incubated in pH7 solution	24	4	43	29	67
6)	incubated in pH7 solution	24	8	54	46	85
7)	dry-add pH7 prior to exposure to ova	24	8	65	55	85
8)	incubated in pH7 solution	48	8	103	0	0
9)	" " " "	48	4	79	48	61
10)	" " " "	72	8	42	18	43
11)	" " " "	72	4	102	9	9

	Conditions	Time (hr)	Temp. (°C)	# ova	# fert.	% fert.
12)	incubated in pH7 solution	24	8	64	51	80
13)	incubated in 0.05 M glycine	24	8	112	0	0
14)	incubated in pH7 - $\frac{\text{sperm}}{\text{buffer}} = \frac{1}{1}$	72	4	40	0	0
15)	" " " " = $\frac{1}{10}$	72	4	45	7	16
16)	as in (14)	72	2	53	0	0
17)	as in (15)	72	2	52	1	2
18)	w/pH7 and 1.25 mM ATP (Sigma Chemicals)	48	2	60	49	82
19)	" "	72	2	54	0	0
20)	" "	48	4	48	4	8
21)	" "	72	4	42	0	0
22)	w/pH 7.5	72	2	45	25	56
23)	w/pH 8.0	72	2	44	0	0
24)	w/pH 8.5	72	2	44	0	0
25)	w/pH 7.5	72	4	45	12	27
26)	w/pH 8.0	72	4	59	0	0
27)	w/pH 8.5	72	4	42	0	0
28)	w/pH 7.5	24	4	44	44	100
29)	w/pH 8.0	24	4	51	0	0
30)	w/pH 7.5	48	4	40	21	53
31)	w/pH 7.5	48	2	23	22	96
32)	w/pH 7.5	72	2	34	11	32
33)	pH 7.5 + 1.25 mM ATP + O <sub>2</sub>	72	2	143	130	91
34)	" " " "	72	2	124	120	97

C. Effect of Buffer, ATP, Temperature, Oxygen and Salinity of Post-Preservation Mixing Medium on Sperm Fertility.

Methods.

1. Sperm were incubated for 72 hr at 2C in a variety of media:
  - a) sperm : pH 7.5 buffer = 1:1
  - b) sperm : pH 7.5 buffer w/ 1.25 mM ATP = 1:1
  - c) " : " " " + O<sub>2</sub> = 1:1
  - d) " " " + O<sub>2</sub> = 1:1
2. All preservation was carried out in sealed plastic capsules capable of holding 0.5 ml total solution.
3. Following incubation sperm was expressed onto fresh ova bathed in the following solutions.
  - a) 10 ‰ sea water
  - b) 20 ‰ sea water
  - c) 35 ‰ sea water
  - d) pH 7 buffer
  - e) pH 7.5 buffer

Results

(see Table 21)

Table 21: Effect of Incubation Media and Mixing Media on Sperm  
Preserved 72 hr at 2C.

Incubation Medium		MIXING MEDIA					TOTALS
		10°/00	20°/00	35°/00	pH 7	pH 7.5	
pH 7.5	#ova	40	41	36	38	36	$\frac{191}{4} = 2\%$
	#fert	0	1	0	3	0	
pH 7.5 1.25 mM ATP	#ova	24	26	27	25	35	$\frac{137}{3} = 2\%$
	#fert	1	0	0	2	0	
pH 7.5 O <sub>2</sub>	#ova	26	17	22	25	26	$\frac{116}{18} = 16\%$
	#fert	7	8	2	0	1	
pH 7.5 O <sub>2</sub>	#ova	18	23	18	18	13	$\frac{90}{62} = 69\%$
	#fert	18	22	14	5	3	
TOTALS		$\frac{26}{108}$ =24%	$\frac{32}{107}$ =30%	$\frac{16}{103}$ =16%	$\frac{10}{106}$ = 9%	$\frac{4}{110}$ = 4%	

D. Effect of ATP Concentration.

Methods.

- Sperm were incubated in the following solutions of ATP in pH 7.5 buffer, at 10C.
  - 1.25 mM ATP
  - 1.0 "
  - 2.0 "
  - 5.0 "

e) 10.0 mM ATP

f) 0.0 " , i.e. pH 7.5 buffer only

2. Sperm were tested after 72 and 94 hrs with fresh ova.

### Results.

Table 22 : Effect of ATP concentration on sperm fertility.

ATP CONC.	72 hr		94 hr	
	# ova	# fertilized	# ova	# fertilized
1.0 mM	48	5	65	6
1.25	57	22	63	0
2.0	39	38	78	0
5.0	39	5	75	0
10.0	46	23	55	0
pH 7.5 only	49	0	58	0

E. Temperature.

### Methods.

1. Sperm were stripped and placed in small plastic capsules in a dry state.

2. Capsules were capped and placed in a Lauda-Brinkman K-2/R circulating water bath.

3. The temperature range tested was 0-20C at 2 degree increments.

4. At intervals of 2, 4, 6, 12, 15 and 24 hr, the sperm were removed from the water bath and mixed with fresh ova.
5. After 5-10 min. the ova were rinsed and later checked for evidence of fertilization.

#### Results.

The relationship between temperature, time of preservation and fertility are given in Table 23 and summarized in Fig. 8. The mortality - abnormality rates resulting from these tests are given in Table 24.

#### F. Salinity.

##### Methods.

1. The salinity range tested was the same as that used for the ova (Sect. II E).
2. Sperm : Sea water = 1:10.
3. Sperm were tested at 25C at intervals of 5, 10, 20, 30, 40 min.
4. Sperm were tested at 2C at intervals of 1, 3, 6, 9 hr.

##### Results.

The relationship between salinity and time of preservation at 25C is given in Table 25 and summarized in Fig. 9 , and at 2C is given in Table 26 and summarized in Fig. 10.

Mortality and abnormality rates from these tests are given in Tables 27 and 28.

Table 23 Percent of fresh ova fertilized with sperm exposed to the range of 0-20°C over various time intervals.

Temp. °C	Exposure Time (Hours)							
	2	4	6	12	15	24	48	72
0	81.5± 10(a)	60.2± 13	19.2± 9	2.0± 1	0	0		
2	95.6± 4	91.0± 6	38.5± 18	13.1± 5	0	25.6± 33	63.9± 49	36.5± 40
4	77.3± 10	26.1± 7	17.6± 13	0	0	0		
6	25.2± 10	13.1± 4	5.3± 2	0	0	0		
8	28.0± 3	14.2± 9	2.4± 1	0	0	0		
10	22.7± 7	12.1± 4	0	0	0			
12	24.7± 9	6.8± 3	0	0	0			
14	28.8± 13	9.7± 5	0	0	0			
16	30.4± 16	8.3± 4	0	0	0			
18	13.3± 3	10.8± 5	0	0	0			
20	8.4± 4	2.4± 2	0	0	0			

(a) Standard deviations have been rounded to nearest whole number.

FIGURE 8. The effect of temperature on the fertilizing capacity of sperm subjected to the range of 0-20C. Each fertilization percent represents the mean of four replicate tests.




Table (24) Percentage abortion of embryos from temperature-treated sperm.

Temp. °C	2	4	6	12
0	4.0(a)	0	6.3	0
2	0	8.0	0	12.0
4	9.3	5.3	0	-(b)
6	0	5.3	0	-
8	10.3	0	3.7	-
10	2.3	0	-	-
12	4.1	9.7	-	-
14	0	10.0	-	-
16	0	0	-	-
18	0	7.7	-	-
20	6.3	3.2	-	-

(a) Each percent represents mortalities and those embryos which exhibited structural deformities.

(b) Fertilization did not occur.

Table 25 Percent of fresh ova fertilized with sperm exposed to the range of 0-35‰ at various time intervals at ambient temperature (22-25°C).

		Exposure Time (Minutes)				
		5	10	20	30	40
SALINITY ‰	0	0	0	0	0	0
	5	26.3 <sup>±</sup> 11(a)	11.4 <sup>±</sup> 6	3.3 <sup>±</sup> 4	1.6 <sup>±</sup> 1	0
	10	97.2 <sup>±</sup> 4	96.0 <sup>±</sup> 5	50.0 <sup>±</sup> 12	30.0 <sup>±</sup>	7.0 <sup>±</sup> 3
	15	95.9 <sup>±</sup> 5	96.0 <sup>±</sup> 4	17.9 <sup>±</sup> 10	0	0
	20	85.6 <sup>±</sup> 7	86.3 <sup>±</sup> 14	27.0 <sup>±</sup> 20	3.2 <sup>±</sup> 2	0
	25	97.8 <sup>±</sup> 2	89.0 <sup>±</sup> 3	15.1 <sup>±</sup> 12	1.9 <sup>±</sup> 2	0
	30	98.8 <sup>±</sup> 3	95.3 <sup>±</sup> 7	34.0 <sup>±</sup> 12	3.4 <sup>±</sup> 2	0
	35	99.2 <sup>±</sup> 2	99.2 <sup>±</sup> 2	29.8 <sup>±</sup> 7	3.4 <sup>±</sup> 3	0

(a) Values represent the means. Standard deviations have been rounded to the nearest whole number.

FIGURE 9. Percent fertilization of fresh ova fertilized with sperm exposed over the salinity range of 0-35<sup>0</sup>/00 at ambient temperature (four replicate tests).




Table (26) Percent fertilization of fresh ova fertilized with sperm exposed to 0-35‰ sea water at 2°C. The pH of sea water was adjusted with pH 10.00 buffer to 8.0-8.4.

SALINITY (‰)	Exposure Time (Hours)		
	1	3	6
0	0	0	0
5	95.2 <sup>±</sup> 8 (a)	12.3 <sup>±</sup> 7	2.3 <sup>±</sup> 2
10	92.6 <sup>±</sup> 6	15.3 <sup>±</sup> 7	3.7 <sup>±</sup> 2
15	45.5 <sup>±</sup> 13	2.8 <sup>±</sup> 2	0
20	41.4 <sup>±</sup> 8	0	0
25	27 <sup>±</sup> 3	0	0
30	0	0	0
35	0	0	0

(a) Values represent the mean values. Standard deviations have been rounded to nearest whole number.

FIGURE 10. Percent fertilization of fresh ova fertilized with sperm exposed at 2C over the salinity range of 0-30<sup>0</sup>/00. pH of sea water adjusted to 8.0-8.4.




Table (27) Percentage abortion of embryos resulting from salinity exposed sperm maintained at 2°C.

SALINITY ‰	Exposure Time (Hours)		
	1	3	6
5	0	5(a)	0
10	1	0	0
15	0	6.7	-(b)
20	0	-	-
25	9.4	-	-

Table (28) Percentage abortion of embryos resulting from salinity exposed sperm maintained at ambient temperature, 22-25°C.

SALINITY ‰	Exposure Time (Minutes)				
	5	10	20	30	40
0	0	0	0	0	-(b)
10	0	2(a)	0	2.3	1.8
15	2	2	5	0	-
20	0	3	0	12.0	-
25	4	4.2	6.5	0	-
30	4.7	0	6.0	3.4	-
35	5.3	1.7	3.3	0	-

(a) Percentage represent both mortalities and embryos which exhibited structural deformities.

(b) Fertilization did not occur.

## G. pH

Methods.

1. pH levels between 6.0 and 9.4 in 10<sup>0</sup>/<sub>00</sub> and 22<sup>0</sup>/<sub>00</sub> sea water were prepared as in Sect. II F.
2. Sperm:pH-adjusted sea water = 1:10.
3. Preservation tests were run at 25C and checked at 5, 10, 20, 30, 40 min.
4. Sperm were also incubated in pH-adjusted 10<sup>0</sup>/<sub>00</sub> sea water at 2C and checked at 1, 3, 6, 9 hr.

Results.

The role of pH and salinity at 25C on sperm preservation are shown in Tables 29 and 30 and summarized in Figs. 11 and 12, and at 2C in Tables 31 and 32 and Figs. 13 and 14. Mortality-abnormality rates are given in Tables 33 and 34.

## H. Whole Testis Preservation.

Methods.

1. Males were sacrificed and the testes excised, placed dry in small plastic capsules and the capsules capped.
2. The capsules were placed in a water bath at temperatures from -4C to 10C at two degree increments, and stored for 24, 48, 72, 96 hr intervals before testing with fresh ova in 20<sup>0</sup>/<sub>00</sub> sea water.
3. Note: Large robust males with good mature reproductive coloration should be used.

Table (29) Percentage fertilization of fresh ova fertilized with sperm exposed over the pH range 6.0-9.5 at 10<sup>6</sup>/oo. Tests were conducted at ambient temperature (22-25°C).

		Exposure Time (Minutes)			
		10	20	30	40
pH	6.0	50.8 <sup>±</sup> 7.0(a)	40.5 <sup>±</sup> 5	3.4 <sup>±</sup> 4	0
	6.5	91.2 <sup>±</sup> 3	85.0 <sup>±</sup> 14	6.4 <sup>±</sup> 5	0
	6.8	90.7 <sup>±</sup> 11	83.0 <sup>±</sup> 9	12.3 <sup>±</sup> 7	0
	7.0	96.7 <sup>±</sup> 4	84.8 <sup>±</sup> 6	42.5 <sup>±</sup> 10	5.9 <sup>±</sup> 3
	7.4	89.4 <sup>±</sup> 5	36.3 <sup>±</sup> 5	5.4 <sup>±</sup> 3	0
	7.8	83.6 <sup>±</sup> 6	15.4 <sup>±</sup> 7	2.0 <sup>±</sup> 2	0
	8.2	86.5 <sup>±</sup> 5	13.1 <sup>±</sup> 5	0	0
	8.6	47.3 <sup>±</sup> 12	5.4 <sup>±</sup> 2	0	0
	9.0	33.7 <sup>±</sup> 7	0	0	0
	9.2	15.8 <sup>±</sup> 3	0	0	0
9.5	0	0	0	0	

(a) Standard deviations have been rounded to the nearest whole number.

Table 30 Percentage fertilization of fresh ova fertilized with sperm exposed to a pH range of 6.0-9.4 at 22°/oo. Tests were conducted at ambient temperature (22-25°C).

		Exposure Time (Minutes)			
		10	20	30	40
pH	6.0	36.2 <sup>±</sup> 8	27.8 <sup>±</sup> 4	0	0
	6.5	84.2 <sup>±</sup> 4	60.8 <sup>±</sup> 7	10.2 <sup>±</sup> 5	0
	6.8	89.1 <sup>±</sup> 6	65.0 <sup>±</sup> 14	9.0 <sup>±</sup> 5	0
	7.0	86.3 <sup>±</sup> 9	68.7 <sup>±</sup> 10	21.8 <sup>±</sup> 8	0
	7.4	72.0 <sup>±</sup> 11	51.2 <sup>±</sup> 9	12.1 <sup>±</sup> 4	0
	7.8	77.4 <sup>±</sup> 9	29.4 <sup>±</sup> 6	2.3 <sup>±</sup> 2	0
	8.2	62.1 <sup>±</sup> 7	19.0 <sup>±</sup> 9	5.1 <sup>±</sup> 1	0
	8.6	38.6 <sup>±</sup> 7	17.1 <sup>±</sup> 5	0	0
	9.0	22.0 <sup>±</sup> 4	0	0	0
	9.2	4.3 <sup>±</sup> 2	0	0	0
	9.4	0	0	0	0

(a) Standard deviations have been rounded to nearest whole number.

FIGURE 11. Percent fertilization of fresh ova fertilized with sperm held at 10<sup>0</sup>/<sub>00</sub> exposed to a pH range of 6.0-9.5. Tests were carried out at 22-25C.




FIGURE 12. Percent fertilization of fresh ova fertilized with sperm treated at 22<sup>0</sup>/<sub>00</sub> sea water over the pH range of 6.0-9.4, at 22-25C.




Table (31) Percent fertilization of fresh ova fertilized with sperm exposed to a pH range of 6.0-9.5 at 2°C and 10‰ sea water. Values represent the means.

		Exposure Time (Hour)				
pH		1	3	6	9	12
	6.0	82.6± 7.6(a)	42.0± 7.2	6.7± 3.4	0	0
	6.5	82.0± 9.4	69.7± 7.5	42.2± 11.4	4.3± 2.1	0
	6.8	79.9± 7.8	40.5± 13.8	17.3± 7.3	4.0± 2.1	0
	7.0	82.2± 9.6	65.7± 11.7	20.1± 6.3	0	0
	7.4	50.2± 11.1	14.1± 5.0	0	0	0
DH	7.8	19.4± 11.8	0	0	0	
	8.2	23.8± 13.7	3.5± 1.1	0	0	
	8.6	16.5± 5.1	0	0	0	
	9.0	5.4± 2.3	0	0	0	
	9.2	0	0	0	0	
	9.5	0	0	0	0	

(a) Standard deviations have been rounded to nearest whole number.

Table 32 Percent fertilization of fresh ova fertilized with sperm exposed to a pH range of 6.0-9.4 at 22°/oo sea water and 2°C. Values represent the means.

	Exposure Time (Hours)			
	1	3	6	9
6.0	69.3± 4 (a)	0	0	0
6.5	91.3± 8	20.4± 10	4.2± 2	0
6.8	93.1± 7	23.5± 11	2.3± 2	0
7.0	95.4± 5	16.1± 8	1.5± 1	0
pH 7.4	95.2± 3	27.0± 9	3.7± 2	0
7.8	28.1± 7	2.1± 1	0	0
8.2	18.3± 5	4.1± 2	0	0
8.6	10.4± 5	1.8± 1	0	0
9.0	13.2± 4	0	0	0
9.2	0	0	0	0
9.4	0	0	0	0

(a) Standard deviations have been rounded to nearest whole number.

FIGURE 13. Percent fertilization of fresh ova fertilized by sperm exposed to  $10^0/00$  at 2C over the pH range 6.0-9.2.




FIGURE 14. Percent fertilization of fresh ova fertilized by sperm exposed to 22<sup>0</sup>/<sub>00</sub> at 2C over the pH range 6.0-9.2.




Table (33) Percent abortion of embryos from pH-treated sperm.  
 A. Sperm exposed to 100/oo sea water at 22-25°C.  
 B. Sperm exposed to 220/oo sea water at 22-25°C.

		Exposure Time (Minutes)			
		10	20	30	40
A.	6.0	0.	0	0	-(b)
	6.5	0.6(a)	2.4	4.7	-
	6.8	0	3.5	9.0	-
	7.0	0	0	5.2	0
	7.4	0	0	0	-
	pH 7.8	3.5	15.0	0	-
	8.2	4.7	8.9	0	-
	8.6	5.7	13.0	-	-
	9.0	20.3	-	-	-
	9.2	40.7	-	-	-
B.	6.0	8.3	5.3	0	-(b)
	6.5	0	5.7	0	-
	6.8	7.4	4.5	10.0	-
	7.0	3.7	3.0	0	-
	pH 7.4	4.0	0	0	-
	7.8	0	5.3	0	-
	8.2	4.3	2.1	0	-
	8.6	0	9.7	-	-
	9.0	20.0	-	-	-
	9.2	24.3	-	-	-

(a) Values represent total mortalities and embryos exhibiting structural deformities.

(b) Fertilization did not occur.

Table 34 Percentage abortion of embryos resulting from pH-treated sperm.

A. Sperm exposed to 10<sup>0</sup>/oo sea water at 20°C.

B. Sperm exposed to 22<sup>0</sup>/oo sea water at 20°C.

		Exposure Time (Hours)			
		1	3	6	9
A.	pH 6.0	5.1 (a)	0	8.5	- (b)
	6.5	4.0	0	3.7	10
	6.8	5.0	0	7.6	0
	7.0	0	0	0	-
	7.4	8.3	0	-	-
	7.8	0	-	-	-
	8.2	0	0	-	-
	8.6	0	-	-	-
	9.0	0	-	-	-
	B.	pH 6.0	0	- (b)	-
6.5		3 (a)	7	0	-
6.8		0	5.3	0	-
7.0		0	3.8	0	-
7.4		6	9	0	-
7.8		5.5	0	-	-
8.2		0	0	-	-
8.6		0	0	-	-
9.0		0	-	-	-
9.5		-	-	-	-

(a) Values include total mortalities and embryos with structural deformities.

(b) No fertilization recorded.

Results.

Table 35 : Percent fertility of sperm from whole testis preservation.

Time	10C	8C	6C	4C	2C	0C	-2C	-4C
24	86	85	96	90	90	98	92	95
48	96	30	91	89	*	99	99	99
72	4	7	95	60	93**	96	99	46
96	0	19	88	70	72	88	83	26
Total # ova	364	247	661	389	373	467	490	553
# fertilized	241	95	610	297	327	445	466	380
% abnormal	3	3	15	6	2	3	7	33

\* not attempted

\*\* average of 3 replicates

#### IV. DISCUSSION OF OVA AND SPERM PRESERVATION TESTS.

##### A. Ova Preservation

Best results for ova preservation can be obtained by "dry" techniques, using only the small amounts of ovarian fluid that accompanies the ova upon stripping to keep the ova moist. Containers containing the ova should be sealable and watertight to prevent both dessication or entry of extraneous fluids. Ideally, the container should be completely filled with ova, with minimum air-ova interface, again to prevent dessication. A gas permeable material would probably be of advantage, since the exchange of  $O_2$  and  $CO_2$  would be allowed.

The temperature of storage is also fairly critical. In the experiments performed, the range 10-14 C proved to be optimum. In this range ova retained fertilizability of up to 67% after 72 hrs., with recorded fertility up to 168 hrs at 12C. Refinement of technique could probably enhance these values. Abnormality and abortion rates were also minimal within this temperature range.

Techniques using other than dry preservation such as salines, pH-adjusted media, sugar solutions, etc. generally gave unsatisfactory results in terms of long time storage and preservation of fertilizability. Ova tended to water harden, become activated, or perhaps die after exposure to any kind of aqueous medium. Non-aqueous media were not tested, however.

## B. Sperm Preservation

Sperm preservation proved to be more difficult since the process of obtaining the sperm, i.e. stripping, also initiated activity and motility in the sperm. Under normal circumstances the motility is short lived, of the order of minutes. Once the sperm lost their motility they would be unable to effect fertilization. Again, all aqueous media tested proved unsatisfactory for preservation, even those which included some ATP as an energy source.

The best results for stripped sperm preservation were obtained using dry storage, i.e. only the sperm and accompanying spermatid fluids, and low temperatures, the latter acting to slow or inhibit the motility. Temperatures below 6C gave good results, with the best preservation at approx. 2C. The preservation time, however, was only of the order of a few hours, e.g. 13% fertility after 12 hrs, rather than days as with the ova. Hence, preservation of stripped sperm was judged to be unfeasible with respect to the objectives of this study.

Hence it was decided to leave the sperm intact, within the testes, and preserve the testes themselves. This would prevent the activation of the sperm caused by stripping. Excellent results were obtained by preserving whole, excised testes, from robust, mature males. Sperm retained fertilizability of over 90% when stored in the temperature range of 0-4C for 72 hrs, and fertility was still in excess of 70% after 96 hrs.

Thus, whole testis preservation at low temperature is regarded

as the best procedure for maintaining sperm fertility. Abnormality and abortion rates of the resulting embryos were also very minimal using this procedure.

## V. EMBRYO PRESERVATION TESTS.

All tests thus far have dealt with the preservation of individual gametes separately. The following tests study the possibility of using temperature to arrest development of already fertilized zygotes. This would provide an alternative method of studying embryogenesis by delaying the onset of embryogenesis until the desired time.

### Methods.

1. Large batches of at least 500 fresh ova were fertilized with fresh sperm.
2. The resulting zygotes were immediately divided into 11 equal lots in cappable test tubes, one lot to act as a control group and the other 10 to act as test groups.
3. The experimental lots were designated as follows:
  - a) freshly fertilized
  - b) germinal disc stage
  - c) 2-cell stage
  - d) 4-cell stage
  - e) 64-cell stage
  - f) 10 hour post fertilization
  - g) 12 " " "
  - h) 18 " " "
  - i) 24 " " "
  - j) 48 " " "

4. Each lot was allowed to develop at 25C until it reached its designated stage.
5. Upon reaching the designated stage a given lot was placed in a water bath at the test temperature, ranging from 2-12C at 2 degree intervals.
6. At 24, 48, 72 hr intervals 1/3 of the embryos in each tube were removed along with 5 ml of test water, placed in a Petri dish allowed to reach 25C, and monitored for subsequent embryonic development, abortions, abnormalities, and hatching.

Results.

(see Tables 36-41)

Table 36 : Effect of storage at 2C on the embryogenesis of  
*F. heteroclitus* zygotes.

	25C	2 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	20	46	30	30	31	30	50	50	52	30	26
% initial mortality	5	100	97	97	100	63	6	8	6	20	42
% total mortality <sup>3</sup>	15	100	97	97	100	70	22	22	23	30	54
% abnormality	0	-	3	0	-	10	12	4	2	7	0
% hatching	85	-	0	3	-	20	66	74	75	63	46

	25C	2 C - 48 HR									
total # zygotes		50	30	30	30	30	49	50	50	29	25
% initial mortality		100	100	100	100	90	53	76	50	48	52
% total mortality		100	100	100	100	94	76	86	72	70	72
% abnormality		-	-	-	-	3	2	0	2	3	0
% hatching		-	-	-	-	3	22	14	26	27	28

	25C	2 C - 72 HR									
total # zygotes		50	30	31	30	32	56	50	47	31	23
% initial mortality		100	100	100	100	100	93	94	98	74	61
% total mortality		100	100	100	100	100	93	94	98	78	65
% abnormality		-	-	-	-	-	0	4	0	3	5
% hatching		-	-	-	-	-	7	2	2	19	30

<sup>1</sup> Fresh Fertilized

<sup>2</sup> Germinal Disc

<sup>3</sup> First 48 hrs after removal from water bath.

Table 37 : Effect of storage at 4C on the embryogenesis of  
F. heteroclitus zygotes.

	25C	4 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	104	30	23	24	24	25	49	30	40	28	24
% initial mortality <sup>3</sup>	3	33	61	71	71	36	12	4	40	4	13
% total mortality	48	53	74	96	75	56	39	32	60	32	33
% abnormality	7	17	9	0	21	16	2	6	35	0	17
% hatching	45	30	17	4	4	28	59	62	5	68	50

	25C	4 C - 48 HR									
total # zygotes		29	25	25	25	25	29	49	51	24	25
% initial mortality		86	92	100	100	88	72	59	27	42	16
% total mortality		90	92	100	100	88	79	73	55	63	48
% abnormality		3	4	-	-	12	7	0	0	0	0
% hatchin.		7	4	-	-	0	14	27	45	37	52

	25C	4 C - 72 HR									
total # zygotes		28	28	27	25	25	30	50	73	21	24
% initial mortality		82	100	100	100	100	97	88	85	81	50
% total mortality		89	100	100	100	100	97	90	92	90	67
% abnormality		0	-	-	-	-	0	4	0	0	0
% hatching		11	-	-	-	-	3	6	8	10	33

ORIGINAL PAGE IS  
OF POOR QUALITY

Table 38: Effect of storage at 6C on the embryogenesis of F. heteroclitus zygotes.

	25C	6 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	168	52	31	31	30	29	38	39	41	30	37
% initial mortality	0	29	94	97	83	55	3	18	2	7	14
% total mortality	21	37	94	97	83	55	11	28	12	13	16
% abnormality	10	7	0	0	0	14	2	3	8	7	14
% hatching	69	56	6	3	17	31	87	69	80	80	70

	25C	6 C - 48 HR									
total # zygotes		51	31	30	30	31	31	39	38	29	29
% initial mortality		82	100	100	97	54	35	8	8	45	17
% total mortality		92	100	100	100	58	48	21	16	48	28
% abnormality		2	-	-	-	19	4	0	0	0	0
% hatching		6	-	-	-	23	48	79	84	52	72

	25C	6 C - 72 HR									
total # zygotes		46	29	28	31	31	34	42	41	29	26
% initial mortality		85	100	100	100	84	71	36	5	55	19
% total mortality		91	100	100	100	94	88	62	61	66	31
% abnormality		9	-	-	-	6	12	0	2	3	4
% hatching		0	-	-	-	0	0	38	37	31	65

Table 39 : Effect of storage at 8C on the embryogenesis of  
F. heteroclitus zygotes.

	25C	8 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	56	26	49	51	50	49	34	52	21	20	19
% initial mortality	0	19	43	35	38	4	0	2	10	5	0
% total mortality	36	73	90	62	76	92	79	31	100	80	100
% abnormality	0	19	0	14	8	8	0	0	-	0	-
% hatching	64	8	10	4	16	0	21	69	-	20	-

	25C	8 C - 48 HR									
	total # zygotes		48	49	36	51	48	36	51	25	20
% initial mortality		17	84	33	100	13	3	4	8	5	19
% total mortality		85	92	92	100	88	28	27	84	85	67
% abnormality		7	4	2	-	8	0	0	0	5	0
% hatching		8	4	6	-	4	72	73	16	10	33

	25C	8 C -72 HR									
	total # zygotes		54	50	43	47	56	40	48	28	20
% initial mortality		100	100	100	100	46	5	8	21	15	50
% total mortality		100	100	100	100	77	60	52	78	35	67
% abnormality		-	-	-	-	11	2	0	2	0	0
% hatching		-	-	-	-	12	38	48	20	65	33

Table 40 : Effect of storage at 10C on the embryogenesis of  
F. heteroclitus zygotes.

	25C	10 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	63	16	15	16	15	10	49	51	54	10	10
% initial mortality	25	25	53	50	60	10	6	2	2	0	0
% total mortality	51	38	60	63	67	60	55	34	50	20	10
% abnormality	9	6	0	0	0	0	0	4	2	0	0
% hatching	40	56	40	37	33	40	45	62	48	80	90

	25C	10 C - 48 HR									
	total # zygotes		16	15	15	15	10	45	53	52	10
% initial mortality		31	87	73	80	40	9	4	10	0	10
% total mortality		38	87	80	80	60	67	57	71	20	50
% abnormality		6	0	0	0	0	2	0	0	0	0
% hatching		56	13	20	20	40	31	43	29	80	50

	25C	10 C - 72 HR									
	total # zygotes		16	18	18	15	10	55	51	50	10
% initial mortality		56	77	83	93	40	4	8	10	0	0
% total mortality		80	77	83	93	50	24	33	34	30	0
% abnormality		2	5	0	0	20	16	0	2	0	0
% hatching		18	18	17	7	30	60	67	64	70	100

ORIGINAL PAGE IS  
OF POOR QUALITY

Table 41: Effect of storage at 12C on the embryogenesis of  
F. heteroclitus zygotes.

	25C	12 C - 24 HR									
	CONTROL	FF <sup>1</sup>	GD <sup>2</sup>	2C	4C	64C	10 hr	12 hr	18 hr	24 hr	48 hr
total # zygotes	76	30	21	20	20	25	40	40	37	24	20
% initial mortality	3	13	24	60	30	0	3	5	19	0	10
% total mortality	8	17	24	60	100*	8	8	8	22	4	10
% abnormality	0	3	0	0	0	0	7	5	5	0	5
% hatching	92	80	76	40	0	92	85	88	73	96	85

	25C	12 C - 48 HR									
total # zygotes		26	20	20	17	20	39	41	38	20	28
% initial mortality		19	45	50	47	5	21	7	11	0	32
% total mortality		23	45	50	71	5	23	10	13	0	32
% abnormality		0	0	0	0	15	5	7	3	5	0
% hatching		77	55	50	29	80	72	83	84	95	68

	25C	12 C - 72 HR									
total # zygotes		32	19	19	20	30	30	34	36	31	27
% initial mortality		34	63	47	65	10	20	24	39	3	4
% total mortality		34	63	47	70	10	20	24	39	6	4
% abnormality		3	5	0	0	20	3	3	8	0	4
% hatching		63	32	53	30	70	77	73	53	94	92

\* Dish killed off by fungus infection.

## VI. ASTP VISUAL ORIENTATION TESTS

## A. BIBLIOGRAPHY OF PERTINENT FISH VISION LITERATURE

1. Bogenschütze, H. 1961. Vergleichende Untersuchungen über die optische Komponente der Gleichgewichtshaltung bei Fischen. Z. Vergl. Physiol. 44: 626.
2. Bowman, R.S., and N.S. Sutherland. 1970. Shape discrimination by goldfish: coding of irregularities. J. Comp. Psychol. 72: 90-97.
3. Butcher, E.O. 1938. The structure of the retina of Fundulus heteroclitus and the regions of the retina associated with the different chromatophoric responses. J. Exp. Zool. 79: 275-297.
4. Clark, E. 1961. Visual discrimination in lemon sharks. Abstr. Symp. Papers. 10th Pac. Sci. Congr., Honolulu, Hawaii p. 176-177.
5. Cronly-Dillon, J.R. 1963. Units sensitive to direction of movement in goldfish optic tectum. Exp. Neurol. 7: 46-64.
6. Cronly-Dillon, J.R. 1964. Units sensitive to direction of movement in goldfish optic tectum. Nature 203: 214-215.
7. Crozier, W.J., E. Wolf, and G. Zerrahn-Wolf. 1936-37. On critical frequency and critical illumination for response to flickered light. J. Gen. Physiol. 20: 211-228.
8. Crozier, W.J., E. Wolf, and G. Zerrahn-Wolf. 1936-37. Temperature and critical illumination for reaction to flickering light, II. Sunfish. J. Gen. Physiol. 20: 411-431.

9. Crozier, W.J., and E. Wolf. 1939-40. The flicking response curve for Fundulus. J. Gen. Physiol. 23: 677-694.
10. Davis, R.E. 1968. Environmental control of memory fixation in goldfish. J. Comp. Physiol. Psychol. 65: 72.
11. Gaffron, M. 1934. Untersuchungen über das Bewegungssehen bei Libellenlarven, Fliegen und Fischen. Z. Vergl. Physiol. 20: 299-337.
12. Goodyear, C.P. 1973. Vision and learning in the orientation of mosquito fish, Gambusia affinis. Behavior 45: 191.
13. Greenway, A.P. 1967. Photokinesis in the teleost fish Danio malabaricus. Z. Vergl. Physiol. 56: 416.
14. Hanyu, I., and M.A. Ali. 1963. Flicker fusion frequency of electroretinogram in light-adapted goldfish at various temperatures. Science 140: 662-663.
15. Harden-Jones, F.R. 1963. The reactions of fish to moving backgrounds. J. Exp. Biol. 40: 437-446.
16. Hasler, A.D. 1956. Influence of environmental reference points on learned orientation in fish (Phoxinus). Z. Vergl. Physiol. 38: 303.
17. Hester, F.J. 1968. Visual contrast thresholds of the goldfish (Carassius auratus) Vis. Res. 8: 1315-1336.
18. Hollyfield, J.G. 1972. Histogenesis of the retina in the killifish, Fundulus heteroclitus. J. Comp. Neur. 144: 373-380.

19. Ingle, D. 1968. Spatial dimensions of fish vision. In: The Central Nervous System and Fish Behavior. ed. D. Ingle. University of Chicago Press. Chicago, Ill. 51-59.
20. Inoue, M., and T. Kondo. 1972. On the optomotor reaction of fish relevant to fishing method. I. Reaction of fish to visual patterns. J. Tokyo Univ. Fish. 58: 9-16.
21. Lyall, A.H. 1957. The growth of the trout retina. Quart. J. Microsc. Sci. 98: 101-110.
22. Lyon, E.P. 1904. On rheotropism. Am. J. Physiol. 4: 77-82.
23. Mackintosh, J., and N.S. Sutherland. 1963. Visual discrimination by the goldfish: the orientation of rectangles. Anim. Behav. 11: 135.
24. Main, R. 1927. Rheotropism in Fundulus, a forced movement. Am. J. Physiol. 81: 496.
25. Matthews, W.A. 1964. Shape discrimination in tropical fish. Anim. Behav. 12: 111.
26. McCleary, R.A., and J.J. Bernstein. 1959. A unique method for the control of brightness cues in the study of color vision in fish. Physiol. Zool. 32: 284-292.
27. Meyer, D.L., and H.O. Schwassmann. 1970. Electrophysiological method for determination of refractive state in fish eyes. Vision Res. 10: 1301-1304.
28. Mühlmann, D. 1967. Untersuchungen zum Ablauf der Dunkeladaptation bei Fischen. I. Optomotorische Reaktionen bei Cichlasoma meeki. Z. Vergl. Physiol. 55: 119-133.

29. Muntz, W.R.A., and D.P.M. Northmore. 1973. Scotopic spectral sensitivity in a teleost fish (Scardinius erythrophthalmus) adapted to different day lengths. Vis. Res. 13: 245-252.
30. O, Gower, A.K., and R.F. Mathewson. 1967. Spectral sensitivity and flicker-fusion frequency of the lemon shark, Negaprion brevirostris. In: Sharks, Skates and Rays. eds. D.P. Rall, P.W. Gilbert, R.F. Mathewson. Johns Hopkins, Baltimore, Maryland. Chap. 29, 433-446.
31. Pavlov, D.S., and Y.N. Sbikin. 1967. Study of the spectral and threshold sensitivity of vision in fish by a method of optomotor reaction. Ikhtiologicheskaya Nauk. SSSR Moscow 74-79.
32. Reddy, S.R., and G. Kote. 1975. Predatory behavior of Gambusia affinis in relation to different light colors. Physiol. Behav. 14: 255-258.
33. Regan, D., N.A.M. Schellart, H. SpekReijse, and T.J.T.P. Vandenberg. 1975. Photometry in goldfish by electrophysiological recording: comparison of criterion response method with heterochromatic flicker photometry. Vis. Res. 15: 799-808.
34. Ritter, J.A., and H.R. MacCrimmon. 1973. Influence of environmental experience on response of yearling rainbow trout (Salmo gairdneri) to a black and white substrate. J. Fish. Res. Bd. Can. 30: 1740-1742.

35. Schwassmann, H.W. 1960. Environmental cues in the orientation rhythm of fish. Cold Spr. Harb. Symp. on Quant. Biol. XXV: 443-450.
36. Schwassmann, H.O. 1965. Functional development of visual pathways in larval sardines and anchovies. Mar. Res. Comm. Cal. Coop. Oceanic Fish. Invest. Rept. 10: 64-70.
37. Shaw, E., and A. Tucker. 1965. The optomotor reaction of schooling carangid fishes. Anim. Behav. 13: 330.
38. Shaw, E., and B.D. Sachs. 1967. Development of the optomotor response in the schooling fish, Menidia menidia. J. Comp. Physiol. Psychol. 63: 685.
39. Tamara, T. 1957. A study of visual perception in fish, especially on resolving power and accomodation. Bull. Jap. Soc. Sci. Fisheries 22(9): 536-557.
40. Tester, A.L., and S. Kato. 1966. Visual target discrimination in blacktip sharks (Carcharhinus melanopterus) and grey sharks (C. menissorah). Pacific Sci. 20: 461-471.
41. Varanelli, C.C., and J.D. McCleave. 1974. Locomotor activity of Atlantic salmon parr (Salmo salar L.) in various light conditions and in weak magnetic fields. Anim. Behav. 22: 178-186.
42. Von Schiller, P. 1934. Kinematoskopisches Sehen der Fische. Z. Vergl. Physiol. 20: 454-462.

43. White, E. L. 1948. An experimental study of the relationship between the size of the eye and the size of the optic tectum in the brain of the developing teleost, Fundulus heteroclitus. J. Exp. Zool. 108: 439-469.
44. Wolf, E., and G. Zerrahn-Wolf. 1935-36. Threshold intensity of illumination and flicker frequency for the eye of the sun-fish. J. Gen. Physiol. 19: 495-502.

## B. Protocol of Visual Acclimation Procedures

### Introduction

This experiment was designed to investigate the relative roles of the vestibular and visual systems in orientation and swimming behavior under zero gravity. Hence, it was decided to rear freshly hatched fry under a predetermined visual parameter, i.e. acclimate the fry to a given visual background. Prior to the ASTP flight, the fry would be packaged in plastic bags with one side containing the background to which they were acclimated. The purpose of the experiment was: a) determine if the fry could use a known visual cue to enable them to orient in the absence of gravitational cues, b) to determine if fry reared with a definite visual background would adapt to zero gravity more rapidly than control or blinded fry.

### Methods

1. Large batches of ova were freshly fertilized, incubated at 22C for 21 days and then mechanically disturbed to obtain maximum simultaneous hatching on day 21.
2. The hatchees were divided into 5 equal lots and each lot placed in a separate 10 gal tank, for acclimation to a different visual background.
3. The fry were acclimated to the following visual parameters.
  - a) control group - no special visual background

- b) horizontal bar background on sides of tank - 1/4" black and white bars
  - c) vertical bar background on sides of tank - 1/4" black and white bars
  - d) black overhead background - roof and top half of sides painted black with light from below
  - e) blinded fish.
4. A new group of fry were hatched every 2 weeks, until ultimately a series of 5 tanks for each condition in (3) above were established.
5. By ASTP liftoff time - fry ranging in age from 3-11 weeks, in 2 week intervals, were available.
6. It was decided that the 3 week old fry were the ideal size for packaging and use on the ASTP mission.
7. Fry were placed in large plastic bags with 500 ml sea water with one side of the bag painted with the acclimatization background for that group. The bags were placed in a light tight container and transported to NASA, Houston, for packaging.
8. ASTP E.I.P. stated that the final package had 5 compartments, each with 6 fry, and that the compartments were designated as follows:
- Compartment 1 - Normal controls
  - Compartment 2 - Vertical bar acclimated
  - Compartment 3 - Horizontal bar acclimated
  - Compartment 4 - Black overhead acclimated
  - Compartment 5 - Blinded

### C. Protocol for Film Analysis

1. A Vanguard Analyser was used to facilitate a detailed analysis of the position and locomotor activity of each fish.
2. The duration of time in seconds and the number of frames a given fish was visible for each day's filming sequence were determined, according to the following information provided by NASA:

#### Flight Film - Sequences in Order Seen

##### Roll CI 28

- a) Day 6 - fish filmed at 24 frames per second
- b) Day 7 - fish filmed at 24 frames per second

##### Roll CI 29

- c) Day 2 - fish lost because of improper setting of film magazine
  - d) Day 8 - fish filmed at 12 frames per second
  - e) Day 9 - fish filmed at 12 frames per second
3. The position of each fish was drawn, using tracing paper, at least every 6 frames (i.e. every  $1/4$ - $1/2$  sec). Thus, a frame-by-frame diagram for each fish was established, showing the orientation of that fish relative to its original starting point and its position  $1/4$ - $1/2$  sec previously.
    - a) The numbers within the fish body outline represent a sequential 6 frame series, starting with 0, each number thereafter representing a 6 frame step.
    - b) Occasionally intermediate frame positions are also drawn, for particularly active fish and these are designated by a subscript

e.g. 5<sub>0</sub>, 5<sub>1</sub>, 5<sub>2</sub>, 5<sub>3</sub>, 5<sub>4</sub>, 5<sub>5</sub>, etc., so that each interval could be divided into 6 subintervals if necessary.

- c) Only position changes are noted. If a fish does not move in a given 6 frame interval, the number is not recorded in the body position outline, but is recorded in the margin next to (rather than below) the previous number.

4. The frame-by-frame diagrams were used to construct summary diagrams for each fish on each day. Each summary diagram shows:

- a) the number of frames the fish was visible
- b) the time the fish was visible
- c) the starting or 0 point, i.e. the position of the fish as it first became visible
- d) the pathway and direction of swimming activity of the fish, as indicated by arrows.

#### D. Results of ASTP Film Analysis

1. Film from Day 2 was missing. Unfortunately this was probably the most critical day since this was the earliest possible time that fish could be monitored in what was probably their most active and aggravated state. All subsequent days were to be compared to Day 2 from the standpoint of
  - a) relative quality (looping vs. orientated) and quantity of activity
  - b) comparison of the effectiveness of the various visual parameters in orientation and adaptation to zero gravity.
2. Analysis of Days 6, 7, 8, 9 gave the following results.
  - a) Tables 42-45 represent an analysis of each day representing the conditions in each compartment on the given day.
  - b) Table 46 analyses each compartment and compares certain behavioral parameters over the 4 day test period.
  - c) Tables 47-66 analyzes the activity of each fish in each compartment for each day's filming. It should be noted, however, that the number designations given to each fish, i.e. fish #1, fish #2, etc., are completely arbitrary from day to day, since it is virtually impossible to actually identify individuals.
  - d) Table 67 compares the total amount of looping behavior for each day of the test in all compartments combined. If anything, there was an increase in looping with increasing time, since the smallest amount of looping occurred on Day 6.

Table 42: Summary of Conditions on Day 6.

	COMPARTMENT				
	1	2	3	4	5
Number of fish visible	5	5	5	5	6
Number of fish alive	5	5	5	5	5
Total mortality	1	1	1	1	1
Number of fish looping spontaneously	0	0	1	1	1
Number of fish looping after disturbance	2	0	0	0	2
Total # of fish looping	2	0	1	1	3
Back always to camera	4	4	2	2	5 <sup>3</sup>
Belly to Camera	1 <sup>1</sup>	1	3 <sup>1</sup>	1 <sup>2</sup> +2 <sup>1</sup>	0
Time Filmed (sec.)	13	10-25	33	19	29

<sup>1</sup>intermittently

<sup>2</sup>entirely

<sup>3</sup>difficult to tell because of black ventral coloration of blinded animals.

Table 43: Summary of Conditions on Day 7.

	COMPARTMENT				
	1	2	3	4	5
Number of fish visible	5	5	5	5	6
Number of fish alive	5	5	5	5	4
Total mortality	1	1	1	1	2
Number of fish looping spontaneously	2	1	2	1	4
Number of fish looping after disturbance	0	0	2	1	4
Total # of fish looping	2	1	2	1	4
Back always to camera	5	3	5	1	4
Belly to camera	0	1+1 <sup>1</sup>	0	4 <sup>1</sup>	0
Time filmed (sec.)	9	22	23	24	30

Table 44: Summary of Conditions on Day 8.

	COMPARTMENT				
	1	2	3	4	5
Number of fish visible	5	5	5	3	6
Number of fish alive	5	5	5	2	4
Total mortality	1	1	1	4	2
Number of fish looping spontaneously	0	1	0	0	0
Number of fish looping after disturbance	3	2	0	0	3
Total # of fish looping	3	3	0	0	3
Back always to camera	5	5	3	2	4
Belly to camera	0	0	1+1 <sup>1</sup>	0	0
Time filmed (sec.)	9	17	17	18	19

Table 45: Summary of Conditions on Day 9.

	COMPARTMENT				
	1	2	3	4	5
Number of fish visible	5	5	5	4	6
Number of fish alive	5	5	5	2	4
Total mortality	1	1	1	4	2
Number of fish looping spontaneously	0	0	0	0	0
Number of fish looping after disturbance	3	4	2	0	1
Total # of fish looping	3	4	2	0	1
Back always to camera	5	4	3	2	4
Belly to camera	0	1 <sup>1</sup>	1+1 <sup>1</sup>	0	0
Time filmed (sec.)	10	22	18	21	24

Time Looping

Table 46: Compartment by Compartment Comparison of Activity of  
ASTP Fish.

	DAY			
	6	7	8	9
<b>A. Compartment 1</b>				
Number of fish living	5	5	5	5
Number of fish looping	2	2	3	3
# of fish with belly to camera	1	0	0	0
# of fish with back to camera	4	5	5	5
<b>B. Compartment 2</b>				
Number of fish living	5	5	5	5
Number of fish looping	0	1	3	4
# of fish with belly to camera	1	2	0	1
# of fish with back to camera	4	3	5	4
<b>C. Compartment 3</b>				
Number of fish living	5	5	5	5
Number of fish looping	1	2	0	2
# of fish with belly to camera	3	0	2	1
# of fish with back to camera	2	5	3	4
<b>D. Compartment 4</b>				
Number of fish living	5	5	2	2
Number of fish looping	1	1	0	0
# of fish with belly to camera	3	4	0	0
# of fish with back to camera	2	1	2	2
<b>E. Compartment 5</b>				
Number of fish living	5	4	4	4
Number of fish looping	3	4	3	1
# of fish with belly to camera	0	0	0	0
# of fish with back to camera	5	4	4	4

Table 47: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 1, Day 6.

DAY 6 - COMPARTMENT 1						
FISH	1	2	3	4	5	6
Time visible (sec.)	13	13	13	13	9	
Total time looping (sec.)	0	0	3.5	3	0	
Percent time looping	-	-	27	23	-	
Number of loops	-	-	8	6	-	
Number of loops/sec.	-	-	2.3	2	-	
Looping to left	-	-	yes	no	-	
Looping to right	-	-	yes	yes	-	

Table 48: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 2, Day 6.

DAY 6 - COMPARTMENT 2						
FISH	1	2	3	4	5	6
Time visible (sec.)	25	25	25	25	10	
Total time looping (sec.)	0	0	0	0	0	
Percent time looping	-	-	-	-	-	
Number of loops	-	-	-	-	-	
Number of loops/sec.	-	-	-	-	-	
Looping to left	-	-	-	-	-	
Looping to right	-	-	-	-	-	

Table 49: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 3, Day 6.

DAY 6 - COMPARTMENT 3						
FISH	1	2	3	4	5	6
Time visible (sec.)	41	32	31	33	33	
Total time looping (sec.)	0	0	23.6	0	0	
Percent time looping	-	-	76	-	-	
Number of loops	-	-	8	-	-	
Number of loops/sec.	-	-	0.3	-	-	
Looping to left	-	-	yes	-	-	
Looping to right	-	-	no	-	-	

Table 50 : Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 4, Day 6.

DAY 6 - COMPARTMENT 4						
FISH	1	2	3	4	5	6
Time visible (sec.)	19	19	19	17	19	
Total time looping (sec.)	0	0	4.75	0	0	
Percent time looping	-	-	25	-	-	
Number of loops			2			
Number of loops/sec.	-	-	0.4	-	-	
Looping to left	-	-	no	-	-	
Looping to right	-	-	yes	-	-	

Table 51: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 5, Day 6.

DAY 6 - COMPARTMENT 5						
FISH	1	2	3	4	5	6
Time visible (sec.)	29	19 <sup>1</sup>	29	29	29	29
Total time looping (sec.)	0	0	2.5	0	9.75	7
Percent time looping	-	-	9	-	34	24
Number of loops	-	-	6	-	11	10
Number of loops/sec.	-	-	2.4	-	1.1	1.4
Looping to left	-	-	yes	-	yes	yes
Looping to right	-	-	yes	-	yes	yes

<sup>1</sup> dead

Table 52: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 1, Day 7.

DAY 7 - COMPARTMENT 1						
FISH	1	2	3	4	5	6
Time visible (sec.)	9	9	9	9	3	
Total time looping (sec.)	0	9	0	0	3	
Percent time looping	-	100	-	-	100	
Number of loops	-	8	-	-	5 <sup>1</sup>	
Number of loops/sec.	-	0.9	-	-	1.7	
Looping to left	yes		-	-	no	
Looping to right	no		-	-	yes	

<sup>1</sup> estimate only - fish last in glare part of time

Table 53: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 2, Day 7.

DAY 7 - COMPARTMENT 2						
FISH	1	2	3	4	5	6
Time visible (sec.)	22	22	4	22	10	
Total time looping (sec.)	0	0	4	0	0	
Percent time looping	-	-	100	-	-	
Number of loops	-	-	6	-	-	
Number of loops/sec.	-	-	1.5	-	-	
Looping to left	-	-	yes	-	-	
Looping to right	-	-	no	-	-	

Table 54 : Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 3, Day 7.

DAY 7 - COMPARTMENT 3						
FISH	1	2	3	4	5	6
Time visible (sec.)	23	24	23	23	23	
Total time looping (sec.)	0	4.8	23	0	0	
Percent time looping	-	20	100	-	-	
Number of loops	-	5	30	-	-	
Number of loops/sec.	-	1.0	1.3			
Looping to left	-	yes	yes	-	-	
Looping to right	-	yes	yes	-	-	

Table 55: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 4, Day 7.

DAY 7 - COMPARTMENT 4						
FISH	1	2	3	4	5	6
Time visible (sec.)	24	24	24	15	24	
Total time looping (sec.)	0	0	0	0	8.25	
Percent time looping	-	-	-	-	34	
Number of loops	-	-	-	-	6	
Number of loops/sec.	-	-	-	-	0.7	
Looping to left	-	-	-	-	no	
Looping to right	-	-	-	-	yes	

Table 56: Fish-by-Fish Analysis of Behavior of Fish in  
 Compartment 5, Day 7.

DAY 7 - COMPARTMENT 5						
FISH	1	2	3	4	5	6
Time visible (sec.)	30	30	30	30	30	13
Total time looping (sec.)	0 <sup>1</sup>	19.75	25.5	8.75	22.25	0 <sup>1</sup>
Percent time looping	-	66	85	29	74	-
Number of loops	-	27	24	19	14	-
Number of loops/sec.	-	1.4	0.9	2.1	0.6	
Looping to left	-	yes	yes	yes	yes	-
Looping to right	-	yes	yes	yes	yes	-

<sup>1</sup>fish is dead

Table 57: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 1, Day 8.

DAY 8 - COMPARTMENT 1						
FISH	1	2	3	4	5	6
Time visible (sec.)	9	9	9	9	9	
Total time looping (sec.)	7.5	0	45	7.5	0	
Percent time looping	83	-	50	83	-	
Number of loops	6	-	3	4	-	
Number of loops/sec.	0.8	-	0.7	0.5	-	
Looping to left	yes	-	yes	no	-	
Looping to right	yes	-	yes	yes	-	

Table 58: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 2, Day 8.

DAY 8 - COMPARTMENT 2						
FISH	1	2	3	4	5	6
Time visible (sec.)	17	11	17	17	17	
Total time looping (sec.)	14	11	0	0	17	
Percent time looping	82	100	-	-	100	
Number of loops	15	7	-	-	12 <sup>2</sup>	
Number of loops/sec.	1.1	0.6	-	-	0.7	
Looping to left	yes	yes	-	-	yes	
Looping to right	yes	yes	-	-	yes	

<sup>2</sup> approximate - lost in glare part of time

Table 59: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 3, Day 8.

DAY 8 - COMPARTMENT 3						
FISH	1	2	3	4	5	6
Time visible (sec.)	17	17	17	17	5	
Total time looping (sec.)	0	0	0	0	0	
Percent time looping	-	-	-	-	-	
Number of loops	-	-	-	-	-	
Number of loops/sec.	-	-	-	-	-	
Looping to left	-	-	-	-	-	
Looping to right						

Table 60: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 4 , Day 8.

DAY 8 - COMPARTMENT 4						
FISH	1	2	3	4	5	6
Time visible (sec.)	17	7.5	17			
Total time looping (sec.)	0	0	0 <sup>1</sup>			
Percent time looping	-	-	-			
Number of loops	-	-	-			
Number of loops/sec.	-	-	-			
Looping to left	-	-	-			
Looping to right	-	-	-			

<sup>1</sup>dead

Table 61: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 5, Day 8.

DAY 8 - COMPARTMENT 5						
FISH	1	2	3	4	5	6
Time visible (sec.)	19.5	18.5	19	20	19	19
Total time looping (sec.)	16.5	0 <sup>1</sup>	0	15	0 <sup>1</sup>	10
Percent time looping	85	-	-	75	-	50
Number of loops	17	-	-	22	-	15
Number of loops/sec.	1.0	-	-	1.5	-	1.5
Looping to left	yes	-	-	yes	-	yes
Looping to right	yes	-	-	yes	-	yes

<sup>1</sup> possibly dead

Table 62: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 1, Day 9.

DAY 9 - COMPARTMENT 1						
FISH	1	2	3	4	5	6
Time visible (sec.)	12	10.5	10.5	10.5	10.5	
Total time looping (sec.)	0	5	8.5	0	8	
Percent time looping	-	48	81	-	76	
Number of loops	-	7	10	-	9	
Number of loops/sec.	-	1.4	1.2	-	1.1	
Looping to left	-	no	yes	-	no	
Looping to right	-	yes	yes	-	yes	

Table 63: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 2, Day 9.

DAY 9 - COMPARTMENT 2						
FISH	1	2	3	4	5	6
Time visible (sec.)	22	22	22	22	22	
Total time looping (sec.)	8	17.5	10.5	12	0	
Percent time looping	36	80	48	55	-	
Number of loops	7	13	9	9	-	
Number of loops/sec.	0.9	0.8	0.9	0.8	-	
Looping to left	yes	yes	yes	yes	-	
Looping to right	yes	yes	yes	yes	-	

Table 64 : Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 3, Day 9.

DAY 9 - COMPARTMENT 3						
FISH	1	2	3	4	5	6
Time visible (sec.)	18	18	18	18	18	
Total time looping (sec.)	7	7	0	0	0	
Percent time looping	39	39	-	-	-	
Number of loops	8	14	-	-	-	
Number of loops/sec.	1.1	2	-	-	-	
Looping to left	yes	yes	-	-	-	
Looping to right	yes	yes	-	-	-	

Table 65: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 4, Day 9.

DAY 9 - COMPARTMENT 4						
FISH	1	2	3	4	5	6
Time visible (sec.)	21	21	21	20		
Total time looping (sec.)	0	0 <sup>1</sup>	0 <sup>1</sup>	0		
Percent time looping	-	-	-	-		
Number of loops	-	-	-	-		
Number of loops/sec.	-	-	-	-		
Looping to left	-	-	-	-		
Looping to right	-	-	-	-		

<sup>1</sup> dead

Table 66: Fish-by-Fish Analysis of Behavior of Fish in  
Compartment 5, Day 9.

DAY 9 - COMPARTMENT 5						
FISH	1	2	3	4	5	6
Time visible (sec.)	24.5	24.5	24.5	12.5	12.5	24.5
Total time looping (sec.)	0	0 <sup>1</sup>	0 <sup>1</sup>	3	0	0 <sup>1</sup>
Percent time looping	-	-	-	24	-	-
Number of loops	-	-	-	6	-	-
Number of loops/sec.	-	-	-	2	-	-
Looping to left	-	-	-	yes	-	-
Looping to right	-	-	-	no	-	-

<sup>1</sup> dead

Table 67: Comparison of Total Looping Behavior on Day-by-Day Basis.

	D 6	D 7	D 8	D 9
Number of Fish Alive	25	24	21	21
Number of Fish Looping	7	10	9	10
Percent of Fish Looping	28	42	43	48

Table 68: Comparison of Total Looping Behavior on Compartment-by-Compartment Basis.

	1	2	3	4	5
Total # of Fish-Days <sup>1</sup>	20	20	20	14	17
Avg. # Fish/Cmpt./Day	5	5	5	3.5	4.25
Total # of Fish-Day Looping <sup>2</sup>	10	8	5	2	11
Avg. # Fish-Looping/Cmpt.	2.5	2	1.25	0.5	2.75
Percent of Fish Looping/Cmpt.	50	40	25	14	65

<sup>1</sup>Total of live fish present in that compartment over 4 day period.

<sup>2</sup>" " - " looping " " " " " " "

The increase in looping behavior thereafter, however, may be due to an increased tendency on the part of the astronauts to "force" the fish to loop by tapping or shaking the bags. Thus, I feel the increased looping is not indicative, since undistrubed fish may have oriented normally.

- e) Table 68 compares the total looping behavior on a compartment-by-compartment basis over the total test period. The greatest amounts of looping occurred in Cmpt. 1 and 5, i.e. the controls and blinded fish. The least amount of looping occurred in Cmpt. 4, i.e. black background fishes. The ability of these fish to orient is also supported by the fact that most of them oriented with their backs to the black background, Table 46 D., Days 6 and 7, unlike most of the fishes in other compartments.

Whether the percent looping behavior of the fishes in Cmpts. 2 and 3 is significantly less than that of the controls could probably be proven statistically. Thus, it would appear that prior visual acclimation did permit those fishes to orient somewhat better than fishes with no visual acclimation.

- f) Tables 69 and 70 present both a day-by-day and compartment-by-compartment analysis of the total time duration of looping and the number of loops made. I can see no discernable patterns here, other than the smaller amounts of time and lesser number of loops made by the fishes in

Cmpt. 4 (Table 70).

Table 69: Day-by-Day Comparison of Duration and Quality of Looping.

	D 6	D 7	D 8	D 9
Total # fish looping	7	10	9	10
Avg. percent time looping	31	71	79	53
Avg. number of loops/sec.	1.4	1.2	0.9	1.2

Table 70: Compartment-by-Compartment Comparison of Duration and Quality of looping.

	Compartment				
	1	2	3	4	5
Average # fish looping/ cmpt./day	2.5	2	1.25	0.5	2.75
Average percent time looping/fish	67	63	55	30	51
Average # loops/sec./fish	1.3	0.9	1.1	0.6	1.5

Conclusions.

The success of the two main objectives of the experiment can be summarized as follows:

- a) No conclusions can be drawn as to whether prior visual acclimation would allow those fishes to become visually oriented more rapidly than non-acclimated fishes. The quantity of disorientation was observed to increase rather than decrease over the test period. But, as pointed out, this may be due to the fact that the astronauts increased their efforts to "force" the fishes to become disoriented by tapping or shaking the bags.
- b) Prior visual acclimation, however, did seem to enable such fishes to orient better than non-acclimated fish, since the least amounts of looping was shown by fishes in Cmpts. 2, 3, and especially 4.

## VII. ELECTRON MICROSCOPY

The relationships between the various ovum envelopes and the mechanism of fertilization was studied by scanning electron microscopy (SEM).

### Methods.

Only mature ova were used for this investigation and were obtained by stripping gravid females. The ova were fixed in 2% glutaraldehyde (in 25<sup>0</sup>/<sub>00</sub> Instant Ocean solution) for 2 hours at room temperature and then rinsed with distilled water; a graded ethanoldehydration sequence prior to 3 changes in amyl acetate and subsequent critical point drying in CO<sub>2</sub> was used to bring the specimens to dryness. The ova were mounted on aluminum stubs, coated with gold/palladium in an ion sputtering apparatus and examined on a JOEL JSM U-3 Scanning Electron Microscope.

### Results.

1. The outer surface of the ovum is covered with a dense pile of short (0.3-0.6 $\mu$  dia) fibrils (Fig. 1a). Several thicker (0.8-1.0 $\mu$  dia.), longer fibrils are present in the area peripheral to the micropyle. These fibrils are part of the outermost, jelly coat layer of the ovum.
2. The chorion lies directly beneath the jelly coat. The outermost portion of the chorion is a thin, homogeneous zone about 0.3-0.4 $\mu$  thick (Figs. 2a, 3c). Immediately below this homogeneous

zone is a heterogeneous portion of the chorion comprised of several lamellae, numbering from a few as 4 in some ova (Fig. 2c) to up to 9-10 in others. Each lamellus appears to be continuous throughout the chorion. Thinner lamellae are found at the inner and outer extremes of the chorion, especially when there are a larger number (Fig. 2a). The entire lamellar zone is about 9-12 $\mu$  thick.

3. Internal to the lamellar zone is a homogeneous, chorionic crystalline zone. This zone was not always detected (Figs. 2c, 3a, c). When present, it varies from a maximum of 13 $\mu$  in thickness (Fig. 2a, d), to a relatively thin 1 $\mu$  (Fig. 3b).

4. The micropyle apparatus consists of a relatively smooth-sided, funnel-shaped vestibule (Fig. 1c) at the bottom of which is a 4-5 $\mu$  dia. opening, the micropyle, surrounded by a slightly elevated lip (Fig. 1b). The micropyle canal traverses the chorion, diminishing in diameter as it approaches the ovoplasm. The inner micropyle aperture rests directly against the ovoplasm (Fig. 1d). The canal sides are irregular, showing evidence of the lamellated structure of the chorion (Figs. 1b, d).

#### Conclusions.

1. The presence of a thick, impenetrable chorion surrounding the plasma membrane precludes any access of sperm to the ovum nucleus other than by means of the micropyle apparatus.




2. The size of the micropyle apertures, outer 4-5 $\mu$  dia., inner 2-3 $\mu$  dia., are such that although several sperm may approach the outer

aperture and possibly enter, the canal becomes progressively narrower and it is conceivable that only one sperm makes contact with the plasma membrane.

3. The absence of fibrils in the region of the micropyle is an adaptation facilitating fertilization. These fibrils are sticky and would collect sperm and prevent their access to the micropyle.




## FIGURE 1.

- a. Surface view of ovum showing fibrils and fibril free vestibule (v) surrounding micropyle (m) X500.
- b. External micropyle aperture with lip (l) and inner view of micropyle canal, X4000.
- c. Side view of vestibule with micropyle at bottom, X500.
- d. Side view of micropyle canal (mc) with portion of cell cytoplasm (c) adjacent to inner micropyle aperture, X2000.



## FIGURE 2.

- a. Transverse view of chorion homogeneous zone (hz), lamellar zone (lz), and tongue of chorion crystalline zone (cz), X1000.
- b. Transverse view of chorion lamellar zone with 7 lamellae, X2000.
- c. Transverse view of chorion lamellar zone with 4 lamellae showing herringbone pattern, X1000.
- d. Transverse view of lamellar and crystalline zones of chorion, X1000.



## FIGURE 3.

- a. Transverse view of lamellar zone of chorion with 10 lamellae and with underlying cytoplasm (c), X2000.
- b. Transverse view of chorion showing extremely thin crystalline zone (cz), X2000.
- c. Chorion homogeneous zone (hz) and lamellar zone (lz) with fibrillar protein matrix, X500.
- d. Chorion lamellar zone (lz) showing fibrillar protein matrix (f = jelly coat fibrils), X1500.

