

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

RECEIVED BY
NASA STI FACILITY
DATE: 9/15/76
DCAF NO. 090211
PROCESSED BY
 NASA STI FACILITY
 ESA - SDS AIAA

Title of Investigation: 28990

Investigation of Environmental Change Pattern in JAPAN.

7.6-10.465
CR-148697

Principal Investigator: Takakazu MARUYASU *eto*
Science University of Tokyo
Noda City, Chiba ken, 278, JAPAN

August 16, 1976

Quarterly Progress Report for period

April-June 1976

Name and Address of National Sponcering Agency	Science and Technology Agency Kasumigaseki 2-2-1 Chiyodaku Tokyo, 100, JAPAN
--	--

**ORIGINAL CONTAINS
COLOR ILLUSTRATIONS**

Original photography may be purchased from:
EROS Data Center
10th and Dakota Avenue
Sioux Falls, SD 57198.

(E76-10465) INVESTIGATION OF ENVIRONMENTAL
CHANGE PATTERN IN JAPAN. 1: INVESTIGATION
OF SOIL EROSION IN HOKKAIDO WHICH IS CAUSED
BY THAWING OF SOIL WATER IN LATE SPRING
Quarterly Progress Report. (Science Univ. G3/43. 00465

N76-31613
HC \$3.50

Unclas

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo

Co-Investigator: Hiroaki OCHIAI
Toba Merchant Marine College

Yasuhiro SUGIMORI
Chief of Disaster Analysis Laboratory,
National Research Center for Disaster Protection
Science and Technology Agency

Daitaro SHOJI
Chief of Hydrographer, Hydrographic Department
Marine Safety Agency

Kaname TAKEDA
Leader of Research Group, National Institute
of Resources.

Kiyoshi TSUCHIYA
National Space Development Agency of JAPAN

Iwao NAKAJIMA
Section Chief of Management Technology, Forest
Experiment Station, Forest Agency

Takamasa NAKANO
Faculty of Science, Tokyo Metropolitan University

Shigechika HAYASHI
Chief of Soil Improvement Laboratory, Hokkaido
National Agricultural Experiment Station
Upland Farming Division, Forest Agency

Seiji HORIKAWA
Faculty of Engineering, University of Tokyo

Shunji MURAI
Institute of Industrial Science, University of
Tokyo

Kantaro WATANABE
Faculty of Marine Science and Technology,
Tokai University

Taisuke WATANABE
Chief of No 2 Resources Section, Tokai Region
Fisheries Research Laboratory, Fishery Agency

Title of Investigation: 28990 Investigation of Environmental
Change Patterns in JAPAN

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo
Noda City, Chiba ken, 278 , JAPAN

Quarterly Progress Report for Period
April 1976-June 1976

Name and address of National Sponcering Agency
Science and Technology Agency
Kasumigaseki 2-2-1 Chiyoda ku,
Tokyo, 100, JAPAN

Subject

1. Investigation of Soil Erosion in Hokkaido which is caused by
Thawing of Soil Water in Late Spring

Co-Investigator: Shigechika HAYASHI
Upland Farming Division,
National Agricultural Experiment
Station,
Memuro, Hokkaido, 082, JAPAN

1. Introduction.

This paper reports the results of the study about the analysis of the changes occurred in the fields of Tokachi district, where we regarded as the test site of this study, using photographic treatment and the density slicing instruments.

This study was performed by following members:

Michikazu Fukuhara of Upland Farming Division, Hokkaido National Agricultural Experiment Station, Yoshizumi Yasuda and Yasubumi Emori of Institute of Color Technology, Faculty of Engineering, Chiba University.

2. Techniques.

B & W bulk positives of 4, 5, 7 bands in the two scenes of LANDSAT data, the one obtained on 11 June 1975, the other, 24 May 1973, were prepared.

After they were enlarged to 1:200,000 scale, the ratio images of 1975/1973 of each band were made.

These images were analysed using slicing instrument (GRAF-COLOR 905), and the color-coded density maps obtained were interpreted.

3. Accomplishment.

1975/1973 ratio images are shown in photo-1 (band 4), photo-2 (band 5) and photo-3 (band 7). In these prints, the red colored portion indicates the increased reflectance, blue-colored, the decreased.

Pattern, expressed in the prints of the color-coded maps are considered to be classified as follows:

1). The place where land management changed. For instance, the places where were allowed fallow in 1973 and became planted in 1975 (photo-3 ^{red} blue parts).

2). Seasonal variation of vegetation. In natural or grazed grasslands, the ground surface is covered with dead grass in May, but in June, it is covered with vigorously grown grass (photo-1 blue, photo-2 blue).

3). Apparent change caused by difference of relative reflectance among objects in images. River beds or street area where no vegetation is seen (photo-1 red). Alluvial soils or volcanic ash soils mixed with subsoils (photo-2 red).

4). The part difficult to interpret (photo-3 blue).

4. Conclusion.

Multidate analysis of LANDSAT data was tested using ratio image method. As by this method, many informations have been obtained which cannot be obtained by means of single scene analysis, this method is expected to have ^{many} ~~great~~ possibility to interpret the imageries of agricultural fields which show large annual fluctuation.

ORIGINAL PAGE IS
OF POOR QUALITY

5




Photo.-1
(BAND 4)




Photo.-2
(BAND 5)



Photo.-3
(BAND 7)

1975/1973 ratio imges.



288017

"Made available Under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information, and without liability
for any use made thereof."

RECEIVED BY
NASA STI FACILITY
DATE: 9/15/76
DCAF NO. [] [] [] [] [] []
PROCESSED BY
 NASA STI FACILITY
 ESA-SDS AIAA

Title of Investigation: 28990

Investigation of Environmental Change Pattern in JAPAN.

E 7.6 - 10.465
NASA-CR-148697

Principal Investigator: Takakazu MARUYASU *etc*
Science University of Tokyo
Noda City, Chiba ken, 278, JAPAN

August 16, 1976

Quarterly Progress Report for period

April-June 1976

Name and Address of
National Sponcering
Agency
Science and Technology Agency
Kasumigaseki 2-2-1 Chiyodaku
Tokyo, 100, JAPAN

Original photography may be purchased from:
EROS Data Center
10th and Dakota Avenue
Sioux Falls, SD 57198

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo

Co-Investigator: Hiroaki OCHIAI
Toba Merchant Marine College

Yasuhiro SUGIMORI
Chief of Disaster Analysis Laboratory,
National Research Center for Disaster Protection
Science and Technology Agency

Daitaro SHOJI
Chief of Hydrographer, Hydrographic Department
Marine Safety Agency

Kaname TAKEDA
Leader of Research Group, National Institute
of Resources.

Kiyoshi TSUCHIYA
National Space Development Agency of JAPAN

Iwao NAKAJIMA
Section Chief of Management Technology, Forest
Experiment Station, Forest Agency

Takamasa NAKANO
Faculty of Science, Tokyo Metropolitan University

Shigechika HAYASHI
Chief of Soil Improvement Laboratory, Hokkaido
National Agricultural Experiment Station
Upland Farming Division, Forest Agency

Seiji HORIKAWA
Faculty of Engineering, University of Tokyo

Shunji MURAI
Institute of Industrial Science, University of
Tokyo

Kantaro WATANABE
Faculty of Marine Science and Technology,
Tokai University

Taisuke WATANABE
Chief of No 2 Resources Section, Tokai Region
Fisheries Research Laboratory, Fishery Agency

Title of Investigation: 28990 Investigation of Environmental
Change Patterns in JAPAN

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo
Noda City, Chiba ken, 278 , JAPAN

Quarterly Progress Report for Period
April 1976-June 1976

Name and address of . Science and Technology Agency
National Sponcering Kasumigaseki 2-2-1 Chiyoda ku,
Agency Tokyo, 100, JAPAN

Subject

1. Investigation of Soil Erosion in Hokkaido which is caused by
Thawing of Soil Water in Late Spring

Co-Investigator: Shigechika HAYASHI
Upland Farming Division,
National Agricultural Experiment
Station,
Memuro, Hokkaido, 082, JAPAN

1. Introduction.

This paper reports the results of the study about the analysis of the changes occurred in the fields of Tokachi district, where we regarded as the test site of this study, using photographic treatment and the density slicing instruments.

This study was performed by following members:

Michikazu Fukuhara of Upland Farming Division, Hokkaido National Agricultural Experiment Station, Yoshizumi Yasuda and Yasubumi Emori of Institute of Color Technology, Faculty of Engineering, Chiba University.

2. Techniques.

B & W bulk positives of 4, 5, 7 bands in the two scenes of LANDSAT data, the one obtained on 11 June 1975, the other, 24 May 1973, were prepared.

After they were enlarged to 1:200,000 scale, the ratio images of 1975/1973 of each band were made.

These images were analysed using slicing instrument (GRAF-COLOR 905), and the color-coded density maps obtained were interpreted.

3. Accomplishment.

1975/1973 ratio images are shown in photo-1 (band 4), photo-2 (band 5) and photo-3 (band 7). In these prints, the red colored portion indicates the increased reflectance, blue-colored, the decreased.

Pattern, expressed in the prints of the color-coded maps are considered to be classified as follows:

1). The place where land management changed. For instance, the places where were allowed fallow in 1973 and became planted in 1975 (photo-3 ^{red} ~~blue~~ parts).

2). Seasonal variation of vegetation. In natural or grazed grasslands, the ground surface is covered with dead grass in May, but in June, it is covered with vigorously grown grass (photo-1 blue, photo-2 blue).

3). Apparent change caused by difference of relative reflectance among objects in images. River beds or street area where no vegetation is seen (photo-1 red). Alluvial soils or volcanic ash soils mixed with subsoils (photo-2 red).

4). The part difficult to interpret (photo-3 blue).

4. Conclusion.

Multidate analysis of LANDSAT data was tested using ratio image method. As by this method, many informations have been obtained which cannot be obtained by means of single scene analysis, this method is expected to have ^{many} ~~great~~ possibility to interpret the imageries of agricultural fields which show large annual fluctuation.




Photo.-1
(BAND 4)




Photo.-2
(BAND 5)




Photo.-3
(BAND 7)

1975/1973 ratio imges.

1
@abs The author has identified the following significant results.

Using MSS channels 5 and 7 and a supervised classification system with a PPD classification algorithm, it was possible to map the exact areal extent of the snow cover and of the transition zone with melting ~~the~~ snow

patches and snow free parts of various sizes over a large area under different aspects such as relief, exposure, ^{and} shadows, etc. A correlation of the data from ground control, areal underflights and earth-resources satellites provided a very accurate interpretation of the melting procedure of snow in high mountains.