

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

03

II

STIF

~~available under NASA sponsorship  
in the interest of early and wide dis-  
semination of Earth Resources Survey  
Program information and without liability  
for any use made thereof.~~

Department of Applied Earth Sciences  
Stanford University  
Stanford, California 94305

April 1, 1977

7.7-10147

CR-152640

Mr. J. C. Broderick  
Goddard Space Flight Center  
Greenbelt, Maryland, 20771

"Made available under NASA sponsorship  
in the interest of early and wide dis-  
semination of Earth Resources Survey  
Program information and without liability  
for any use made thereof."

Dear Mr. Broderick

Attached is a progress report for the investigation number  
28150, on behalf of Mr. Muftah M. Unis.

Please contact me if you have any questions. Your comments and  
suggestions are greatly appreciated and I look forward to hearing from  
you.

My best regards,

Sincerely,



A. A. Missallati

28150

(E77-10147) SIMULTANEOUS USE OF LANDSAT  
DIGITAL DATA WITH THAT OF GEOPHYSICAL AND  
GEOLOGICAL SURVEYS IN GEOMATHEMATICAL  
MODELLING FOR MINERAL EXPLORATION PREDICTION  
Progress Report, 2 Nov. 1976 - 2 Apr. 1977

E77-21513  
HC A02/MFA01  
Unclas  
G3/43 00147

RECEIVED

CC: Mr. Muftah M. Unis  
CC: Mrs. Lisa Robok

APR 07 1977

JS/902.6

**LANDSAT TYPE 1 REPORT (April 2, 1977)**

---

**A-TITLE OF THE INVESTIGATION:** Simultaneous use of Landsat digital data  
----- with that of geophysical and geological  
surveys in geomathematical modelling for  
mineral exploration prediction.

**B-INVESTIGATION #:** 28150  
-----

**C-PRINCIPAL INVESTIGATORS:**  
-----

**A. A. Missallati**  
-----

Department of Applied Earth Sciences  
Stanford University  
Stanford, California, 94305

Phone (415) 447-2747

**R. J. P. Lyon**  
-----

Department of Applied Earth Sciences  
Stanford University  
Stanford, California, 94305

Phone (415) 447-2747

**D-TECHNICAL MONITOR:**  
-----

**J. C. Broderick**  
-----

Goddard Space Flight Center  
Greenbelt Maryland, 20771

Phone (301) 982-4826

**E-PERIOD:**  
-----

November 2, 1976 to April 2, 1977

**F-ACTION REQUIRED:**  
-----

None

---

Remote Sensing Laboratory  
Stanford University  
Stanford, California, 94305


G-ABSTRACT:

A geological and mineral exploration survey program with particular emphasis on uranium mineralization in three areas in southern Libya (Fig. 1) was initiated by The Industrial Research Center (Department of Geological Researches and Mining) in Libya in 1974. The geological and geophysical studies in these three areas indicated the favorability of finding radioactive minerals and several anomalous zones were defined.

Area 2 (Fig. 1) has been selected as a test site for the present investigation which involves developing methods (through the application of geomathematical models) to combine and use information on regional geology, geophysics, Landsat digital data and geomathematics in computer-assisted, regional uranium resource evaluation. This investigation is now being conducted at the Stanford Remote Sensing Laboratory, School of Applied Earth Sciences.

In this investigation 43 variables, including age and type of rocks, contact relationships, structural relationships, aero-radiometric and aero-magnetic anomalies and the four Landsat-2 quantified channels (and their 6 ratios) are being used as input for the geomathematical models. One of the objectives of the study is to reveal the interrelationship among the variables and its relation to uranium occurrences. This will define which variable or variables (geological, geophysical or Landsat-2 digital data) are more significant indicators of uranium occurrences in this locality, and consequently, which may be disregarded.

# LOCATION MAP



The anticipated results of this project are:

1-To locate specific target areas for uranium mineralization

2-To establish stronger criteria for the detection of radioactive mineral deposits, under differing conditions, by defining the variable or variables which are statistically more indicative of uranium occurrences. Such variables may include the selection of most-suitable MSS band imagery combinations, for the detection of radioactive minerals in these type of rocks.

3-To justify, and/or improve the derived criteria for the use of such techniques through applications in similar areas nearby and elsewhere.

H-PROBLEMS:

The delay in receiving the computer compatible tapes from EROS Data Center was the only and the main problem faced during the progress of this project.

Tapes were ordered on November 11, 1976 and were received on February 17, 1977, a delay of more than 3 months.

# I-ACCOMPLISHMENTS:

1-SELECTION OF VARIABLES: 43 variables were selected, these

include the following parameters

## A-GEOLOGICAL: Include,

age and type of rocks	8 variables
contact relationships	22 variables
structural relationships	1 variable

## B-GEOPHYSICAL: Include,

aero-radiometric	1 variable
aero-magnetic	1 variable

## C-LANDSAT-2 DIGITAL DATA: Include,

(4) Channels	4 variables
(6) Ratios	6 variables

Total 43

## 2-SELECTION OF PILOT AREA:

An area 26 by 28 Km (728 sq. Km)

containing all variables was selected as a pilot area.

## 3-CONSTRUCTION OF A GRID SYSTEM:

A grid system oriented along the Landsat azimuth was constructed. The grid lines were printed on the geophysical and geological maps ( 1:50,000 scale ) at intervals of 500M by 450M cells (1cm by 0.9cm) representing 6 by 8 pixels to record the geological and geophysical data. The 500M by 450M cells were each subdivided further into 90 (1mm by 1mm) subcells to quantify the geological data according to percentage covered in each cell.

**J-SIGNIFICANT RESULTS:**

-----  
A computer test file was created for the analysis of two rows in the pilot area. The data were organized into two groups "with" and "without" radioactivity above 1200 counts. The first 42 samples correspond to the geographic cells which are anomalous, and the rest of the samples (70) are without anomalous values.

Analysis of the output shows that all samples were properly classified.

**K-PLANNING FOR NEXT PERIOD:**

-----  
Assembling all data (geological, geophysical and Landsat-2 digital data) in a computer master file to make the final analysis for the pilot area.

**L-PUBLISHED MATERIALS:**

-----  
"Satellite Remote Sensing Seeks Radioactive Mineral Deposits"

The Stanford Earth Scientist  
Vol. 5 No. 3 Winter, 1977 (enclosed)

**M-RECOMMENDATIONS:**

-----  
None

**N-CHANGING IN STANDING ORDER FORMS:**

-----  
None

**O-DATA REQUEST FORMS SUBMITTED:**

-----  
None

P-LANDSAT-2 IMAGES ACQUIRED OVER TEST AREA:

<u>Type of coverage</u>	<u>Film source</u>	<u>Scene ID</u>	<u>CCT</u>
LandSat-2 (MSS) Scene	B&W FCC	2492-08250	Y
LandSat-2 (MSS) Scene	B&W FCC	2492-08252	Y
LandSat-2 (MSS) Scene	B&W FCC	2564-08230	Y
LandSat-2 (MSS) Scene	B&W FCC	2564-08232	Y

S-MAILING LIST

J. C. Broderick  
Goddard Space Flight Center  
Greenbelt Maryland, 20771

Mr. Muftah M. Unis  
Directorate of Surveys  
Ministry of Planning and Scientific Affairs  
P. O. Box 600  
Tripoli, Libya

Lisa Robock  
International Planning and Programs  
Office of International Affairs  
NASA Washington D.C. 20546

Chairman of the Geology Department  
University of Tripoli  
Tripoli, Libya



# THE STANFORD EARTH SCIENTIST

Volume 5, Number 3

Winter, 1977

## Satellite Remote Sensing Seeks Radioactive Mineral Deposits

Previous geological, geophysical, and geochemical studies indicated the favorability of finding radioactive mineral deposits in the Tibesti Region of southern Libya. Satellite remote sensing studies, to compare the results of these previous airborne and ground-based surveys, with the LANDSAT 2 digital data are now being carried out.

The main objective is to develop and apply methods of combining geological, geophysical, and geochemical information together with LANDSAT (as digital data matrices) to assist in locating areas of potential uranium mineralization. In combining all these information sources a geomathematical model is being used, as outlined above and developed by Dr. Alfredo Prelat, now at the Remote Sensing Laboratory, School of Earth Sciences.

In this project 43 variables, including age and type of rocks, contact relationships, structural relationships, spectrometric and magnetic anomalies, and LANDSAT-2 four quantified channels (and their ten ratios) are being used as input for the geomathematical model. One of the objectives of the study is to reveal the interrelationship among the variables and its relation to uranium occurrence. This will define which variables (geological, geophysical, or LANDSAT-2 digital data) are more significant indicators of uranium occurrence in this locality.

The anticipated results of this project are:

1. To locate target areas for uranium mineralization.
2. To establish stronger criteria for the detection of radioactive mineral deposits, under differing conditions by defining the variables which are statistically more indicative of the uranium occurrences. Such variables may include the selection of the most suitable MSS band imagery combinations, for the detection of radioactive minerals in this type of rocks.
3. To justify, and/or improve the derived criteria for the use of such techniques through applications in similar areas, nearby and elsewhere.

This project is being sponsored by a grant to Dr. A. Missillati from the University of Tripoli in Libya.

### Brief Biographies of Drs. A. Prelat and A. Missillati


**Dr. Alfredo Prelat:** B.Sc. Geology 1967, Universidad de la Plata, Argentina; M.S. Geology 1971, Stanford University, Stanford, Calif. Thesis: Application of Markov Chains to Oil Exploration; Ph.D. Geology 1974, Stanford University, Stanford, Calif. Thesis: Statistical Estimation of Oil Discovery Probabilities; Post-Doctoral 1974-76, Royal Norwegian Scientific and Industrial Research Fellowship, taken at the Norwegian Geological Survey (NGU), Trondheim, Norway; Post-Doctoral 1977, Remote Sensing Laboratory, Stanford University, Stanford, Calif.

Professional: Field Geologist 1966, United Nations, in the

Andean Ranges, Argentina; Research Assistant, Computer Applications in Geology, Kansas Geological Survey, Summers 1971, 1972, 1973; Assistant Geologist, Oil Exploration, Argentina, 1968; and Research Associate 1974, Remote Sensing Laboratory, Stanford (1/74 to 7/74).

**Dr. Amin Missillati:** B.Sc. Geology 1965, University of Libya; M.S. Geology 1967, Columbia University, N.Y.C., N.Y.; Ph.D. Geology 1972, Stanford University, Stanford, Calif.; Post-Doctoral 1977, Remote Sensing Laboratory, Stanford University, Stanford, Calif.

Professional: Assistant Professor, 1972 to date, University of Tripoli, Libya; 1973-74, Chairman, Geology Department, University of Tripoli; 1974-76 Vice-Dean, Faculty of Science, University of Tripoli; Consultant in Industrial Scientific Research at the Industrial Research Center, the General Water Authority, the Atomic Energy Commission, and other private firms in Libya.



Mineral occurrence probability map for the Meråker mapsheet.