

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.



1. Report No. NASA TM-75539	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle INFRARED RADIATION OF VENUSIAN CLOUDS		5. Report Date October 1978	6. Performing Organization Code
		8. Performing Organization Report No.	
7. Author(s) L. V. Ksanfomaliti, et. al.		10. Work Unit No.	
		11. Contract or Grant No. NASW-3198	
9. Performing Organization Name and Address SCITRAN Box 5456 Santa Barbara, CA 93108		13. Type of Report and Period Covered Translation	
		14. Sponsoring Agency Code	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546		15. Supplementary Notes Translation of: "Infrakrasnoye Izlucheniye Oblakov Venery", Academy of Sciences USSR, Institute of Space Research, Moscow, Report Pr-289, 1976, pp. 1-24.	
16. Abstract The thermal infrared emission of Venus measured by Venera-9 and Venera-10 displays significant asymmetry in the day-night direction. The emission of the night side corresponds to a brightness temperature of 244 °K. The brightest temperature of the day side is 233-234 °K. The extent of the upper layer of clouds, in which the thermal emission is formed, is 4-6 km. The altitude of the emitting layer above the surface of the planet (64-67 km) is determined from the brightness temperature and the existing models of the atmosphere of Venus. In some cases, correlation is noted between the inhomogeneity and the details of the ultraviolet image. The day side temperatures strangely coincide with the freezing point of sulfuric acid at a concentration of 66-77%.			
17. Key Words (Selected by Author(s))		18. Distribution Statement Unclassified - Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 23	22.

## INFRARED EMISSION OF THE CLOUDS OF VENUS

L. V. Ksanfomaliti, Ye. V. Dedova,  
L. F. Obukhova, N. V. Temnaya,  
G. F. Filippov

The thermal infrared emission of Venus measured by Venera-9 and Venera-10 displays significant asymmetry in the day-night direction. The emission of the night side corresponds to a brightness temperature of  $244^{\circ}\text{K}$ . The brightest temperature of the day side is  $233\text{--}234^{\circ}\text{K}$ . The extent of the upper layer of clouds, in which the thermal emission is formed, is 4-6 km. The altitude of the emitting layer above the surface of the planet (64-67 km) is determined from the brightness temperature and the existing models of the atmosphere of Venus. In some cases, correlation is noted between the inhomogeneity and the details of the ultraviolet image. The day side temperatures strangely coincide with the freezing point of sulfuric acid at a concentration of 66-77%.

The two radiometers of the same type mounted on board the Venera-9 and Venera-10 have an angular resolution of 1.2 deg (0.02 rad), and threshold sensitivity of about  $70^{\circ}\text{K}$  and a range of 8-13 and 18-23 $\mu$ . Rejection of the interval 10-18  $\mu$  prevents distortion of the results due to emission of the higher layers of the atmosphere in the carbon dioxide band at 15  $\mu$ . The wavelength of the maximum of the emission of Venus occurs at the middle of the radiometer range. 43\*

The thermal emission of Venus is formed in the upper layer of the cloud cover of the planet. The brightness temperature of the emitting layer is determined from the measured brightness in the indicated range. The radiation balance of the planet depends on this temperature. Among the problems of the experiment were the study of the limb darkening effect, investigation of local inhomogeneities and

---

\*Numbers in the margin indicate pagination of original foreign text.

their relation with phenomena observed at other wavelengths and study of longitude and latitude effects.

The altitude of the upper part of the cloud layer, from which the emission received by the radiometer originates, was found on the basis of the brightness temperatures obtained in the experiment with the help of the models of the atmosphere of Venus proposed by Marov and Ryabov [1] and Moroz [2]. This altitude is 64-67 km and refers to the level where the optical thickness in the infrared reaches unity. The physical properties of the emitting medium are tentatively the following: The pressure is 50-130 mb, the density is  $(0.13-0.18) \cdot 10^{-3} \text{ gm/cm}^3$ . The concentration of particles of the cloud layer is  $40-200 \text{ cm}^{-3}$  [3], their average diameter is about  $2.2 \mu$ , while the dispersion of the sizes is very small [4]. The most probable composition of the particles is sulfuric acid with a concentration of about 75% [5]. Its total concentration with respect to  $\text{CO}_2$  is about  $3 \cdot 10^{-5}$ . The infrared emission is formed in a rather extended layer, whose brightness temperature  $T_B$  practically coincides with its kinetic temperature [6].

Since the temperature of the outer part of this layer is somewhat lower, the limb-darkening effect arises similar to that described in the literature. It was shown in the work of Pollack and Sagan [7] that within definite limits of the cosine  $\mu_2$  of the zenith distance of the observer, all darkening models can be reduced to the law:

$$B_{\text{IR}}(\mu_2) = k \cdot \mu_2^\alpha \quad (1)$$

where  $k$  and  $\alpha$  are constants. Another version of representing  $B_{\text{IR}}(\mu_2)$  which follows from the approximate solution of the radiation transmission equation, was used by Chase, et al. [8]. The darkening coefficient  $D$  defined by the "boundary temperature"  $T_0$  for  $\mu_2 = 0$  was introduced here:

$$B_{\text{IR}}(T_0) = B(1.0) \cdot (1 - D) \quad (2)$$

where  $B(1.0)$  is the brightness at the point with  $\mu_2 = 1$ . Then:

$$B_{\text{IR}}(\mu_2) = B(1.0) \cdot [(1 - D) + D \mu_2] \quad (3)$$

Neglecting scattering, which is evidently valid for this problem, and using the Planck nature of the source function, one can find the absorption coefficient per unity path length. In any case, the contribution of the emission from colder layers increases with decreasing  $\mu_2$ , which also determines the characteristic form of the curves  $B_{IR}(\mu_2)$  or  $B_{IR}$  as a function of the distance from the center of the disk of the planet.

A few words about the models with which the altitude of the clouds was evaluated. Figure 1 presents five temperature-altitude curves for various models and also for the measurements of Venera-9, Venera-10 and the radio eclipse data of Mariner-10. Our estimates of the altitudes are based on the section of the curve indicated by points, closest in position to the models of Marov and Ryabov, Moroz and NASA-11. It is possible that the estimate of the altitudes will vary somewhat with further refinement of the models. /5

It was indicated in the express-communication about our experiment [9], in analyzing the obtained data, that their interpretation leads to the alternative: either to assume the evening temperature to be significantly greater than the day or to use two darkening laws different  $D$  for the day and evening sides. The first results obtained with the help of Venera-9 and Venera-10 showed that a temperature of about  $233^\circ\text{K}$  is consistently recorded in the day zone for values of  $\mu_2 \approx 1$  (Figs. 2 and 3). These measurements practically coincided with the averaged results of previous (ground-based) determinations [6]. However, the radiometric experiment on Mariner-10 in 1974 gave a temperature of  $255^\circ\text{K}$  in the range  $45\mu$  [8,10], while the first of these works indicated a small probability of significant divergences for the brightness temperatures at 10 and  $45\mu$ .

There are evidently two causes of the differences in the determination of the temperatures. One of them was shown in [9] in favor of our measurements. Its sense is that measurements from spacecraft require determination of two unknowns: the brightness temperature of the emitting layer and the law of limb darkening. The temperature can be determined strictly if the cosine of the zenith angle of the craft

$\mu_2$  is unity. As far as we know, this condition was not satisfied on Mariner-1. As was shown later, the darkening law for Venus cannot be determined by an isolated measurement. Another cause of the divergence is the thermal asymmetry of Venus, which is the content of this article.

Two different darkening laws for day and night were used in work [9]. It was learned soon after that this does not eliminate all difficulties. While all remained comparatively favorable from the day zone and the data fit well in dependence (3), this approximation was unsatisfactory in the zone of the night terminator (evening) (Fig. 4). Measurements of Venera-10 of November 6, 1975, are presented here. The temperatures in the day zone lie at the level  $233^\circ\text{K}$ . However, with consideration of  $\mu_2$ , they are higher in the evening zone. Dependence (1) presented in logarithmic form in Fig. 5,  $\lg B_{\text{IR}}$  as a function of  $\lg \mu_2$ , shows that the value of  $\alpha$  does not remain constant even in narrow zones of  $\mu_2$ . Further operation of the apparatus of Venera-9 and Venera-10 did not leave doubt about the stable increase in the night temperatures over the day. Figures 6 and 7 show the results of measurements obtained with increasing phase angle of the planet up to  $59$  and  $64^\circ$  (November 1 and 9, 1975, respectively). Already in Figure 6, the night temperatures sharply increase near the terminator for the same values of  $\mu_2$  in the day and night zones. This effect is expressed significantly more strongly in Fig. 6. With consideration of  $\mu_2$  in the evening zone, the temperatures reach  $236^\circ\text{K}$  for  $232^\circ\text{K}$  in the day zone. The asymmetry of the curve is common for all the profiles of Figs. 2, 3, 6 and 7. A completely new pattern can be seen in Fig. 8. The curve is symmetric. This is the night zone of the planet. The values of  $\mu_2 = 0.999$  made it possible to determine reliably the night temperature as  $244^\circ\text{K}$  with close darkening laws both in deep night as well as in the evening zones. Representation of

$B_{\text{IR}}(\mu_2)$  in the form of (3) gives a good approximation with:

$$B_{\text{IR}}(\mu_2) = B(I,0)(0,50 + 0,50 \cdot \mu_2) \quad (4)$$

for the day zone and:

$$B_{\text{IR}}(\mu_2) = B(I,0)(0,60 + 0,40 \cdot \mu_2) \quad (5)$$

for the night zone, while the approximation is better for the day zone. The form of (1) also represents well the results for the night zone:

$$B_{\text{IR}}(\mu_2) = B(I, 0) \cdot \mu_2^{\alpha} \quad (6) \quad \underline{71}$$

Values of  $\alpha$  from 0.30 to 0.43 were obtained on different days for the day part of the planet. The extent of the layer corresponding to an optical thickness of unity is 4-6 km. These results are considered in more detail in [11].

The transition of the cloud layer to the night mode of temperatures begins already for a drop of the sun to  $30-85^\circ$  above the horizon. It still remains unclear what the conditions are in the zone of the morning terminator, there are practically no possibilities for large differences: the left part of the zone represented in Fig. 8 was under night conditions for about 25 earth days. We consider it most likely that because of the great length of the solar day, the temperature distribution is quasi-stationary in nature and is similar for the evening and morning zones. (It would be more speculative to make the same assumption about the high latitude regions.) Thus, the "night mode" probably encompasses significantly more than  $180^\circ$  in longitude. The dependence of the brightness temperature on longitude, latitude and angle of the sun will undoubtedly be of interest. The studied traces encompass a zone of  $\pm 35^\circ$  in latitude. The investigations of the polar zones, which were performed from orbit by Venera-9 and Venera-10 only under conditions of relatively small  $\mu_2$  leave the possibility of some ambiguity in interpretation. A satellite of Venus in a circular polar orbit would be the ideal apparatus for solving this problem.

Thus, the thermal emission of Venus is asymmetric (Fig. 9). The temperatures of the day zone in the region of latitudes  $\pm 35^\circ$  are  $233-234^\circ\text{K}$ , and  $244^\circ\text{K}$  in the night zone at the same latitudes. This is the second reason for the differences in measurements on the apparatus of Venera-9 and Venera-10 on the one hand and Mariner-10 on the other, since the route of Mariner-10 was mainly over the night side

of the planet.

The symmetry of both branches of the curves  $B_{IR}(\mu_2)$  obtained /8  
by Taylor [10] is evidently a consequence of the significant coverage  
of the planet by the "night mode." In our measurements, this symmetry  
is observed only in the night zone. There were sequences in which day,  
evening and night were covered, which gave curves with strongly diverg-  
ing branches (Fig. 10).

The stable difference of the day and night temperatures has an  
obvious relation with the interception of part of the solar radiation  
by the cloud layer. We assume a probable cause of the decrease in day  
temperatures to be the appearance of intense convective currents in  
the day zone, which carry part of the radiating substance into the  
zone above the clouds at an altitude of 3-4 km. This leads to a  
change in the exponent  $\alpha$ . It remains unclear whether this cause is  
natural. There is a strange coincidence noticed by L. Ksanfomaliti.  
Figure 11 presents the phase diagram of the sulfuric acid solution  
as a function of its concentration. The curve is taken from the work  
of A. Young [5]. The level of the day temperature lies precisely at  
the sharp bend in the curve at the point with the sulfuric acid con-  
centration of 66-77%. The refractive index for the solution with  
concentration 70% and temperature  $233^{\circ}\text{K}$  is 1.44, which coincides  
ideally with [4]. However, the temperature of  $250^{\circ}\text{K}$  taken by A. Young  
distinguishes practically nothing with these points of the curve,  
whereas, our values ( $233^{\circ}\text{K}$ ) can be interpreted as evidence that the  
upper boundary of the clouds of Venus is somehow related just to this  
point of the phase diagram. There are evidently mechanisms stabiliz-  
ing this level, at least in temperature. One follows from the temper-  
ature-pressure dependence of water vapor over sulfuric acid presented  
in the same work of A. Young. The pressure of water vapor over the  
acid with a concentration of 68% is three times greater than for a  
concentration of 70%. But the freezing point at 68% is above that for  
70%. The difference in pressures provides for the motion of water /9  
vapor downward to the main mass of the absorber. The freezing drops  
with lower concentration transmit the excess water vapor to other  
drops, thereby enriching the acid and maintaining the liquid state

(the arrow in Fig. 11). The large latent heat of the phase transition in the drops of  $H_2SO_4$  together with the intense mixing imparts a steady-state nature to the process, which is also aided by the sharp increase in the rate of precipitation of drops from the more rarified upper layers according to the data of [3]: while  $10^7$  sec are required for their drop by one altitude scale height (5 km) at an altitude of 68 km, only  $10^6$  sec is required at an altitude of 78 km. The capability of sulfuric acid for intense supercooling is also an important fact noted by Pollack in the discussion to work [12].

The mechanism described above permits a completely independent evaluation of the concentration of water vapor (at the altitude of the emitting layer) from its pressure. This quantity is  $0.9 \cdot 10^{-4}$  with respect to the pressure of carbon dioxide above sulfuric acid with a concentration of 70% and coincides well with other determinations, for example, [13]. The temperature of  $250^\circ K$  taken by many American investigators would give an unacceptably high pressure, almost an order of magnitude higher.

Let us now turn to the details in the curves  $B_{IR}$ . Extended regions with temperatures decreased by  $0.8-1.4^\circ$  are observed in several sequences. The length of these zones is up to 5000 km; we see such a pattern in Fig. 3. There is not doubt about their nature: these are local elevations of the emitting layer. According to the estimate of A. Young, such elevations can exceed the scale height. It remains surprising in this case that the mechanism stabilizing the temperature operates so effectively.

There is also great interest in the fine-scale details of the curves. They are clearly traced on individual profiles, but are occasionally absent on other traces investigated with the same resolution. Their amplitude does not usually exceed  $2^\circ K$ . According to the available statistics, the resolution under orbital conditions permits obtaining details, curves of three types are observed:

- with weakly expressed details (Fig. 7);
- with a few periodic details (Fig. 3; Fig. 6).

In these cases, the small waves on the curves have a spatial extent of 150-180 km and are separated by intervals of about 1500 km; their

correlation with ultraviolet details is observed. As was shown in [9], one can assume a relation between these waves and the dynamics of the atmosphere of Venus;

-- with a large number of details with unclear period. The typical dimension of the details (50-80 km) is an interesting fact. A sharp increase in resolution does not give any additional details (Fig. 8).

Further tasks of the experiment are the study of the night and morning zones and also of the high-latitude regions of the planet.

#### REFERENCES

1. M. Ya. Marov, L. L. Ryabov. Model' atmosfery Venery ("Model of the Atmosphere of Venus"). IMP preprint No. 112, Moscow, 1974.
2. V. I. Moroz. Rabochaya model' atmosfery Venery ("Working Model of the Atmosphere of Venus"). IKI preprint No. 162, Moscow, 1973.
3. P. Gierash. In the Coll: The Atmosphere of Venus, 2-7, publ. NASA Sp 382, 1975.
4. J. E. Hansen, J. W. Hovenier, J. Atm. Sci., 31, 1137-1160, 1974.
5. A. T. Young, Icarus, 18, 564-582, 1973.
6. V. I. Moroz. Uspekhi fizicheskikh nauk, 104, 2, 255-296, 1971.
7. J. B. Pollack, C. Sagan, J. Geophys. Res., 70, 18, 4403-4426, 1965.
8. S. C. Chase, E. D. Miner, D. Morrison, G. Münch, G. Neugebauer, Science, 183, 1291-1292, 1974.
9. L. V. Ksanfomaliti. Pis'ma v AZh, 2, 2, 11-17, 1976.
10. F. W. Taylor, J. Atm. Sci., 32, 6, 1101-1106, 1975.
11. L. V. Ksanfomaliti. Pis'ma v AZh, in press.
12. S. Wofsi, In the Coll: The Atmosphere of Venus, 25-27. publ. NASA, Sp.382, 1975.
13. W. Rosson, C. Sagan, J. Atm. Sci., 32, 6, 1164-1176, 1975.

- Fig. 1. Temperature-altitude curves for different models. The points indicate sections of the curve on which the altitude estimates in the article are based.
- Fig. 2. Curve of brightness  $B_{IR}$  and brightness temperature  $T_B$  in the infrared region according to the data of Venera-10, October 29, 1975. The approximation by two curves of the type (3) indicated by points deviates significantly from the curve at the edges. The lower curve  $B_{IR}$  is the reading of the control ultraviolet photometer. The circles in the upper part is the resolution scaled on the time axis in terms of the velocity normal to the sighting line and the slant range. The day temperature is  $234^{\circ}\text{K}$ . The phase angle is  $57^{\circ}$ .
- Fig. 3. October 26, 1975, Venera-9 transmitted measurements of the brightness  $B_{IR}$  and temperature  $T_B$  after composing the condensed profile with a decrease in temperature by  $1-1.4^{\circ}$ . The typical periodic waves on the curve are separated by intervals of about 1500 km. The waves are evidently correlated with details of the ultraviolet profile. The phase angle is  $55^{\circ}$ .
- Fig. 4. Curves of  $B_{IR}$  and  $B_{UV}$  of Venera-10 for November 6, 1975. The left, evening part of the profile cannot be represented in the form (3). The day temperature is  $233^{\circ}\text{K}$ . The phase angle is  $62^{\circ}$ .
- Fig. 5. Change in the exponent  $\alpha$  in formula (1) represented as  $\Delta \lg B_{IR} / \Delta (\lg M_2 + 1)$ . Venera-10, November 6, 1975.  $B_{IR}$  in arbitrary units.
- Fig. 6. Measurements of Venera-9 of November 1, 1975. The growth in  $B_{IR}$  and  $T_B$  to the left from the maximum value is well noted. Periodic waves are noted. The phase angle is  $59^{\circ}$ .
- Fig. 7. Sharp growth in  $B_{IR}$  and  $T_B$  to the left and the maximum value and the terminator. Venera-9, November 9, 1975. The phase

angle is  $64^\circ$ .

Fig. 8. Night side of Venus. The curve is practically symmetric on both sides from  $\mu_{2\max}$ . Venera-9, November 13, 1975, the phase angle is  $122^\circ$ . The maximum value of  $\mu_2$  corresponds to  $244^\circ\text{K}$ . The measurements were performed with high resolution.

Fig. 9. Diagram of the distribution of the constant components of the brightness of the thermal emission of Venus from the surface of the cloud layer in the region of latitudes  $+35^\circ$ . The variation of the brightness is approximately linear in the transition zone. View from the North Pole. /13

Fig. 10. Day and night branches of the profiles  $B_{\text{IR}}$ , presented in Fig. 3 and Fig. 6 as a function of  $\mu_2$ . Curve 1 is the same as in Fig. 8, 2 is the same as in Fig. 4.

Fig. 11. Phase diagram of the water-sulfuric acid system [4]. The day temperatures of  $233^\circ$  coincide with the position of the minimum of the curve at 70-76% solution. The arrow indicates the change in composition of the drops and the decrease in its freezing temperature.


Models of the atmosphere

of Venus:

- 1. Marov, Ryabov, 1972
- 2. Marov, Ryabov, 1974
- 3. Moroz, 1973
- 4. NASA I
- 5. NASA II

Data:

- Mariner 10
- Venera-9
- Venera-10



ORIGINAL PAGE IS  
OF POOR QUALITY

Figure 1



Y-557, 21 82 24 1075,




Figure 3

9-62- 910 106 64125




Figure 4





Figure 5

Venera-9  
November 1  
1975

Phase angle 53°



CM. u. ster  
2

$B_{UV}$   
20  
10

$t_{MIN}$   
25  
20  
15  
10  
05  
8'00"  
55  
50  
45  
04 05 06 07 08 09 095 105 115 125 135 145 155 165 175 185 195 205 215 225 235 245 255

Figure 6

KNOX, M. A. & S. J. R.

cm<sup>2</sup> meter  
mW

B<sub>UV</sub>  
-20

10




Figure 7






Figure 9




Figure 10




Figure 11