

N79-13626

THE EFFECTS OF STRONG SHOCK WAVES ON MORTALITY RATES
AND PERCENTAGES OF PULMONARY LESIONS IN RATS AS
A FUNCTION OF THE NUMBER OF EXPOSURES

Patrick Vassout and Georges Parmentier

Translation of "Etude des effets des ondes de choc fortes sur
les taux de letalite et les pourcentages de lesions
pulmonaires chez le rat en fonction du nombre d'expositions",
Institut franco-allemand de recherches (I.S.L.) Sant-Louis
(France), Unpublished report, 1978, pp. 1-13.



1. Report No. NASA TM-75598	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle THE EFFECTS OF STRONG SHOCK WAVES ON MOR- TALITY RATES AND PERCENTAGES OF PULMONARY LESIONS IN RATS AS A FUNCTION OF THE NUMBER		5. Report Date December 1978	6. Performing Organization Code
7. Author(s) OF EXPOSURES. Patrick Vassout and Georges Parmentier Institut of Franco German Research (I.S.L.) Sant-Louis, France		8. Performing Organization Report No.	9. Work Unit No.
9. Performing Organization Name and Address Leo Kanner Associates Redwood City, California 94063		11. Contract or Grant No. NASW-3199	13. Type of Report and Period Covered Translation
12. Sponsoring Agency Name and Address National Aeronautics and Space Adminis- tration, Washington, D.C. 20546		14. Sponsoring Agency Code	
15. Supplementary Notes Translation of "Etude des effets des ondes de choc fortes sur les taux de letalite et les pourcentages de lesions pulmonaires chez le rat en fonction du nombre d'expositions", Institut franco-allemand de recherches (I.S.L.) Sant-Louis (France), Unpublished report, 1978, pp. 1-13.			
16. Abstract This study of pulmonary lesions and mortality rates caused in rats by exposure to strong shock waves of constant duration revealed that with regard to the pulmonary lesions, twice the number of exposures is compensated for by quartering the overpressure of the wave crest, and with regard to the mortal- ity rates, it revealed that halving the overpressure of the wave crest is offset by a 20-fold increase in the number of exposures.			
17. Key Words (Selected by Author(s))		18. Distribution Statement Unclassified - Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages	22. Price

THE EFFECTS OF STRONG SHOCK WAVES ON MORTALITY RATES AND PERCENTAGES
OF PULMONARY LESIONS IN RATS AS A FUNCTION OF THE NUMBER OF
EXPOSURES

Patrick Vassout and Georges Parmentier

Introduction

The effects of strong blast waves have been studied in particular ^{*/2} since the Second World War. In general, authors have described the relationships which exist between such and such a parameter of the blast wave and the percentages of pulmonary or intestinal lesions, as well as the mortality rates, observed in subjects of a given species for a single exposure.

The literature reviews in [1] and [2] show that so far the parameter "number of exposures" has not been dealt with by the several investigators in question. Indeed, it would not seem very realistic to imagine that an individual could be subjected to a succession of pure blasts. The appearance of new weapons based on the principle of gas detonation, as well as the development of weapons which produce very high overpressure levels at the start of the blast pose the problem concerning the effect of the number of exposures to a strong blast wave on the extent of pulmonary lesions and on mortality rates.

We have therefore undertaken to study the "number of exposures" parameter on pulmonary lesions and mortality rates in the rat [3].

1. Experimental Apparatus

The experimental animals were female rats with an average body weight of 100 g. Each animal was placed in a wire mesh cage [1]. The cages were placed symmetrically around the explosive charge. With this arrangement it was possible to apply the same blast wave to each animal while changing as little as possible the pressure profile of the wave.

The explosive used was "plastit" made by Dottikom Co., Switzerland.

*Numbers in the margin indicate pagination in the foreign text.

1.1. Determination of the Amounts of Explosive Required

By applying the laws of similarity valid for spherical explosives [4] [5], we determined the amounts of explosive necessary to obtain the desired overpressures while maintaining a constant duration of the positive first phase t_0 .

1.2. Recording Pressure Changes

The pressure signals (Fig. 2) were recorded by means of KISTLER /3 6031 piezoelectric dynamic pick-ups placed in the tapered waves. These pick-ups were connected to BIOMATION 805 transistorized analog digital signal recorders. The pick-ups were installed at the same distance from the explosive charge as the experimental animals and measured the pressure in free space. The animals, placed on [word illegible], were exposed to the incident pressure and the pressure reflected by the ground. The pressure levels discussed here always refer to those which existed at the level of the animals. (Overpressure measured in free space multiplied by the reflection factor)

2. Results

2.1. Mortality Rates as a Function of the Number of Exposures

2.1.1. Determination of the Median Lethal Dose for Rats

The median lethal dose (LD_{50}) is the crest overpressure which for a given length of time T_0 for the incident blast wave causes death in 50% of the animals exposed at the end of a given length of time after exposure, for example 24 hours.

Based on the studies of Richmond and White (1962) [6], done on many species of animals, we can estimate the crest overpressure corresponding to the LD_{50} for rats for a given t_0 of the blast wave ($LD_{50} = 3$ bars for $t_0 = 2$ ms). We have chosen as [word illegible] of the blast wave the value $t_0 = 2$ ms. Indeed, according to these authors, beyond this length of time this parameter no longer has any effect on the mortality rate in rats.

For this 2-ms length of time we have studied the mortality rate in rats for a single exposure to an overpressure blast wave between

3 and 5 bars (Fig. 3). It should be noted that all of the overpressure less than 3 bars did not cause any mortality in our experiments.

It appears that the LD_{50} for rats, 24 hours after a single exposure, is located around 3.5 bars. We also find that the difference in overpressure necessary to go from a mortality rate of 10% to one of 90% five minutes after exposure is 1.5 bars in rats.

2.1.2. Effect of the Number of Exposures on Rat Mortality Rates 74

For the purpose of determining relationships between mortality rates, the number of exposures and crest overpressure, series of 40 animals were exposed to blast waves of $t_0 = 2$ s. produced at regular 4-minute intervals.

Fig. 4 shows the results obtained in the overpressure range between 2 and 4 bars. The dotted line curve which represents the LD_{50} reveals that a twofold increase in the level of the crest overpressure is offset by a 20-fold decrease in the number of exposures.

2.2. Percentages of Pulmonary Lesions Due to Exposure to Strong Blast Waves

2.2.1. Definition of Lesion Criteria Used in our Experiments

Fig. 3 shows that a single exposure to a wave overpressure blast wave less than 3 bars does not cause any mortality. However, autopsy reveals that the animals have hemorrhagic pulmonary lesions ranging from simple petechia to a completely hemorrhagic lobe. We have taken as the lesion threshold the small hemorrhage isolated on a single lobe according to the criteria proposed by Richmond et al. [7].

2.2.2. Determination of the Overpressure which Causes Pulmonary Lesions in 50% of the Experimental Animals

As in previous experiments, we first of all studied the influence of a single exposure to varied crest overpressures on the rate of pulmonary lesions.

The overpressure which causes the appearance of pulmonary lesions in 50% of the experimental animals is situated around 500 mbars (Fig. 5). An overpressure of 2.5 bars lasting for the same amount of time (2 ms) causes lesions to appear in all of the animals exposed.

2.2.3. Effect of the Number of Exposures on the Percentage of Animals Showing Pulmonary Lesions

/5

Groups of 40 animals were exposed to series of blast waves all lasting $t_0 = 2$ ms and of overpressures ranging between 32 and 500 mbars. The results obtained are plotted in Fig. 6. Note that for a constant percentage of animals with lesions (dotted line curve), a two-fold increase in the number of exposures must be compensated for by a quartering of the crest overpressure. This result shows a different relationship between the wave overpressure and the number of exposures experienced from [word illegible] observed on the basis of mortality rates.

3. Trial Extrapolation for a Theoretical Animal Weighing 70 kg and Conclusion

Many authors such as Zuckerman 1941 [10], White 1959 [9] Richmond 1968 [7] and Jones 1969 [9] think that it is possible to extrapolate the results obtained in animals to man. They indicate that the LD_{50} applicable to man would be on the order of 10 bars for a blast wave lasting 2 ms (t_0). These authors also think that the threshold for the appearance of pulmonary lesions in man surround one-fourth of the LD_{50} , or around 2.5 bars.

As for the effect of the number of exposures on mortality rate, extrapolations of our results to man indicates that twenty exposures to a crest overpressure of 5 bars lasting 2 ms (t_0) experienced at intervals on the order of 5 [word illegible] would cause death in 50% of the subjects exposed.

As for the effect of the number of exposures on the appearance of pulmonary lesions, we fear that lesions might appear in man for crest overpressure values plainly less than those currently accepted (2.5 bars for a t_0 of [word missing] where the subject is exposed to a series of strong blast waves).




Fig. 1. Arrangement of Rat Cages




Fig. 2. Typical recording of pressure signals for an explosive mass.

A Poids de "Plastit" [g]	Distance [cm]	Ap [bar]	B Temps écoulé après une exposition							c nbre d'ani- maux
			5mn	30mn	1h	2h	4h	24h		
1096	278	3	0	8	8	11	28	36		
1200	270	3.5	9	15	34	38	53	68		
1371	267	4	35	35	35	45	58	31		
1560	267	4.5	49	74	77	89	91	35		
1770	269	5	97	100				35		

Fig. 3. Mortality rates observed after an exposure to blast waves of variable Δe and constant t_0 (2 ms) in rats weighing 100 g.

Key: a. Weight of "plastit" in grams
b. Elapsed time after explosion
c. Number of animals




Fig. 4. Effect of the number of blast waves experienced as a function of overpressure with a constant t_0 (2 ms) on mortality rate in rats.

- Key:
- a. Number of blast waves
 - b. LD₅₀
 - c. Mortality rate
 - d. 24 hours after exposure




Fig. 5. [Caption illegible]
 Key; a. Percent of animals with lesions




Fig. 6. Effect of the number of blast waves experienced as a function of the overpressure of constant t_0 (2 ms) on the percentage of rats with lesions.

- Key:
- a. Number of blast waves
 - b. 50% of animals with lesions
 - c. Percentage of rats with pulmonary lesions

References

1. Rigaud, P., "Effect of Blast Waves on Living Creatures in General." Paper presented to the Group of French-Speaking Acousticians (GALF) in May 1970. Notice ISL NB 2/71, (1971).
2. Rigaud, P. and A. Dancer, "Effect of Blast Waves on Living Creatures," Notice ISL NB 6/73, (1973).
3. Vassout, P., G. Paramentier and A. Dancer, "The Effect of Strong Blast Waves on the Mortality Rate and the Appearance of Pulmonary Lesions in Rats," ISL Report R 922/78, (1978).
4. Froböse, M., "Propagation Functions of Spherical Aerial Blast Waves," ISL Report R 3/68 (1968).
5. Froböse, M. and O. Wünsche, "Mortality of Albino Rats Exposed to an Aerial Blast Wave as a Function of the Direction of the Blast and the Shape of the Pressure Curve," ISL Report R 2/59, (1959).
6. [No No. 6 listed]
7. Richmond, D.R., E.G. Damon, G.I. Bowen, E. Fletcher and C.S. White, "Air Blast Studies with Eight Species of Mammals," Lovelace Foundation for Medicine Education and Research, New York, USA (1966) DASA 1854.
"The Relationship Between Selected Blast Wave Parameters on the Response of Mammals Exposed to Air Blast," A. of New York Acad. of Sciences 152, 103-121, (1968).
"A Tentative Estimation of Man's Tolerance to Overpressures From Air Blast," Lovelace Foundation for Medicine Education and Research, New York, USA (1962) DASA 1335.
8. White, C.S. and D. R. Richmond, "Blast Biology," Lovelace Foundation for Medical Education and Research, New Mexico, T.I.D. 5764, (1959).
9. Jones, R.D., D.R. Richmond, and R.E. Flechter, "A Reappraisal of Man's Tolerance to Indirect Air Blast Injury," Lovelace Foundation for Medicine Education and Research, New Mexico, USA, (1969).
10. Zuckerman, S., "Experimental Study of Blast Injury to the Lungs," Lancet II, pp. 219-238, (1940).