

N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED
IN THE INTEREST OF MAKING AVAILABLE AS MUCH
INFORMATION AS POSSIBLE

"Page missing from available version"

CHANGES IN GAS EXCHANGE, TISSUE RESPIRATION
AND GLYCOLYSIS IN RATS DURING HYPOKINESIA

(NASA-TM-76149) CHANGES IN GAS EXCHANGE,
TISSUE RESPIRATION AND GLYCOLYSIS IN RATS
DURING HYPOKINESIA (National Aeronautics and
Space Administration) 11 p HC A02/MF A01

N80-32055

Unclass
CSCL Doc G3/51 28000

Translation of "Zminy Hazoobminu, Tkanunnoho Dykhannya ta
Hlikolizu u Shehuriv pry Hipokinezii", *Fiziolohichnyy Zhurnal*,
Vol. 22, No. 3, 1976, pp 414-417



STANDARD TITLE PAGE

1. Report No. NASA TM-76149	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle CHANGES IN GAS EXCHANGE, TISSUE RESPIRATION AND GLYCOLYSIS IN RATS DURING HYPOKINESIA.		5. Report Date MAY 1980	6. Performing Organization Code
		7. Author(s) L. V. ZORYA	8. Performing Organization Report No.
9. Performing Organization Name and Address BOITRAN Box 5456 Santa Barbara, CA 93108		10. Work Unit No.	11. Contract or Grant No. NASW-2198
		12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546	
13. Type of Report and Period Covered Translation		14. Sponsoring Agency Code	
15. Supplementary Notes Translation of "Zminy Hazoobminu, Tkanunnoho Dykhannya ta Hlikolizu u Shchuriv pry Hipokinezii", Fiziolohichnyy Zhurnal, Vol. 22, No. 3, 1976, pp 414-417			
16. Abstract This article presents the results of an experiment which studied changes in oxygen balance under conditions of hypokinesia in rats. The experimental method used and results obtained are discussed. In summary, the effect of the stress during hypokinesia is expressed most clearly in the changes of general gas exchange, and in the intensity of liver and myocardial tissue respiration.			
17. Key Words (Selected by Author(s))		18. Distribution Statement Unclassified - Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 11	22. Price

L. V. Zorya

Changes in Gas Exchange, Tissue Respiration and Glycolysis
In Rats During Hypokinesia

*/1

Extreme restriction of man's kinetic activity causes changes in oxygen balance of the organism, which basically brings about a lowering of oxygen consumption. There are published reports of a decrease in oxygen consumption and basic metabolism of healthy men bedridden for 15-20 days /4, 6, 16/, 70 days /10/, and 120 days /11/. However, there are conflicting reports describing a 57% increase in oxygen consumption of sportsmen's organisms following a 10-day hypokinesia /1/.

The reports of gas exchange studies in animals under condition of experimental hypokinesia are even more conflicting. On one hand, it was observed that a 30-day, and a 60-day restriction of locomotion in rats did not bring about appreciable changes in the overall gas exchange /12, 14/, or blood composition /8/. On the other hand, a 36% /13/, and a 50% /3/ increase in oxygen consumption after a 30-day hypokinesia has been reported.

It follows from the published data that a prolonged hypokinesia causes a change in the organism's metabolism. However, the nature of these alterations needs clarification. The duration of the gas exchange studies in animals varies between 30 and 120 days of hypokinesia in the reports of other authors. However, there is no description of gas exchange dynamics during the first month of restricted mobility.

We have studied oxygen consumption of the entire organism, and also tissue respiration and glycolysis in the myocardium, skeletal muscle and liver of rats 4, 8, 16 and 30 days after the commencement of hypokinesia.

*Numbers in margin indicate pagination of foreign text.

Experimental Methods

12

Experiments were performed with 233 white, noninbred rats of both sexes with initial body weight 130-180 g. In order to create conditions of hypokinesia, animals were placed in specially constructed individual cages which drastically reduced locomotive activity. Control rats were kept in the same room, in cages of an ordinary vivarian type. Animals were fed an ordinary rat food. The dynamics of animal body weight was observed.

Oxygen consumption was determined according to the procedure of Kalabukhov /5/ on days 4, 8, 16 and 30 of hypokinesia. The intensity of tissue respiration and glycolysis were studied in the Warburg apparatus at the same intervals.

Results and Discussion

Experiments have demonstrated that wavelike variations occurred in the general gas exchange in the rat during the 30-day hypokinesia. It can be seen in the Table that the mean value of oxygen consumption on day four of restricted mobility increased from the initial value of 0.159 ± 0.0079 L/100 g/hr to 0.186 ± 0.0127 L/100 g/hr ($p = 0.05$) which represents a 16.98% increase. During days 8 and 16 of the experiment, oxygen consumption was somewhat reduced, relative to the initial level, and during day 30 it went up. However, these variations proved not to be significant.

Table. Oxygen consumption ($M \pm m$) by the organism of rats during a 30-day hypokinesia (in L/100 g body weight/hr)

Experimental conditions	Day of Experiment			
	4	8	16	30
Initial values	0.159 ± 0.0079	0.183 ± 0.0097	0.177 ± 0.0161	0.138 ± 0.0123
after Hypokinesia	0.186 ± 0.0127	0.169 ± 0.0114	0.155 ± 0.0198	0.164 ± 0.0094
p	=0.05	>0.05	>0.05	>0.05

Similarly, wavelike but better defined changes in oxygen consumption were found when the intensity of tissue respiration was studied in the individual tissues (Figure 1). The most significant elevation in oxygen consumption from 3.00 ± 0.309 to $4.52 \pm 0.295 \mu\text{L O}_2/\text{mg dry tissue}/30 \text{ min}$ ($p < 0.01$) occurred in the liver during the fourth day of hypokinesia. Later, a tendency for the normalization of liver tissue respiration was observed, with an insignificant increase during day 30. Oxygen consumption increased in the myocardium from 1.96 ± 0.176 to $2.53 \pm 0.185 \mu\text{L O}_2/\text{mg dry tissue}/30 \text{ min}$ ($p < 0.01$) on day 16 of hypokinesia. During day 30 of the experiment, myocardial tissue respiration decreased to values below the initial ones ($1.76 \pm 0.135 \mu\text{L O}_2/\text{mg dry tissue}/30 \text{ min}$; $p > 0.05$). During the initial 16 days of hypokinesia there were insignificant variations of oxygen consumption values for the skeletal muscle. However, a lowering of tissue respiratory intensity from 1.63 ± 0.094 to $1.13 \pm 0.106 \mu\text{L O}_2/\text{mg dry tissue}/30 \text{ min}$ ($p < 0.01$) took place by day 30 of the experiment.

Changes in glycolysis were of a different nature: its intensity was gradually decreasing during hypokinesia in all experimental tissues (Figure 2). A definite decrease in the intensity of glycolysis on day four of the experiment was observed only in the myocardium (from 2.41 ± 0.225 to $1.81 \pm 0.100 \mu\text{L CO}_2/\text{mg dry tissue}/30 \text{ min}$; $p < 0.02$). Myocardial glycolysis decreased even more to 1.55 ± 0.082 ($p < 0.001$) during day eight, and on day 30 it reached $1.33 \pm 0.121 \mu\text{L CO}_2/\text{mg dry tissue}/30 \text{ min}$ ($p < 0.002$). In the liver and in the skeletal muscle, a definite lowering of glycolytic activity began on day eight of hypokinesia

in agreement with observations of Barbashova /2/, and are another indication of the condition of a harsh stress during this period. Portugalov et al. /15/ have noticed that the endocrine glands of




Fig. 1. Changes in the intensity of tissue respiration during a 30-day hypokinesia. Vertical axis, $\mu\text{L O}_2/\text{mg}$ dry tissue/30 min; abscissa, days of hypokinesia. a, liver; b, heart; c, skeletal muscle.




Fig. 2. Changes in the intensity of tissue glycolysis during 30-day hypokinesia. Vertical axis, $\mu\text{L CO}_2/\text{mg}$ dry tissue per 30 min. Abscissa, days of hypokinesia.

the organism work with maximal intensity during the first five days, followed by a period of adaptation to the new condition. Kovalenko et al./7/, studying the system adrenal cortex - hypophysis during hypokinesia concluded that changes during the initial period are the result of the emotional tension. Furthermore, the restriction of mobility becomes evident at the end of the second week.

In summary, the effect of the emotional stress during the initial period of hypokinesia is expressed most clearly in the changes of general gas exchange, and in the intensity of liver and myocardial tissue respiration. Changes in the metabolism of the skeletal muscle were of somewhat different nature. There

was no evidence of wavelike alterations: a lowering in the intensity of tissue respiration was detected only on day 30. However, lowered glycolysis was observed in this tissue much earlier, beginning on day eight of hypokinesia.

References

1. Ashcheulova, E.N., Yakhnovich, G.G., and Pudov, V.I. Dynamics of gas exchange of man under conditions of restricted movement of variable duration. In: Fiziologicheskiye problemy detrenirovanosti ("Physiological problems of decreased training"), Moscow, 1970, pp. 109-117.
2. Barbashova, Z.I., Zhukov, E.K., Baklanova, S.M., Borisova, L. Ya., Grigoryeva, G.I., Nesterov, V.P., Simanovskiy, L.N., Tarakanova, O.I., and Fedorov, V.V. Change in the resistance of the organism, functional condition of tissues and their biochemical processes during hypodynamia. In: Adaptatsiya k myshechnoy deyatelnosti i gipokinesiyi ("Adaptation to muscular activity and hypokinesia"). Proceedings of a symposium, Novosibirsk, 1970, p. 26-35.
3. Galushko, Yu.S., Abakumova, I.A., and Nitochkina, I.A. Connection between oxygen utilization and the level of unoxidized products in the urine of rats during hypokinesia. In: Aktualnyye voprosy kosmich. biol. i med. ("Current problems in cosmic biology and medicine"), Moscow, 1971, p. 77-78.
4. Kakurin, L.I., Katkovskiy, B.S., Machinskiy, G.V., Pometov, Yu.D., and Smirnova, G.I. Alterations in the gas and energy exchange of healthy man under the influence of low-calory

- nutrition and hypokinesia. In: Fiziol. problemy detrenirovannosti ("Physiological problems of reduced training"), Moscow, 1970, p. 121-126.
5. Kalabukhov, N.I. Metodika eksperimentalnykh issledovaniy po ekologiyi nazemnykh pozvonochnykh zhivotnykh ("Methods of experimental studies in ecology of terrestrial vertebrates"), Moscow, 1951. p. 75-77.
 6. Kotkovskiy, B.S. Effect of hypokinesia on the oxygen regimen of man during physical work. In: Kislородnyy rezhim organisma i yego regulirovaniye ("Oxygen regimen of the organism and its regulation"), Kiev, 1966, p. 285-290.
 7. Kovalenko, E.A., Popkov, V.L., Kondratyev, Yu.I., Mailyan, E.S., Galushko, Yu. S., Prokhonchukov, A.A., Kazaryan, V.A., Morozova, R.S., Serova, L.V., Potapov, A.N., Romanov, V.S., and Pishik, V.B. Changes in the functions of the organism during prolonged hypokinesia. Patol. fiziol. i eksper. terapiya, 1970, 6, p. 3-9.
 8. Lunev, I.Ya. Gas content and acid-base equilibrium of blood during a 10-day hypokinesia. In: Aktualnyye voprosy kosmich. biol. i med. ("Current problems in cosmic biology and medicine"), Moscow, 1971, p. 174-175.
 9. Mailyan, E.S., and Kovalenko, E.A. On tissue respiration under the influence of prolonged hypokinesia and hypergravitation. In: Sovremennyye problemy biokhimii dykhaniya i klinika ("Current problems in the biochemistry of respiration and its clinical aspects"). Proceedings, II All-Union Conference, Ivanovo, 1972, 1, p. 160-162.
 10. Mikheev, M.I., Sokolov, V.I., and Tikhonov, M.A. Some peculiarities of external respiration and gas exchange during

- prolonged hypodynamia. In: Problemy kosm. biol. ("Problems of cosmic biology"), Moscow, 1969, p. 71-78.
11. Motorin, Yu.A. Effect of a 120-day long clinico-statical hypokinesia on basal metabolism of man. In: Aktualnyye voprosy kosmich. biol. i med. ("Current problems of cosmic biology and medicine"), Moscow, 1971, p. 189-190.
 12. Nitochkina, I.A., and Galushko, Yu.S. Comparative evaluation of changes in gas exchange in white rats determined by two different methods during a prolonged hypokinesia lasting 60 and 100 days. In: Materialy III nauchn konfer. molodykh spetsialistov ("Proceedings, III Scientific conference of young specialists"), Moscow, 1969, p. 91-93.
 13. Nitochkina, I.A., Ryazhskiy, A.V., and Galushko, Yu.S. Change in body weight and oxygen utilization in rats of various age during a prolonged hypokinesia. In: Aktualnyye voprosy kosmich. biol. i med. ("Current problems in cosmic biology and medicine"), Moscow, 1971, p. 200-201.
 14. Popkov, V.L., Mailyan, E.S., Galushko, Yu.S., Kovalenko, E.A., Zaytseva, E.I., Nitochkina, I.A., Stalova, L.V., and Ryazhskiy, A.V. Changes in gas exchange, gas homeostasis and tissue respiration in rats during prolonged hypokinesia. Fiziol. zhurnal SSSR, 1970, 56, #12, p. 1808-1812.
 15. Portugalov, V.V., Ilyina-Kakuyeva, E.I., Artyukhina, T.V., Gotlib, V.L., and Starostin, V.I. Changes in some endocrine glands and secretion of hypothalamus nuclei during hypokinesia. In: Eksper. issledovaniya gipokinezii, izmenennoy gas. sredy, uskoreniy, peregruzok i dr. faktorov ("Studies on hypokinesia, altered gaseous environment, accelerations, overloading and other factors"), Moscow, 1968, p. 29-33.

16. Smirnova, G.I. Changes in oxygen utilization by the organism of man under the influence of organic nutrition, hypokinesia and acceleration in the centrifuge. In: Aktualnyye voprosy kosmich. biol. i med. ("Current problems in cosmic biology and medicine"), Moscow, 1971, p. 242-243.

Chair of Pathological Physiology
Tarnopil Medical Institute

Received
14 March, 1975