

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

DRA

ORIGINAL PAGE IS  
OF POOR QUALITY

(NASA-CR-172879) PLANETARY ATMOSPHERES  
PROGRAM Final Technical Report, 15 May 1982  
- 15 May 1983 (Lewis (John S.) Associates,  
Inc.) 71 p HC A04/MF A01 CSCL 03B

N83-30339

Unclas  
10114

G3/91

FINAL TECHNICAL REPORT

to the

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

Planetary Atmospheres Program

on

CONTRACT NASW-3625

for the year


15 May 1982 to 15 May 1983

from

J. S. Lewis Associates, Inc.

5010 W. Sweetwater Drive

Tucson, AZ 85745



ORIGINAL PAGE IS  
OF POOR QUALITY

CONTENTS

PAGE

GENERAL INTRODUCTION.....2

SECTION I: Non-Solar Compositional Models of the  
Troposphere of Jupiter.....3

SECTION II: Venus:Halide Cloud Condensation and Volatile  
Element Inventories.....

SECTION III: Shock-Wave Processing of Interstellar  
Cloud Materials.....

ORIGINAL PAGE IS  
OF POOR QUALITY

## General Introduction

This is the final report on a program of research designed for execution on microcomputers. At its original inception, the program was intended to run for a period of three years, and to involve

- a. theoretical work on compositional trends in the systems of the Jovian planets, including both the radial chemical variations within each satellite system and the compositional trends from Jupiter through Saturn and Uranus to Neptune,
- b. geochemistry of the moderately volatile elements on Venus, for the purpose of determining whether the Pioneer Venus and Venera compositional measurements within the Venus troposphere can reveal whether Venus originally had a volatile element inventory comparable to that of Earth, or whether it has always been deficient in volatiles, and
- c. examination of several types of chemical processes of astrophysical interest, including condensation in cool stellar atmospheres and low-density nova shells, in which grains condense and cool in isolation, and grain-grain reaction products may be negligible, and also including the disequilibrium chemistry of such ejected interstellar matter under the influence of supernova shock waves in star-forming regions.

The original program of research was shortened by NASA to two years. In that time we have pursued all three of these projects

ORIGINAL PAGE IS  
OF POOR QUALITY

to a considerable degree, but of course have left a number of loose ends that will be picked up under the continuation of this research at the University of Arizona. Our achievements are as follows:

a. We developed detailed physico-chemical models of the Jovian subnebula (as described in last years report) which were completed at the same time that Lunine and Stevenson of Cal Tech submitted a very large paper on the same subject. Although the chemical aspects of the problem were more closely studied in our work, and the physical aspects treated more thoroughly in Lunine and Stevenson's paper, the overall results are quite closely comparable. The largest differences in the results arise from relatively minor details of the mass and mass distribution in the Jovian subnebula, and are not important. We then turned our attention to the planetary side of the problem, examining the evidence for departures from solar composition in the envelopes of Jupiter and Saturn. Such departures may be due to fractionation processes during the era of planetary accretion, or to some form of differentiation within the planets once they have formed. For Jupiter, the evidence favors minor enrichment of carbon (and possibly nitrogen) over solar composition, and possible severe depletion of oxygen below its solar abundance by a factor of 10 to 100. We have examined the molecular composition of a suite of Jovian model atmospheres of non-solar composition, including studying the chemical effects of lightning discharges on the atmospheric chemistry, using the Voyager lightning results as the standard on which the calculations were based. The major results

ORIGINAL PAGE IS  
OF POOR QUALITY

of this study are included in Section I of the present report. They will be submitted for publication in *Icarus* in the near future. We find that modest enhancements of C and N may serve to explain the observed presence of HCN in the Jovian troposphere, and that the production of organic matter by disequilibrium processes (shock wave heating) is only modestly larger than in a solar-composition atmosphere.

b. Our study of the geochemistry of Venus has culminated in a paper published in *Science* during the past year. A copy of that paper is included as Section II of the present report. Further planned work on the evolution of the hydrogen isotopic evolution of the Venus atmosphere, taking into account the known meteoritic and cometary mass fluxes through the inner solar system, has been deferred for future investigation at the University of Arizona. With the arrival of Shailendra Kumar at the U. of A. next year, and with the interest and involvement of Don Hunten, this project should continue to move forward.

c. We reported our initial work on the subject of condensation of preplanetary matter in low-density astrophysical settings in last year's report. Since then we have concentrated our efforts on the carbon chemistry of shocked interstellar cloud material, in an attempt to see whether conditions in star-forming regions are dependent upon photochemistry for the production of complex organic matter, or whether shock-wave heating from supernova explosions might be important in dense (optically opaque) clouds. The results, although still not complete, are very promising. Some of the features of this exploratory work are described in

Section III of the present report. Both aspects of this study (isolated-grain condensation and shock chemistry of the interstellar medium) will culminate in publications within the next few months.

After two of the originally planned three years, two of our three main projects have resulted in papers, one of which has already appeared in Science. We believe that work of this sort, done in this manner, is potentially a very cost-effective way for NASA or other agencies to get basic research done. It was our original purpose to start up this project for the purpose of providing a base of operations for young recent Ph. D.'s who are unable to find academic appointments under the present condition of the academic job market, and who would be lost to space science permanently if they had to seek employment elsewhere. We still believe that this principle is valid, but the premature termination of this productive project renders such a goal impossible.

ORIGINAL PAGE IS  
OF POOR QUALITY

I

Non-Solar Compositional Models of the  
Troposphere of Jupiter

Since 1969 it has been usual to assume that the elemental compositions of the tropospheres of Jupiter and Saturn are very close to that of the Sun. However, in recent years a number of interpretations of Earth-based and spacecraft-based infrared spectra of these planets have appeared which suggest significant departures from solar composition.

The most impressive such departure that has been suggested is for water vapor. Interpretations of IR spectra in the 5 $\mu$ m transmission window suggest that there is insufficient water to saturate the atmosphere at temperatures above about 220K (Hanel, et al., 1979). When the 5 $\mu$ m flux was qualitatively regarded as due to emission from small regions devoid of clouds, and therefore presumably dominated by very dry descending air, this observation presented no obvious conflict with the assumption of overall solar composition for the troposphere. However, recent work indicates that the 5 $\mu$ m emission may originate from a deep level within the troposphere over most or all of the planetary surface, and that this flux is attenuated by a thick, variable overlying atmosphere that must be dry everywhere on the planet down to levels where the temperature is 250 K or higher. The most careful available interpretations of the 5 $\mu$ m spectrum (Bjoraker, et al., 1981; Drossart and Encrenaz, 1982) suggest that water vapor must be depleted by a factor of 100 or more below the solar abundance ratio of H<sub>2</sub>O:H<sub>2</sub> = 1.3x10<sup>-3</sup> throughout the troposphere. This would have an important effect on both the atmospheric molecular composition and the cloud mass and structure. It also would greatly change the amount and composition of organic compounds

ORIGINAL PAGE IS  
OF POOR QUALITY

produced by disequilibrating processes such as lightning.

A second suggested departure from solar composition has been reviewed by Wallace and Hunten (198x), who summarize and criticize evidence from spectroscopic studies by Cochran (1977), Sato and Hansen (1979), and others. Recent work by Encrenaz and Combes (1982) and other post-Voyager results summarized therein suggest a 2- to 4-fold enrichment of methane in Jupiter and up to a 6-fold enrichment of methane in Saturn (Boriez and de Bergh, 1981).

A third, more subtle issue, is whether ammonia might be slightly enriched on Jupiter. It is likely that an enhancement of more than about a factor of two would make itself known through its effect on the microwave opacity of the troposphere, resulting in lower brightness temperatures in the centimeter region (Gulkis and Poynter, 1976). We shall consider ammonia abundances of 1 to 2 times the solar abundance.

Finally, phosphine appears to be enriched above solar proportions in Saturn by a factor of 1.5 or 2 (Tokunaga, et al., 1980; Larson, et al., 1980), but to have about 0.5 to 1 times its solar abundance in the upper troposphere of Jupiter (Drossart and Encrenaz, 1982). The latter authors conclude that the spectroscopic determinations of phosphine in the 5 $\mu$ m and 10 $\mu$ m regions are consistent with a solar phosphine abundance in the lower troposphere, modified by destruction of phosphine near the tropopause by solar UV photolysis (Prinn and Lewis, 1975).

We must recall that the observed amounts of phosphine on both Jupiter and Saturn near their tropopauses are not expected on the basis of strict equilibrium calculations on a solar-composition

atmosphere, since a solar water vapor abundance would cause complete oxidation of phosphine to condensable phosphates at all temperatures below about 600K (Barshay and Lewis, 1978; Lewis and Fegley, 1981). However, it is kinetically plausible that, in a rapidly convecting atmosphere, oxidation of phosphine would quench near 800K, leaving most of the phosphine unoxidized (Prinn and Lewis, 1975).

The only other chemically active element so far identified spectroscopically, germanium (as the hydride germane), appears to have slightly less than its solar abundance on Jupiter, in accord with theoretical expectations (Lewis and Fegley, 1981).

The present paper reports on the molecular composition of Jovian model atmospheres in which carbon is enriched by a factor of five, and nitrogen by a factor of two, relative to the Sun. The consequences of several different oxygen depletion factors, from 10 to  $10^5$  times below solar abundance, are explored.

We shall examine the equilibrium composition of such a non-solar-composition atmosphere along a plausible Jovian adiabat, the likely effects of quenching of equilibration in rising parcels of gas, and the effects of lightning discharges on the abundances of trace constituents of such an atmosphere. Our method of calculation is basically that used by Lewis (1969), Barshay and Lewis (1975), and Lewis and Fegley (1981), adapted to run in BASIC on a microcomputer system. The method of treatment of the chemical effects of lightning is that used by Lewis (1980), scaled to the level of lightning activity deduced by Borucki, et al. (1982).

ORIGINAL PAGE IS  
OF POOR QUALITY

### Equilibrium Composition

The mole fractions of a number of species at equilibrium along a Jovian adiabat are displayed in Figs. 1 and 2 for two different degrees of depletion of oxygen. All important oxygen compounds (water vapor, CO, formaldehyde, etc.) are depleted by the same factor that total oxygen is depleted: there are thus no important changes in the distribution of oxygen between its compounds, only in the overall abundance of this subset of compounds. Nitrogen compounds are raised in abundance by a factor of two for each N atom per molecule: thus ammonia is enhanced in abundance by a factor of two, and the relatively rare molecular nitrogen by a factor of four. Carbon compounds are enhanced by a factor of five for each C atom per molecule. Thus  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CN}$  would be enhanced by a factor of  $625 \times 2 = 1250$  relative to its abundance in a solar-composition gas. Since some models of the atmospheres of Uranus and Neptune assume C:H ratios up to 1000 times the solar ratio, the extension of such calculations to these planets can be expected to yield much higher equilibrium concentrations of complex species than any chemical models of Jupiter and Saturn have included to date.

The model atmosphere in Fig. 1 (with a  $10^5$ -fold depletion of oxygen) could not generate a distinct water cloud layer, and the ammonia clouds would be about twice as dense as in standard solar-composition models. Saturation of both solid ammonia and solid ammonium hydrosulfide would occur at a slightly higher temperature than in the solar-composition case. Aside from the

ORIGINAL PAGE IS  
OF POOR QUALITY

Caption to Fig. 1

Molecular abundances along a Jovian adiabat at chemical equilibrium, with oxygen depleted by a factor of  $10^5$ , carbon enriched by a factor of 5, and nitrogen enriched by a factor of 2. The results for many other compounds under these conditions and at a variety of other conditions are given in the Appendices.



trivial enrichments of methane and ammonia and depletion of water, the only departures from the composition of an equilibrium solar-composition atmosphere visible at temperatures below 300K would be an increased N<sub>2</sub> abundance. However, the molecular nitrogen abundance in the upper troposphere is surely not regulated by local equilibrium, but rather by quenching of nitrogen reduction in rising parcels of gas (Barshay and Lewis, 1978).

The atmosphere in Fig. 2, with only a 10-fold depletion of water, would indeed form a water ice cloud layer with significant optical thickness. However, it may also provide more than the observed amount of water vapor near the ~250K level. Aside from these effects due solely and directly to the water vapor abundance, the observable properties of these two model atmospheres would be very similar. They would, however, differ significantly under the effects of lightning discharges: oxygen-bearing products such as carbon monoxide and formaldehyde would be produced in significant amounts in the latter case, but not in the former.

Note that the presence of phosphine at  $T < 600\text{K}$  is fully compatible with complete chemical equilibration if insufficient water vapor is available to oxidize all the phosphine. This will be the case if the water vapor abundance is less than 1.5 times the phosphine abundance. This occurs if water is depleted by a factor of  $10^3$  or more. Thus any atmospheric composition which provides a guarantee of phosphine stability would be incapable of generating detectable traces of CO due to rapid overturn. In this case, an exogenic source for CO must be sought (see, for example,


ORIGINAL PAGE IS  
OF POOR QUALITY

Caption to Fig. 2

Molecular abundances along a Jovian adiabat at chemical equilibrium, with oxygen depleted by a factor of 10, carbon enriched by a factor of 5, and nitrogen enriched by a factor of 2. The dashed branches of the curves for phosphine and phosphorus sesquioxide assume exact equilibration, and are not likely to be attained during the convective overturn time. See the text for discussion.

15  
Fig. 2

ORIGINAL PAGE IS  
OF POOR QUALITY



Prather and McElroy, 1980; Strobel and Yung, 1980).

The observed presence of a trace of germane could easily be removed by reaction with hydrogen sulfide to precipitate GeS in any of these models, irrespective of the degree of water depletion. In all these models, as in the solar composition case (Barshay and Lewis, 1978; Lewis and Fegley, 1981), the observed presence of GeH<sub>4</sub> near the tropopause must be attributed to departures from equilibrium at low temperatures.

#### Quenching During Vertical Mixing

Kinetic arguments have been presented which suggest that quench temperatures near 1000K are reasonable for the reactions destroying many high-temperature species, such as CO, GeH<sub>4</sub>, PH<sub>3</sub>, HCN, acetylene, etc. (Prinn and Lewis, 1975; Prinn and Barshay, 1979; Lewis and Fegley, 1981; and the review by Lewis, 1983). For nitrogen, quench temperatures of 1000 to 1500K would result in a mole fraction of 4 to 12 ppm N<sub>2</sub>. Quench of C<sub>2</sub>H<sub>6</sub> destruction at 1000 to 1200K would provide 1.5 to 10 ppb of ethane at higher altitudes.

The only three observed species that can plausibly be attributed to quenching of hot lower tropospheric gases due to rapid vertical mixing in an adiabatic solar-composition atmosphere are CO, PH<sub>3</sub> and GeH<sub>4</sub>. We have already seen that a sufficiently low oxygen abundance renders CO negligible while making PH<sub>3</sub> unconditionally stable, while the status of germane is essentially unaffected by the oxygen abundance. But what other species may plausibly be formed in greater quantities in a low-oxygen atmosphere

ORIGINAL PAGE IS  
OF POOR QUALITY

than in a solar-composition gas? After N<sub>2</sub> and ethane, are any of these species potential indicators of the oxygen content of the lower troposphere?

Of the many other organic species that are present in trace amounts at higher temperatures along the Jovian adiabat, methyl amine, ethylene, methyl mercaptan and HCN are the only species that even remotely approach ethane in abundance. Near 1200K all of these species have about 1% of the abundance of ethane (~0.1 ppb vs. 10 ppb for ethane). These levels are far too low for spectroscopic detection of ethylene, methyl amine and methyl mercaptan, which are subject to overwhelming interference from methane, ethane and NH<sub>3</sub>. Further, photochemical production of ethylene and methyl amine may mask such a deep-mixing source, and photochemical destruction of methyl mercaptan, CH<sub>3</sub>SH, should be rapid. The spectroscopic detection limit for HCN is on the order of 1 ppb, and indeed its detection in the upper troposphere has been reported (Tokunaga, et al., 1980). Within the uncertainty of our knowledge of the elemental abundances, thermal structure, and kinetic behavior of the atmosphere, we conclude that it is conceivable that the observed HCN originates from quenching of deep tropospheric gases by rapid convective overturn.

#### Effects of Lightning Discharges

The importance of electrical discharge activity on Jupiter has long been a matter of debate. Bar-Nun (1975) advocated an enormous rate of dissipation of energy by Jovian lightning, equivalent to at least the entire internal heat flux of the

planet. Lewis (1976) argued that the efficiency of conversion of the total convective heat flux into lightning certainly could not reasonably be greater than unity. He advocated the use of the terrestrial conversion efficiency, about  $10^{-4}$ , in the absence of any direct evidence for lightning on Jupiter. Calculations of the chemical effects of this amount of lightning on the atmosphere near the water cloud layer showed that the most abundant lightning-produced species in a solar-composition gas, CO, could not be produced in sufficient quantity to provide the observed amount of CO. These calculations also showed that that only very small concentrations of CO ( $2 \times 10^{-14}$ ), N<sub>2</sub> ( $2 \times 10^{-15}$ ), HCN ( $3 \times 10^{-16}$ ) and C<sub>2</sub>H<sub>2</sub> ( $1 \times 10^{-16}$ ) would be found mixed into the bulk atmosphere (Lewis, 1980a). No other shock wave product would have a mole fraction higher than  $10^{-17}$ .

Bar-Nun has continued to advocate enormous conversion efficiencies, and has claimed that lightning must be localized in the topmost (ammonia) cloud layer where the yields of organic products from shock events are much higher than in a solar-composition gas (where O is more abundant than C, and CO forms in preference to organic matter). However, the Voyager detection of lightning flashes and whistlers permits the conclusion that high-altitude lightning is several orders of magnitude rarer than Bar-Nun claims (Lewis, 1980b). A careful assessment of the total body of evidence regarding Jovian lightning (Borucki, et al., 1982) has led to the conclusion that the conversion efficiency of atmospheric convective energy flux into lightning is between  $0.27 \times 10^{-4}$  and  $0.5 \times 10^{-4}$ , in excellent agreement with the

ORIGINAL PAGE IS  
OF POOR QUALITY

prejudices of Lewis (1976).

Because the relative importance of CO and organic compounds is a sensitive function of the C:O ratio, and because all of the models explored herein have C:O ratios greater than unity, it is clearly important to explore the consequences of lightning activity in these models. Note that, since C:O is everywhere greater than unity, there is no longer a fundamental distinction between the chemical consequences of high-altitude and low-altitude discharges.

ORIGINAL PAGE IS  
OF POOR QUALITY

We present first the mole fractions of a number of distinctive shock products in the shocked gas at a temperature of 2000K for the cases of:

1. O depleted 10x; peak shock pressure 50 b
2. O depleted 100x; peak shock pressure 50 b
3. O depleted 10x; peak shock pressure 10 b
4. O depleted 100x; peak shock pressure 10 b.

The graphite activity is indicated for each run: the abundances of organic compounds generally reach a maximum where the activity of graphite approaches unity. Note the great insensitivity of the results to the degree of depletion of O once O is significantly less abundant than C. The bottom line in Table I gives the percent of total carbon converted into organic matter. The two 10-bar examples feature 33 and 35% conversion: it is obvious that no process whatsoever can be more than three times as efficient as this, no matter what assumptions are made!

Also note that the graphite activity reaches 0.7 in the latter runs. The highest possible value for the graphite activity, and hence for the abundances of carbon compounds, occurs almost exactly at the 2000K, 10 bar point illustrated. Under no circumstances can the activity of graphite surpass 1.000.

Several organic compounds, notably  $C_2H_2$ , HCN and  $C_2H_4$ , attain mole fractions greater than  $10^{-5}$  in the shocked gas. The total mole fraction of all organic species approaches  $6.5 \times 10^{-4}$ . The total yield of organic compounds thus can be as high as about 30 times the yield of organics in a shocked solar composition gas, in which the favored high-temperature carbon compound is CO.

ORIGINAL PAGE IS  
OF POOR QUALITY

Table I

Mole Fractions of Gases Shocked to 2000K  
vs. (O depletion factor, shock pressure)

Species	(10x, 50b)	(100x, 50b)	(10x, 10b)	(100x, 10b)
CO	$1.15 \times 10^{-4}$	$1.16 \times 10^{-5}$	$1.19 \times 10^{-4}$	$1.19 \times 10^{-5}$
N <sub>2</sub>	$1.82 \times 10^{-4}$	$1.82 \times 10^{-4}$	$1.48 \times 10^{-4}$	$1.46 \times 10^{-4}$
HCN	$3.74 \times 10^{-5}$	$3.85 \times 10^{-5}$	$1.16 \times 10^{-4}$	$1.18 \times 10^{-4}$
C <sub>2</sub> H <sub>2</sub>	$3.73 \times 10^{-5}$	$3.98 \times 10^{-5}$	$4.41 \times 10^{-4}$	$4.69 \times 10^{-4}$
C <sub>2</sub> H <sub>4</sub>	$1.73 \times 10^{-5}$	$1.84 \times 10^{-5}$	$4.09 \times 10^{-5}$	$4.31 \times 10^{-5}$
C <sub>2</sub> H <sub>6</sub>	$4.75 \times 10^{-7}$	$5.07 \times 10^{-7}$	$2.25 \times 10^{-7}$	$2.37 \times 10^{-7}$
CH <sub>3</sub> CN	$2.17 \times 10^{-8}$	$2.31 \times 10^{-8}$	$4.62 \times 10^{-8}$	$4.85 \times 10^{-8}$
HCHO	$5.25 \times 10^{-9}$	$5.25 \times 10^{-10}$	$1.08 \times 10^{-9}$	$1.08 \times 10^{-10}$
CH <sub>3</sub> NH <sub>2</sub>	$3.22 \times 10^{-10}$	$3.22 \times 10^{-10}$	$4.00 \times 10^{-11}$	$4.09 \times 10^{-11}$
CH <sub>3</sub> SH	$1.67 \times 10^{-10}$	$1.72 \times 10^{-10}$	$1.11 \times 10^{-10}$	$1.14 \times 10^{-10}$
CH <sub>3</sub> OH	$2.47 \times 10^{-10}$	$2.47 \times 10^{-11}$	$1.01 \times 10^{-12}$	$1.01 \times 10^{-13}$
CO <sub>2</sub>	$8.72 \times 10^{-11}$	$8.45 \times 10^{-13}$	$5.32 \times 10^{-12}$	$5.19 \times 10^{-14}$
a (graphite)	0.202	0.208	0.692	0.710
% C converted	5	5	33	35

Obviously, when the oxygen abundance is insufficient to use up more than a small portion of the available carbon, other compounds must be formed from that carbon. Since the only available elements with abundances greater than or comparable to C are H and N, obviously the compounds formed must be hydrocarbons, nitriles, amines, etc. The hydrocarbon abundances are enhanced by about a factor of 100, and HCN is enriched by about a factor of 10, in the O-depleted gas. However, it is important to realize that the 2-fold enrichment of N and the 5-fold enrichment of C assumed in these calculations account for an enrichment of C<sub>2</sub> hydrocarbons by a factor of 25 and HCN by a factor of 10. Thus for the heavier hydrocarbons the effects of C enrichment and O depletion are comparable.

It is not sufficient simply to know how much of these products are present in the shocked gas. What we need to know is the over-all concentration of these species in the bulk atmosphere after quenching and mixing of the shocked gas with the unheated background gas. The method of calculation of Lewis (1980), modified to take into account the best available estimates of the lightning energy dissipation rate (Borucki, et al., 1982) will be used.

Borucki, et al., find that the rate of energy dissipation by Jovian lightning is about  $0.36 \text{ erg/cm}^2 \text{ s}$ . We shall assume that this entire energy flux is employed in heating parcels of the Jovian atmosphere to exactly the optimum pressure and temperature conditions for synthesis of organic matter. Too low a shock temperature produces too little of the organic products, while

shocking to too high a temperature wastes energy without increasing the yield of interesting products. The 10-bar points in Table I are very close to that optimum. The energetic cost of heating the parcel to the optimum temperature (2000 to 2500K) is about  $3 \times 10^{11}$  erg/g. Thus the available lightning energy flux can shock about  $1.2 \times 10^{-12}$  g/cm<sup>2</sup> s to these optimum conditions. The total mass flux through the region of the water clouds is given by the product of the density times the mean vertical overturn speed,  $v$ . But  $v$  is just the ratio of the eddy diffusion coefficient  $K$  to the scale height  $H$ . For  $K = 3 \times 10^8$ ,  $H = 5 \times 10^6$ , and a density of  $10^{-4}$  g/cm<sup>3</sup>, we calculate a turbulent mass flux of  $8 \times 10^{-3}$  g/cm<sup>2</sup> s through this region. The mass fraction of shocked gas in the total flow is then  $1.5 \times 10^{-10}$ . Thus a species produced with a mole fraction of  $10^{-4}$  in the shocked gas (for example, HCN) would have a mole fraction of  $\sim 10^{-14}$  in the bulk atmosphere.

Table II summarizes the mole fractions of lightning-produced gases after dilution by turbulent mixing with the unshocked background atmosphere. In several ways, this is a generous estimate. First, the largest plausible enrichment factors for C and N are used. Second, water is assumed to be depleted to much less than the C abundance. Third, the very low-pressure conditions used tend to maximize the yield of organic matter per erg, but realistic shock events near the water saturation level on Jupiter (200 to 280K for the range of water abundances from  $10^{-5}$  of solar to solar) will more typically run between 15 and 50 bars. At such high pressures the optimum conditions for formation of organic matter occur at much higher temperatures, and the

Table II

Mole Fractions of Lightning-Produced Gases  
after mixing with Unshocked Atmosphere

C <sub>2</sub> H <sub>2</sub>	$7.0 \times 10^{-14}$
N <sub>2</sub>	$2.7 \times 10^{-14}$
HCN	$1.8 \times 10^{-14}$
C <sub>2</sub> H <sub>4</sub>	$6.4 \times 10^{-15}$
C <sub>2</sub> H <sub>6</sub>	$3.7 \times 10^{-17}$
CH <sub>3</sub> CN	$7.3 \times 10^{-18}$
CH <sub>3</sub> SH	$1.7 \times 10^{-20}$

energetic cost of achieving these higher temperatures will decrease the net energetic efficiency of the process. This will of course result in lower mole fractions of the products in the bulk atmosphere after dilution.

One cannot simply postulate that all lightning occurs at the ammonia cloud layer and therefore gives vastly better yields: in fact, the Voyager optical flash data place very severe limits on such flashes, and the very best of them would be only about twice as efficient in generating organic matter as the examples used in Table II. The main advantage of postulating such lightning is that the production of organic matter may be assigned to a region where the eddy mixing rate is low (say,  $10^6 \text{ cm}^2/\text{s}$ ), and may thus result in greatly enhanced abundances of products. However, this is the logical equivalent of postulating extensive lightning at the tropopause on Earth. Such lightning is of course very rare, and is invariably associated with massive, highly turbulent nimbus towers that are so buoyant (due to latent heat release) that they penetrate and overshoot the tropopause. On Jupiter this kind of behavior would be ruled out on energetic grounds by a low water vapor abundance: if the requisite amount of water were present (perhaps 10 times the solar abundance), and if these motions remained coherent from the saturation level ( $\sim 300\text{K}$ ) to the tropopause ( $\sim 130\text{K}$ ) over a vertical travel of 80 km, then the lightning products would be those for a wet atmosphere, rich in  $\text{CO}$ .

The observational implications of this discussion are easy to express: lightning discharges on Jupiter do not produce any

products in observable amounts. So far as we can address the HCN abundance problem, the best available source appears to be quenching of hot gases at equilibrium near the 1200 to 1400K level along a Jovian adiabat.

ORIGINAL PAGE IS  
OF POOR QUALITY

## References

- A. Bar-Nun (1975). *Icarus* 24, 197-210.
- S. Barshay and J. S. Lewis (1978). *Icarus* 33, 593-611.
- G. Bjoraker, U. Fink, H. P. Larson and V. Kunde (1981). *Bull. Am. Astron. Soc.* 13, 735.
- W. J. Borucki, A. Bar-Nun, F. L. Scarf, A. F. Cook II and G. E. Hunt (1982). *Icarus* 52, 492-502.
- J. C. Buriez and C. de Bergh (1981). *Astron. Astrophys.* 94, 382-390.
- W. D. Cochran (1977). *Icarus* 31, 325-347.
- P. Drossart and T. Encrenaz (1982). *Icarus* 52, 483-491.
- T. Encrenaz and M. Combes (1982). *Icarus* 52, 54-61.
- S. Gulikis and R. Poynter (1976). *Phys. Earth Planet. Interiors* 6, 36-43.
- R. Hanel, M. Conrath, M. Flasar, V. Kunde, P. Lowman, W. Maguire, J. Pearl, J. Pirraglia, R. Samuelson, D. Gautier, P. Gierasch, S. Kumar and C. Ponnampertuma (1979). *Science* 204, 972-976.
- H. P. Larson, U. Fink, H. A. Smith and D. S. Davis (1980). *Astrophys. J.* 240, 327-337.
- J. S. Lewis (1969). *Icarus* 10, 393-409.
- J. S. Lewis (1976). in *The Exobiology of Jupiter*, C. Ponnampertuma, ed., Academic Press, N. Y. 13-25.
- J. S. Lewis (1980a). *Icarus* 43, 85-95.
- J. S. Lewis (1980b). *Science* 210, 1351-1353.
- J. S. Lewis and B. Fegley (1983). *Space Sci. Revs.*, in press.

- M. J. Prather, J. A. Logan and M. B. McElroy (1978). *Astrophys. J.* 223, 1072-1081.
- R. G. Prinn and S. S. Barshay (1977). *Science* 198, 1031-1034.
- R. G. Prinn and J. S. Lewis (1975). *Science* 190, 294-296.
- M. Sato and J. E. Hansen (1979). *J. Atmos. Sci.* 36, 1133-1167.
- D. F. Strobel and Y. L. Yung (1979). *Icarus* 37, 256-263.
- A. T. Tokunaga, R. F. Knacke and S. T. Ridgway (1980). *Icarus* 44, 93-101.

## APPENDICES

	Page
Appendix 1: Listing of PROGRAM JUPITER.....	A 1
Appendix 2: Molecular abundances at equilibrium along an in a solar-composition Jovian model atmosphere	A 4
Appendix 3: Molecular abundances as in Appendix 2, but with a ten-fold depletion of O and C and N enrichment	A 8
Appendix 4: Molecular abundances as in Appendix 3, but with oxygen depleted by a factor of 100.....	A12
Appendix 5: Molecular abundances as in Appendix 4, but with oxygen depleted by a factor of $10^5$ .....	A15
Appendix 6: Molecular abundances in a gas shocked in the vicinity of the water clouds: O depleted 10x...	A19
Appendix 7: Molecular abundances as in Appendix 6, but with oxygen depleted by a factor of 100.....	A21
Appendix 8: Composition of a gas shocked in the vicinity of the ammonia clouds, with O depleted 10x.....	A22
Appendix 9: Composition as in Appendix 8, but with oxygen depleted by a factor of 100.....	A24

```

1 REM * PROGRAM JUPITER
10 READ ABH, ABHE, ABO, ABC, ABN, ABSU, ABNE
20 FOR NT = 1 TO 10
22 INPUT "Temperature = ", TR
25 READ T, KH, KH2O, KCO, KCH4, KCO2
26 READ T, KOH, KO, KC, KH2S, KHS
27 READ T, KCN, KHCN, KNH3, KSO2, KS
28 READ T, KC2H6, KC2H4, KC2H2, KC3H8
29 READ T, KCH3OH, KHCHO, KHCOOH, KCH3SH
30 READ T, KCH3NH2, KCH3CN, KC2N2, KN2H4, KN
36 IF TR = T THEN 40 ELSE 25
40 INPUT "Use adiabatic (1) or insert pressures (2)"; I
41 IF I = 1 THEN 45 ELSE 50
45 PR = 4.2E-08*T^3.33333
46 GOTO 60
50 INPUT "Total pressure (bars) = ", PR
60 ABH = .001*PR*ABH
61 ABHE = .001*ABHE*PR
62 ABO = .001*ABO*PR
63 ABC = .001*ABC*PR
64 ABN = .001*ABN*PR
65 ABSU = .001*ABSU*PR
66 ABNE = .001*ABNE*PR
80 REM INITIAL GUESS OF ELEMENTAL ACTIVITIES
81 RH = -(KH/4)+SQR(KH^2+8*ABH)
82 RN = -(KN/4)+SQR(KN^2+8*ABN)
83 RO = (2*ABO)/(3*KH2O*RH^2)
84 PH2 = RH^2
85 GR = ABC/((KCO*RO)+(KCH4*PH2^2))
86 RS = ABSU/((KSO2*RO^2)+(KH2S*PH2))
99 INDEX = 0
200 FOR ITER = 1 TO 100
201 INDEX = INDEX+1
202 PH = KH*RH
203 PH2 = RH^2
204 PHE = ABHE
205 PNE = ABNE
210 PCO = KCO*GR*RO
211 PCH4 = KCH4*GR*PH2^2
212 PCO2 = KCO2*GR*RO^2
213 PC = KC*GR
214 PC2H6 = KC2H6*PH2^3*GR^2
215 PC2H4 = KC2H4*PH2^2*GR^2
216 PC2H2 = KC2H2*PH2*GR^2
217 PC3H8 = KC3H8*PH2^4*GR^3
220 POH = KOH*RH*RO
221 PH2O = KH2O*PH2*RO
222 PO = KO*RO
223 PO2 = RO^2
224 PCH3OH = KCH3OH*GR*RO*PH2^2
225 PHCHO = KHCHO*GR*RO*PH2
226 PHCOOH = KHCOOH*GR*PO2*PH2
230 PCN = KCN*GR*RN
231 PHCN = KHCN*GR*RH*RN
232 PNH3 = KNH3*RN*PH2*RH
233 PN2 = RN^2
234 PN = RN*KN
235 PC2N2 = KC2N2*PN2*GR^2
236 PCH3CN = KCH3CN*RN*RH*PH2*GR^2
237 PN2H4 = KN2H4*PN2*PH2^2
238 PCH3NH2 = KCH3NH2*GR*RN*RH*PH2^2
240 PH2S = KH2S*RS*PH2
241 PSO2 = KSO2*RS*PO2
242 PHS = KHS*RH*RS

```

ORIGINAL PAGE IS  
OF POOR QUALITY

```

243 PS = KS*RS
244 PS2 = RS^2
245 PCH3SH = KCH3SH*GR*RS*PH2^2
400 REM CALCULATE ELEMENTAL SUMS
401 SH = (PH+POH+PHS+PHCN)+2*(PH2+PH2O+PH2S+PC2H2+PHCHO+PHCOOH)+3*(PCH3CN+PNH3)+
4*(PC2H4+PCH3OH+PCH3SH+PN2H4+PCH4)+5*PCH3NH2+6*PC2H6
402 SC = PCO+PCO2+PCH4+PC+PHCN+PCN+PCH3OH+PHCHO+PHCOOH+PCH3NH2+PCH3SH+2*(PC2H2+P
C2H4+PC2H6+PCH3CN+PC2N2)
403 SO = PH2O+PCO+POH+PO+PCH3OH+PCH3OH+2*(PO2+PCO2+PSO2+PHCOOH)
404 SN = PN+PNH3+PCN+PHCN+PCH3NH2+PCH3CN+2*(PN2+PC2N2+PN2H4)
405 SS = PS+PHS+PSO2+PH2S+PCH3SH+2*PS2
419 P = PH2+PH+PH2O+PCO+PCO2+PCH4+PN2+PNH3+POH+PH2S+PHE+PO+PNE+PHS
420 EH = ABS(SH-ABH)/ABH
421 EO = ABS(SO-ABO)/ABO
422 EC = ABS(SC-ABC)/ABC
423 EN = ABS(SN-ABN)/ABN
424 ES = ABS(SS-ABSU)/ABSU
440 IF EH<.001 THEN IF EO<.001 THEN IF EC<.001 THEN IF EN<.001 THEN IF ES<.001 T
HEN GOTO 501 ELSE GOTO 449
449 REM COMPARE ELEMENTAL SUMS TO ABUNDANCES
450 IF INDEX = 1 THEN 452
451 GOTO 455
452 RH = RH*SQR(ABH/SH)
453 GOTO 500
455 IF INDEX = 2 THEN 457
456 GOTO 460
457 RO = RO*(ABO/SO)
458 GOTO 500
460 IF INDEX = 3 THEN 462
461 GOTO 465
462 GR = GR*(ABC/SC)
463 GOTO 500
465 IF INDEX = 4 THEN 467
466 GOTO 470
467 RN = RN*SQR(ABN/SN)
468 GOTO 500
470 IF INDEX = 5 THEN 472
471 GOTO 488
472 RS = RS*(ABSU/SS)
488 INDEX = 0
500 NEXT ITER
501 REM OUTPUT RESULTS OF ITERATION
502 PLAY "mlt22002c8d8e8f8g8a8b803c202b16a16g16f16e16d16c2"
505 PRINT "Temperature = ";T, "Pressure = ";P;"bars      ";ITER;"iterations"
506 PRINT
507 PRINT "                Pressures of gases in bars"
508 PRINT "H =";PH;" H2 =";PH2;" He =";PHE;" Ne =";PNE
509 PRINT "O =";PO;" O2 =";PO2;" OH =";POH;" H2O =";PH2O
510 PRINT "N =";PN;" N2 =";PN2;" NH3 =";PNH3;" CN =";PCN
511 PRINT "C =";PC;" CO =";PCO;" CO2 =";PCO2;" CH4 =";PCH4
512 PRINT "S =";PS;" S2 =";PS2;" HS =";PHS;" H2S =";PH2S
513 PRINT "C2H6 = ";PC2H6;" C2H4 = ";PC2H4;" C2H2 = ";PC2H2
514 PRINT "CH3OH = ";PCH3OH;" HCHO = ";PHCHO;" HCOOH = ";PHCOOH
515 PRINT "CH3NH2 = ";PCH3NH2;" CH3CN = ";PCH3CN;" C2N2 = ";PC2N2
516 PRINT "N2H4 = ";PN2H4;" HCN = ";PHCN;" CH3SH = ";PCH3SH;" SO2 = ";PSO2
519 PRINT
520 PRINT "INPUT ABUNDANCES OF ELEMENTS"
521 PRINT "H =";ABH;" O =";ABO;" C =";ABC;" N =";ABN;" S =";ABSU
522 PRINT "                Graphite activity ="; GR
523 PRINT "CALCULATED ABUNDANCES OF ELEMENTS"
524 PRINT "H =";SH;" O =";SO;" C =";SC;" N =";SN;" S =";SS
545 INPUT "Type 1 to continue, 0 to end:", GOON
546 IF GOON = 1 THEN 600
547 GOTO 3000
600 NEXT NT
1000 DATA 1756,122.,1.192.,.6516,0.2065,0.0221,0.2295

```

ORIGINAL PAGE IS OF POOR QUALITY

1201 DATA 2000, 1.622E-3, 3467.4, 2.944E7, 3.908E-4, 2.254E10  
 1202 DATA 2000, 0.5808, 6.637E-4, 3.342E-11, 0.614, 0.1033  
 1203 DATA 2000, 2.183E-7, 1.472E-2, 2.046E-5, 4.498E5, 2.891E-3  
 1204 DATA 2000, 7.063E-9, 1.121E-5, 1.054E-3, 1.033E-13  
 1205 DATA 2000, 3.289E-3, 30.52, 3.674E4, 5.408E-7  
 1206 DATA 2000, 6.683E-11, 9.736E-7, 1.647E-6, 5.768E-15, 9.5E-10  
 1211 DATA 1900, 7.907E-4, 7691, 4.275E7, 5.236E-4, 7.907 E10  
 1212 DATA 1900, 1.932, 2.958E-4, 3.467E-12, .817, 4.2E-2  
 1213 DATA 1900, 1.972E-07, 9.705E-3, 2.438E-5, 1.406E6, 1.442E-3  
 1214 DATA 1900, 9.496E-9, 9.669E-6, 5.164E-4, 1.484E-13  
 1215 DATA 1900, 6.349E-3, 44.51, 1.230E5, 7.412E-6  
 1216 DATA 1900, 7.366E-11, 7.470E-7, 6.166E-7, 4.328E-15, 9.496E-9  
 1221 DATA 1800, 3.565E-4, 1.862E4, 6.471E7, 1.342E-3, 3.192E11  
 1222 DATA 1800, 2.203, 1.005E-4, 2.805E-13, 1.122, 3.147E-2  
 1223 DATA 1800, 4.345E-8, 9.462E-3, 2.965E-5, 5E6, 6.653E-4  
 1224 DATA 1800, 1.318E-8, 8.204E-6, 2.333E-4, 2.223E-13  
 1225 DATA 1800, 1.318E-2, 67.70, 4.712E5, 1.052E-5  
 1226 DATA 1800, 8.204E-11, 5.559E-7, 2.070E-7, 3.148E-15, 3.656E-11  
 1251 DATA 1500, 1.754E-5, 5.309E5, 3.054E8, 2.630E-3, 6.324E13  
 1252 DATA 1500, 3.622, 4.037E-6, 1.949E-17, 3.758, 1.1E-2  
 1253 DATA 1500, 1.368E-10, 1.663E-3, 6.252E-3, 6.237E8, 3.556E-5  
 1254 DATA 1500, 4.592E-8, 4.387E-6, 1.141E-5, 1.031E-12  
 1255 DATA 1500, .2110, 3.332E2, 7.757E7, 3.984E-5  
 1256 DATA 1500, 1.231E-10, 1.817E-7, 3.265E-9, 9.354E-16, 6.067E-14  
 1281 DATA 1200, 1.963E-7, 7.925E7, 3.013E9, 1.581E-2, 1.750E17  
 1282 DATA 1200, 7.709, 2.489E-8, 1.114E-23, 23.12, 2.198E-3  
 1283 DATA 1200, 2.339E-14, 1.227E-4, 1.923E-4, 8.77E11, 4.446E-7  
 1284 DATA 1200, 2.992E-7, 1.721E-6, 1.234E-7, 1.029E-11  
 1285 DATA 1200, 13.59, 3.637E3, 1.638E11, 2.935E-4  
 1286 DATA 1200, 2.265E-10, 3.390E-8, 6.470E-13, 1.517E-16, 4.197E-18  
 1301 DATA 1000, 2.259E-9, 1.153E10, 2.877E10, 9.75E-2, 4.786E20  
 1302 DATA 1000, 16.51, 1.56E-10, 6.31E-30, 141.6, 4.457E-4  
 1303 DATA 1000, 3.9E-18, 9.036E-6, 5.848E-4, 1.245E15, 5.623E-9  
 1304 DATA 1000, 1.945E-6, 6.714E-7, 1.334E-10, 1.028E-10  
 1305 DATA 1000, 873., 3.972E4, 3.459E14, 2.163E-3  
 1306 DATA 1000, 4.169E-10, 6.324E-9, 1.282E-14, 2.46E-17, 2.965E-22  
 1321 DATA 800, 2.891E-12, 1.945E13, 8.204E11, 1.4, 6.761E25  
 1322 DATA 800, 52.481, 7.925E-14, 0, 2.099E3, 4.0467E-5  
 1323 DATA 800, 8.166E-24, 1.853E-7, 2.999E-3, 6.683E19, 8.185E-12  
 1324 DATA 800, 3.228E-5, 2.042E-7, 1.603E-12, 4.808E-9  
 1325 DATA 800, 5.794E5, 1.66E6, 3.767E19, 4.325E-2  
 1326 DATA 800, 1.361E-9, 6.012E-10, 1.081E-18, 1.82E-18, 1.803E-28  
 1341 DATA 600, 4.613E-17, 4.295E18, 2.08E14, 100.2, 2.541E34  
 1342 DATA 600, 367.3, 2.667E-19, 0, 4887., 2.004E-8  
 1343 DATA 600, 2.71E-33, 2.786E-10, 4.169E-2, 1.406E26, 4.335E-18  
 1344 DATA 600, 3.485E-3, 2.404E-8, 2.028E-7, 2.168E-6  
 1345 DATA 600, 2.449E10, 7.464E8, 8.670E27, .153  
 1346 DATA 600, 8.222E-9, 1.057E-11, 1.791E-24, 2.228E-20, 8.298E-39  
 3000 END

ORIGINAL PAGE IS  
 OF POOR QUALITY

Ok  
 RUN  
 Temperature = 2000  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 2000

ORIGINAL PAGE IS  
 OF POOR QUALITY

Pressure = 4231.117 bars 55 iterations

Pressures of gases in bars

$\checkmark$ H = 9.872717E-02  $\checkmark$ H2 = 3704.859  $\checkmark$ He = 516.4525  $\checkmark$ Ne = .9715232  
 $\checkmark$ O = 2.603932E-10  $\checkmark$ O2 = 1.539273E-13  $\checkmark$ OH = 1.38698E-05  $\checkmark$ H2O = 5.040033  
 $\checkmark$ N = 1.672887E-10  $\checkmark$ N2 = 3.100888E-02  $\checkmark$ NH3 = .8124676 CN = 1.972217E-11  
 $\checkmark$ C = 1.714606E-14  $\checkmark$ CO = 5.925889E-03  $\checkmark$ CO2 = 1.78003E-06  $\checkmark$ CH4 = 2.752047  
 $\checkmark$ S = 1.18569E-07  $\checkmark$ S2 = 1.682077E-09  $\checkmark$ HS = 2.578749E-04  $\checkmark$ H2S = 9.329599E-02  
 $\checkmark$ C2H6 = 9.454096E-05  $\checkmark$ C2H4 = 4.050091E-05  $\checkmark$ C2H2 = 1.027846E-06  
 $\checkmark$ CH3OH = 9.087049E-06  $\checkmark$ HCHO = 2.275998E-05  $\checkmark$ HCOOH = 1.07494E-08  
 $\checkmark$ CH3NH2 = 5.044307E-06  $\checkmark$ CH3CN = 1.017646E-08 C2N2 = 1.344298E-14  
 $\checkmark$ N2H4 = 2.455018E-09  $\checkmark$ HCN = 8.094584E-05  $\checkmark$ CH3SH = 1.561928E-07 SO2 =  
 2.839605E-12

INPUT ABUNDANCES OF ELEMENTS

H = 7433.529 O = 5.045995 C = 2.758364 N = .8741592 S = 9.355409E-02  
 Graphite activity = 5.130479E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 7433.529 O = 5.045994 C = 2.758365 N = .8745714 S = 9.355414E-02  
 Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREFN

Ok  
 RUN  
 Temperature = 1900  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 1900

Pressure = 3566.139 bars 55 iterations

Pressures of gases in bars

H = 4.418467E-02 H2 = 3122.624 He = 435.2879 Ne = .8188406  
 O = 5.234005E-11 O2 = 3.13092E-14 OH = 1.910306E-05 H2O = 4.249506  
 N = 1.529486E-09 N2 = 2.594236E-02 NH3 = .685201 CN = 1.444064E-11  
 C = 1.576262E-15 CO = 3.439115E-03 CO2 = 1.125533E-06 CH4 = 2.321206  
 S = 4.452742E-08 S2 = 9.535084E-10 HS = 7.247219E-05 H2S = 7.877784E-02  
 C2H6 = 5.97653E-05 C2H4 = 1.948814E-05 C2H2 = 3.333153E-07  
 CH3OH = 4.980299E-06 HCHO = 1.118119E-05 HCOOH = 5.467281E-09  
 CH3NH2 = 2.939073E-06 CH3CN = 4.339645E-09 C2N2 = 3.306453E-15  
 N2H4 = 1.094804E-09 HCN = 3.971319E-05 CH3SH = 1.014639E-06 SO2 =  
 1.359313E-12

INPUT ABUNDANCES OF ELEMENTS

H = 6265.291 O = 4.252977 C = 2.324865 N = .7367782 S = 7.885133E-02  
 Graphite activity = 4.546474E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 6265.291 O = 4.252976 C = 2.324865 N = .7371283 S = 7.885136E-02  
 Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Ok  
 RUN  
 Temperature = 1800  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 1800 Pressure = 2978.005 bars 55 iterations

ORIGINAL PAGE IS  
 OF POOR QUALITY

Pressures of gases in bars

H = 1.820473E-02 H2 = 2607.649 He = 363.5012 Ne = .6837993  
 O = 7.349113E-12 O2 = 5.34734E-15 OH = 8.226365E-06 H2O = 3.550567  
 N = 5.309413E-12 N2 = .0210902 NH3 = .573375 CN = 1.341747E-12  
 C = 5.964491E-17 CO = 1.006192E-03 CO2 = 3.629453E-07 CH4 = 1.9404  
 S = 1.496487E-08 S2 = 5.059538E-10 HS = 3.614738E-05 H2S = 6.581084E-02  
 C2H6 = 1.056681E-05 C2H4 = 2.522348E-06 C2H2 = 2.750711E-08  
 CH3OH = 1.393552E-06 HCHO = 2.745031E-06 HCOOH = 1.397117E-09  
 CH3NH2 = 8.796895E-07 CH3CN = 4.860623E-10 C2N2 = 1.973932E-16  
 N2H4 = 4.514542E-10 HCN = 1.492067E-05 CH3SH = 3.421448E-07 SO2 =  
 6.013996E-13

INPUT ABUNDANCES OF ELEMENTS

H = 5232.033 O = 3.551585 C = 1.941454 N = .6152704 S = 6.584734E-02  
 Graphite activity = 2.126379E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 5232.031 O = 3.551585 C = 1.941454 N = .6155712 S = 6.584735E-02  
 Type 1 to continue, 0 to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

Ok  
 RUN  
 Temperature = 1500  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 1500 Pressure = 1621.74 bars 70 iterations

Pressures of gases in bars

H = 6.609692E-04 H2 = 1420.049 He = 197.9564 Ne = .3723851  
 O = 1.035603E-14 O2 = 6.580658E-18 OH = 3.501344E-07 H2O = 1.933974  
 N = 6.079574E-17 N2 = 1.004149E-06 NH3 = .3352532 CN = 2.732402E-17  
 C = 3.884824E-21 CO = 1.561576E-04 CO2 = 8.295081E-08 CH4 = 1.057113  
 S = 2.38929E-10 S2 = 4.514545E-11 HS = 2.785168E-06 H2S = 3.585643E-02  
 C2H6 = 5.224327E-06 C2H4 = 3.514738E-07 C2H2 = 6.43736E-10  
 CH3OH = 2.17562E-07 HCHO = 2.41937E-07 HCOOH = 1.44486E-10  
 CH3NH2 = 1.868421E-09 CH3CN = 3.871041E-13 C2N2 = 1.302568E-22  
 N2H4 = 1.894096E-15 HCN = 1.251706E-08 CH3SH = 1.07595E-07 SO2 =  
 2.757732E-14

INPUT ABUNDANCES OF ELEMENTS

H = 2849.273 O = 1.934131 C = 1.057282 N = .3350654 S = 3.585931E-02  
 Graphite activity = 1.993239E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 2849.272 O = 1.934131 C = 1.057282 N = .3352552 S = 3.585932E-02  
 Type 1 to continue, 0 to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

OK  
 RUN  
 Temperature = 1200  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 1200

ORIGINAL PAGE IS  
 OF POOR QUALITY

Pressure = 770.8155 bars 60 iterations

Pressures of gases in bars

H = 5.09985E-06 H2 = 674.954 He = 94.08841 Ne = .1769942  
 O = 4.277621E-19 O2 = 2.953622E-22 OH = 3.442014E-09 H2O = .9192864  
 N = 1.930491E-19 N2 = 2.115718E-03 NH3 = .1551027 CN = 7.506413E-20  
 C = 7.772461E-28 CO = 3.61285E-06 CO2 = 3.606337E-09 CH4 = .5025201  
 S = 4.855953E-13 S2 = 1.192916E-12 HS = 6.236913E-08 H2S = 1.704382E-02  
 C2H6 = 4.47848E-07 C2H4 = 3.816592E-09 C2H2 = 4.054488E-13  
 CH3OH = 7.423667E-09 HCHO = 2.943528E-09 HCOOH = 2.278328E-12  
 CH3NH2 = 8.603116E-09 CH3CN = 1.331026E-13 C2N2 = 6.663599E-24  
 N2H4 = 1.462149E-13 HCN = 1.023019E-08 CH3SH = 1.018907E-08 SO2 =  
 2.829173E-16

INPUT ABUNDANCES OF ELEMENTS

H = 1354.256 O = .9192901 C = .5025246 N = .1592562 S = 1.704388E-02  
 Graphite activity = 6.977075E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 1354.256 O = .9192901 C = .5025246 N = .1593342 S = 1.704389E-02

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

OK  
 RUN  
 Temperature = 1000  
 Use adiabat (1) or insert pressures (2)? 1  
 Temperature = 1000

Pressure = 419.7687 bars 60 iterations

Pressures of gases in bars

H = 4.330945E-08 H2 = 367.5638 He = 51.23862 Ne = .0963874  
 O = 1.842793E-23 O2 = 1.395416E-26 OH = 3.739084E-11 H2O = .5006265  
 N = 6.181932E-24 N2 = 4.347094E-04 NH3 = 8.592231E-02 CN = 1.689322E-24  
 C = 1.310925E-34 CO = 7.060576E-08 CO2 = 1.387474E-10 CH4 = .2736645  
 S = 1.002772E-15 S2 = 3.180299E-14 HS = 1.523853E-09 H2S = 9.281751E-03  
 C2H6 = 4.168833E-08 C2H4 = 3.915106E-11 C2H2 = 2.11634E-17  
 CH3OH = 2.894542E-10 HCHO = 3.582961E-11 HCOOH = 3.68583E-14  
 CH3NH2 = 4.677476E-10 CH3CN = 4.010396E-16 C2N2 = 2.405381E-27  
 N2H4 = 1.444773E-15 HCN = 7.50396E-11 CH3SH = 1.08269E-09 SO2 =  
 3.098182E-18

INPUT ABUNDANCES OF ELEMENTS

H = 737.5001 O = .5006266 C = .2736646 N = 8.672766E-02 S = 9.28175E-03  
 Graphite activity = 2.077535E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 737.4998 O = .5006266 C = .2736646 N = 8.679173E-02 S = 9.281754E-03

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Ok  
RUN

ORIGINAL PAGE IS  
OF POOR QUALITY

Temperature = 800

Use adiabat (1) or insert pressures (2)? 1

Temperature = 800 Pressure = 199.5163 bars 65 iterations

Pressures of gases in bars

H = 3.821188E-11 H2 = 174.7031 He = 24.35378 Ne = 4.581306E-02  
 O = 5.549603E-30 O2 = 4.903715E-33 OH = 4.857525E-14 H2O = .2379484  
 N = 1.071984E-30 N2 = 3.534962E-05 NH3 = 4.117369E-02 CN = 1.477951E-31  
 C = 0 CO = 1.748826E-10 CO2 = 1.00924E-12 CH4 = .1300732  
 S = 9.847014E-20 S2 = 1.447344E-16 HS = 6.434833E-12 H2S = 4.411629E-03  
 C2H6 = 1.594969E-09 C2H4 = 5.775289E-14 C2H2 = 2.595081E-21  
 CH3OH = 3.769647E-12 HCHO = 6.182015E-14 HCOOH = 9.823807E-17  
 CH3NH2 = 9.937118E-12 CH3CN = 7.648546E-20 C2N2 = 3.54101E-34  
 N2H4 = 1.96362E-18 HCN = 4.432783E-14 CH3SH = 4.834278E-11 SO2 =  
 3.942599E-21

INPUT ABUNDANCES OF ELEMENTS

H = 350.5348 O = .2379484 C = .1300732 N = 4.122177E-02 S = 4.411628E-03  
 Graphite activity = 3.044096E-06

CALCULATED ABUNDANCES OF ELEMENTS

H = 350.5348 O = .2379484 C = .1300732 N = 4.124439E-02 S = 4.411629E-03

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

RUN

Temperature = 600

Use adiabat (1) or insert pressures (2)? 1

Temperature = 600 Pressure = 76.47441 bars 65 iterations

Pressures of gases in bars

H = 3.774874E-16 H2 = 66.9635 He = 9.334786 Ne = 1.756011E-02  
 O = 0 O2 = 0 OH = 9.531462E-19 H2O = 9.120545E-02  
 N = 0 N2 = 4.791225E-07 NH3 = 1.581293E-02 CN = 0  
 C = 0 CO = 7.319215E-15 CO2 = 0 CH4 = 4.985694E-02  
 S = 2.239988E-26 S2 = 2.670011E-17 HS = 8.473705E-16 H2S = 1.690974E-03  
 C2H6 = 1.288488E-11 C2H4 = 1.327315E-18 C2H2 = 1.672128E-19  
 CH3OH = 3.864259E-15 HCHO = 1.758778E-18 HCOOH = 0  
 CH3NH2 = 2.317276E-14 CH3CN = 4.936496E-26 C2N2 = 0  
 N2H4 = 4.786719E-23 HCN = 1.751077E-19 CH3SH = 3.933742E-13 SO2 = 0

INPUT ABUNDANCES OF ELEMENTS

H = 134.3597 O = 9.120545E-02 C = 4.985694E-02 N = 1.580027E-02 S =  
 1.690973E-03

Graphite activity = 1.109639E-07

CALCULATED ABUNDANCES OF ELEMENTS

H = 134.3597 O = 9.120545E-02 C = 4.985694E-02 N = 1.581389E-02 S =  
 1.690974E-03

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N  
 Temperature = 2000 Pressure = 4219.798 bars 50 iterations

## Pressures of gases in bars

H = 9.847902E-02 H2 = 3686.258 He = 516.4525 Ne = .9715232  
 O = 2.604543E-11 O2 = 1.539996E-15 OH = 1.383819E-06 H2O = .5015906  
 N = 3.166214E-10 N2 = .1110793 NH3 = 1.526163 CN = 1.888116E-10  
 C = 8.672909E-14 CO = 2.998167E-03 CO2 = 9.008069E-08 CH4 = 13.78111  
 S = 1.191657E-07 S2 = 1.69905E-09 HS = 2.585213E-04 H2S = 9.329476E-02  
 C2H6 = 2.382665E-03 C2H4 = 1.025873E-03 C2H2 = 2.616634E-05  
 CH3OH = 4.551488E-06 HCHO = 1.145746E-05 HCOOH = 5.41256E-10  
 CH3NH2 = 4.768817E-05 CH3CN = 4.89094E-07 C2N2 = 1.232093E-12  
 N2H4 = 8.706225E-09 HCN = 7.72993E-04 CH3SH = 7.860851E-07 SO2 =  
 2.855236E-14

## INPUT ABUNDANCES OF ELEMENTS

H = 7433.529 O = .5045995 C = 13.79182 N = 1.748319 S = 9.355409E-02  
 Graphite activity = 2.595125E-03

## CALCULATED ABUNDANCES OF ELEMENTS

H = 7433.527 O = .5045994 C = 13.79182 N = 1.749142 S = 9.355418E-02  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

O depleted by 10x; 5x solar C; 2x solar N  
 Temperature = 1800 Pressure = 2970.04 bars 50 iterations

## Pressures of gases in bars

H = 1.815896E-02 H2 = 2594.554 He = 363.5012 Ne = .6837993  
 O = 7.37766E-13 O2 = 5.388962E-17 OH = 8.237557E-07 H2O = .3546459  
 N = 1.006977E-11 N2 = 7.586244E-02 NH3 = 1.079275 CN = 1.285769E-11  
 C = 3.01365E-16 CO = 5.103685E-04 CO2 = 1.84811E-08 CH4 = 9.705944  
 S = 1.504007E-08 S2 = 5.110518E-10 HS = 3.62377E-05 H2S = 6.580941E-02  
 C2H6 = 2.657194E-04 C2H4 = 6.374863E-05 C2H2 = 6.987104E-07  
 CH3OH = 6.997667E-07 HCHO = 1.385364E-06 HCOOH = 7.078366E-11  
 CH3NH2 = 8.324452E-06 CH3CN = 2.335737E-08 C2N2 = 1.812662E-14  
 N2H4 = 1.607633E-09 HCN = 1.426224E-04 CH3SH = 1.72002E-06 SO2 =  
 6.091266E-15

## INPUT ABUNDANCES OF ELEMENTS

H = 5232.033 O = .3551585 C = 9.707269 N = 1.230541 S = 6.584734E-02  
 Graphite activity = 1.074385E-03

## CALCULATED ABUNDANCES OF ELEMENTS

H = 5232.031 O = .3551585 C = 9.707269 N = 1.231151 S = 6.584739E-02  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

ORIGINAL PAGE IS  
OF POOR QUALITY

O depleted by 10x; 5x solar C; 2x solar N  
Temperature = 1500 Pressure = 1617.344 bars 70 iterations

Pressures of gases in bars

H = 6.592866E-04 H2 = 1412.828 He = 197.9564 Ne = .3723851  
O = 1.040552E-15 O2 = 6.643705E-20 OH = 3.509121E-08 H2O = .1933336  
N = 1.22525E-16 N2 = 4.078506E-06 NH3 = .6705075 CN = 2.781847E-16  
C = 1.962495E-20 CO = 7.926305E-05 CO2 = 4.230569E-09 CH4 = 5.286049  
S = 2.401472E-10 S2 = 4.5607E-11 HS = 2.792243E-06 H2S = 3.585601E-02  
C2H6 = 1.312996E-04 C2H4 = 8.8785E-06 C2H2 = 1.634437E-08  
CH3OH = 1.093108E-07 HCHO = 1.221789E-07 HCOOH = 7.331453E-12  
CH3NH2 = 1.878143E-08 CH3CN = 1.975754E-11 C2N2 = 1.350136E-20  
N2H4 = 7.615127E-15 HCN = 1.271113E-07 CH3SH = 5.407673E-07 SO2 =  
2.798348E-16

INPUT ABUNDANCES OF ELEMENTS

H = 2849.273 O = .1934131 C = 5.286409 N = .6701309 S = 3.585931E-02  
Graphite activity = 1.006924E-03

CALCULATED ABUNDANCES OF ELEMENTS

H = 2849.272 O = .1934131 C = 5.286409 N = .6705157 S = 3.585934E-02  
Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N  
Temperature = 1200 Pressure = 768.7373 bars 55 iterations

Pressures of gases in bars

H = 5.086935E-06 H2 = 671.5398 He = 94.08841 Ne = .1769942  
O = 4.2993E-20 O2 = 2.983635E-24 OH = 3.450697E-10 H2O = 9.192717E-02  
N = 3.792444E-19 N2 = 8.16508E-03 NH3 = .3023898 CN = 7.448318E-19  
C = 3.925844E-27 CO = 1.834086E-06 CO2 = 1.840058E-10 CH4 = 2.512599  
S = 4.88063E-13 S2 = 1.205072E-12 HS = 6.252734E-08 H2S = 1.704378E-02  
C2H6 = 1.125311E-05 C2H4 = 9.638732E-08 C2H2 = 1.029159E-11  
CH3OH = 3.730642E-09 HCHO = 1.486742E-09 HCOOH = 1.156589E-13  
CH3NH2 = 8.428991E-08 CH3CN = 6.620383E-12 C2N2 = 6.560852E-22  
N2H4 = 5.585853E-13 HCN = 1.012531E-07 CH3SH = 5.120423E-08 SO2 =  
2.872445E-18

INPUT ABUNDANCES OF ELEMENTS

H = 1354.256 O = .091929 C = 2.512623 N = .3185124 S = 1.704388E-02  
Graphite activity = 3.524097E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 1354.255 O = .091929 C = 2.512623 N = .3187201 S = .0170439  
Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O\*depleted by 10x; 5x solar C; 2x solar N  
 Temperature = 1000 Pressure = 418.6334 bars 60 iterations

Pressures of gases in bars

H = 4.319944E-08 H2 = 365.6989 He = 51.23862 Ne = .0963874  
 O = 1.852189E-24 O2 = 1.409682E-28 OH = 3.748603E-12 H2O = 5.006262E-02  
 N = 1.233176E-23 N2 = 1.72982E-03 NH3 = .1700957 CN = 1.702163E-23  
 C = 6.621639E-34 CO = 3.584567E-08 CO2 = 7.079948E-12 CH4 = 1.368321  
 S = 1.007885E-15 S2 = 3.212818E-14 HS = 1.527733E-09 H2S = 9.281751E-03  
 C2H6 = 1.047521E-06 C2H4 = 9.887828E-10 C2H2 = 5.372195E-16  
 CH3OH = 1.454649E-10 HCHO = 1.809796E-11 HCOOH = 1.871249E-15  
 CH3NH2 = 4.653476E-09 CH3CN = 2.025583E-14 C2N2 = 2.442087E-25  
 N2H4 = 5.690929E-15 HCN = 7.541793E-10 CH3SH = 5.441053E-09 SO2 =  
 3.145818E-20

INPUT ABUNDANCES OF ELEMENTS

H = 737.5001 O = 5.006265E-02 C = 1.368323 N = .1734553 S = 9.28175E-03  
 Graphite activity = 1.049388E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 737.5 O = 5.006266E-02 C = 1.368323 N = .1735553 S = 9.281758E-03  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 800 Pressure = 198.9755 bars 60 iterations

Pressures of gases in bars

H = 3.811463E-11 H2 = 173.815 He = 24.35378 Ne = 4.581306E-02  
 O = 5.577958E-31 O2 = 4.953954E-35 OH = 4.869919E-15 H2O = 2.379484E-02  
 N = 2.157579E-30 N2 = 1.431997E-04 NH3 = 8.223911E-02 CN = 1.502573E-30  
 C = 0 CO = 8.878852E-11 CO2 = 5.150129E-14 CH4 = .6503657  
 S = 9.897338E-20 S2 = 1.462175E-16 HS = 6.451257E-12 H2S = 4.411633E-03  
 C2H6 = 4.007795E-08 C2H4 = 1.458614E-12 C2H2 = 6.587657E-20  
 CH3OH = 1.894454E-12 HCHO = 3.122673E-14 HCOOH = 4.987581E-18  
 CH3NH2 = 9.97476E-11 CH3CN = 3.897904E-18 C2N2 = 3.659975E-32  
 N2H4 = 7.873866E-18 HCN = 4.495161E-13 CH3SH = 2.429492E-10 SO2 =  
 4.003347E-23

INPUT ABUNDANCES OF ELEMENTS

H = 350.5348 O = 2.379484E-02 C = .6503658 N = 8.244354E-02 S =  
 4.411628E-03  
 Graphite activity = 1.537642E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 350.5346 O = 2.379484E-02 C = .6503658 N = 8.252551E-02 S =  
 4.411634E-03  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 10x; 5x solar C; 2x solar N  
 Temperature = 600 Pressure = 76.26707 bars 65 iterations

Pressures of gases in bars

H = 3.765265E-16 H2 = 66.623 He = 9.334786 Ne = 1.756011E-02  
 O = 0 O2 = 0 OH = 9.555786E-20 H2O = 9.120545E-03  
 N = 0 N2 = 1.945733E-06 NH3 = 3.162349E-02 CN = 0  
 C = 0 CO = 3.716005E-15 CO2 = 0 CH4 = .2492847  
 S = 2.251438E-26 S2 = 2.697377E-17 HS = 8.495338E-16 H2S = 1.690976E-03  
 C2H6 = 3.237684E-10 C2H4 = 3.352293E-17 C2H2 = 4.244742E-18  
 CH3OH = 1.942004E-15 HCHO = 8.884006E-19 HCOOH = 0  
 CH3NH2 = 2.328944E-13 CH3CN = 2.518906E-24 C2N2 = 0  
 N2H4 = 1.924185E-22 HCN = 1.777929E-18 CH3SH = 1.976925E-12 SO2 = 0

INPUT ABUNDANCES OF ELEMENTS

H = 134.3597 O = 9.120544E-03 C = .2492847 N = 3.160054E-02 S =  
 1.690973E-03

Graphite activity = 5.605051E-07

CALCULATED ABUNDANCES OF ELEMENTS

H = 134.3596 O = 9.120545E-03 C = .2492847 N = 3.162738E-02 S =  
 1.690976E-03

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 100x; 5x solar C; 2x solar N  
 Temperature = 2000 Pressure = 4219.793 bars 50 iterations

## Pressures of gases in bars

H = 9.848499E-02 H2 = 3686.705 He = 516.4525 Ne = .9715232  
 O = 2.604231E-12 O2 = 1.539627E-17 OH = 1.383737E-07 H2O = 5.015913E-02  
 N = 3.165769E-10 N2 = .1110481 NH3 = 1.526226 CN = 1.887765E-10  
 C = 8.672514E-14 CO = 2.997672E-04 CO2 = 9.005502E-10 CH4 = 13.78383  
 S = 1.191513E-07 S2 = 1.69864E-09 HS = 2.585057E-04 H2S = 9.329479E-02  
 C2H6 = 2.383315E-03 C2H4 = 1.026029E-03 C2H2 = 2.616713E-05  
 CH3OH = 4.551839E-07 HCHO = 1.145695E-06 HCOOH = 5.411673E-12  
 CH3NH2 = 4.769375E-05 CH3CN = 4.890698E-07 C2N2 = 1.231635E-12  
 N2H4 = 8.70589E-09 HCN = 7.728962E-04 CH3SH = 7.861448E-07 SO2 =  
 2.854208E-16

## INPUT ABUNDANCES OF ELEMENTS

H = 7433.529 O = 5.045995E-02 C = 13.79182 N = 1.748319 S = 9.355409E-02  
 Graphite activity = 2.595007E-03

## CALCULATED ABUNDANCES OF ELEMENTS

H = 7433.528 O = 5.045995E-02 C = 13.79182 N = 1.749143 S = .0935542  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N  
 Temperature = 1800 Pressure = 2970.039 bars 50 iterations

## Pressures of gases in bars

H = 1.816007E-02 H2 = 2594.873 He = 363.5012 Ne = .6837993  
 O = 7.376757E-14 O2 = 5.387644E-19 OH = 8.237055E-08 H2O = 3.546461E-02  
 N = 1.006833E-11 N2 = 7.584067E-02 NH3 = 1.079319 CN = 1.28533E-11  
 C = 3.013053E-16 CO = 5.10205E-05 CO2 = 1.847292E-10 CH4 = 9.706406  
 S = 1.503823E-08 S2 = 5.109267E-10 HS = 3.623548E-05 H2S = 6.580944E-02  
 C2H6 = 2.657121E-04 C2H4 = 6.373903E-05 C2H2 = 6.985194E-07  
 CH3OH = 6.997143E-08 HCHO = 1.38509E-07 HCOOH = 7.076101E-13  
 CH3NH2 = 8.324165E-06 CH3CN = 2.334907E-08 C2N2 = 1.811424E-14  
 N2H4 = 1.607567E-09 HCN = 1.425824E-04 CH3SH = 1.719891E-06 SO2 =  
 6.08903E-17

## INPUT ABUNDANCES OF ELEMENTS

H = 5232.033 O = 3.551585E-02 C = 9.707269 N = 1.230541 S = 6.584734E-02  
 Graphite activity = 1.074172E-03

## CALCULATED ABUNDANCES OF ELEMENTS

H = 5232.031 O = 3.551585E-02 C = 9.707269 N = 1.231151 S = 6.584741E-02  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N  
 Temperature = 1500 Pressure = 1617.343 bars 70 iterations

## Pressures of gases in bars

H = 6.593272E-04 H2 = 1413.002 He = 197.9564 Ne = .3723851  
 O = 1.040424E-16 O2 = 6.642071E-22 OH = 3.508905E-09 H2O = 1.933336E-02  
 N = 1.225023E-16 N2 = 4.076999E-06 NH3 = .6705073 CN = 2.780686E-16  
 C = 1.962038E-20 CO = 7.923487E-06 CO2 = 4.228545E-11 CH4 = 5.286119  
 S = 2.401176E-10 S2 = 4.559576E-11 HS = 2.792071E-06 H2S = .035856  
 C2H6 = 1.312869E-04 C2H4 = 8.876554E-06 C2H2 = 1.633878E-08  
 CH3OH = 1.092988E-08 HCHO = 1.221505E-08 HCOOH = 7.328848E-14

CH3NH2 = 1.877937E-8 CH3CN = 1.974834E-11 C2N2 = 1.34901E-20  
N2H4 = 7.614188E-15 HCN = 1.270661E-07 CH3SH = 5.40708E-07 SO2 =  
2.797316E-18

INPUT ABUNDANCES OF ELEMENTS

H = 2849.273 O = 1.934131E-02 C = 5.286409 N = .6701309 S = 3.585931E-02  
Graphite activity = 1.00669E-03

CALCULATED ABUNDANCES OF ELEMENTS

H = 2849.272 O = 1.934131E-02 C = 5.286408 N = .6705155 S = 3.585934E-02  
Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N  
Temperature = 1200 Pressure = 768.7372 bars 55 iterations

Pressures of gases in bars

H = 5.087248E-06 H2 = 671.6225 He = 94.08841 Ne = .1769942  
O = 4.298771E-21 O2 = 2.9829E-26 OH = 3.450484E-11 H2O = 9.192716E-03  
N = 3.791813E-19 N2 = 8.162363E-03 NH3 = .3023953 CN = 7.445251E-19  
C = 3.924881E-27 CO = 1.833411E-07 CO2 = 1.839154E-12 CH4 = 2.512601  
S = 4.880029E-13 S2 = 1.204775E-12 HS = 6.25235E-08 H2S = 1.704378E-02  
C2H6 = 1.125174E-05 C2H4 = 9.636375E-08 C2H2 = 1.028781E-11  
CH3OH = 3.730186E-10 HCHO = 1.486377E-10 HCOOH = 1.156163E-15  
CH3NH2 = 8.428112E-08 CH3CN = 6.617255E-12 C2N2 = 6.555451E-22  
N2H4 = 5.585369E-13 HCN = 1.012177E-07 CH3SH = 5.119796E-08 SO2 =  
2.871384E-20

INPUT ABUNDANCES OF ELEMENTS

H = 1354.256 O = .0091929 C = 2.512623 N = .3185124 S = 1.704388E-02  
Graphite activity = 3.523232E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 1354.255 O = .0091929 C = 2.512624 N = .3187202 S = .0170439  
Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N  
Temperature = 1000 Pressure = 418.6333 bars 60 iterations

Pressures of gases in bars

H = 4.320209E-08 H2 = 365.7439 He = 51.23862 Ne = .0963874  
O = 1.851961E-25 O2 = 1.409335E-30 OH = 3.748373E-13 H2O = 5.006262E-03  
N = 1.232957E-23 N2 = 1.729206E-03 NH3 = .1700969 CN = 1.701442E-23  
C = 6.62001E-34 CO = 3.583243E-09 CO2 = 7.076463E-14 CH4 = 1.368321  
S = 1.007761E-15 S2 = 3.212027E-14 HS = 1.527639E-09 H2S = 9.281752E-03  
C2H6 = 1.047392E-06 C2H4 = 9.885394E-10 C2H2 = 5.370211E-16  
CH3OH = 1.45447E-11 HCHO = 1.809351E-12 HCOOH = 1.870558E-17  
CH3NH2 = 4.652938E-09 CH3CN = 2.024601E-14 C2N2 = 2.440019E-25  
N2H4 = 5.690312E-15 HCN = 7.539063E-10 CH3SH = 5.440383E-09 SO2 =  
3.144657E-22

INPUT ABUNDANCES OF ELEMENTS

H = 737.5001 O = 5.006266E-03 C = 1.368323 N = .1734553 S = 9.28175E-03  
Graphite activity = 1.04913E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 737.5 O = 5.006266E-03 C = 1.368323 N = .1735553 S = 9.281759E-03  
Type 1 to continue, 0 to end:

Ok

ORIGINAL PAGE IS  
OF POOR QUALITY

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Use adiabat (1) or insert pressures (2)? 1  
 O depleted by 100x; 5x solar C; 2x solar N  
 Temperature = 800 Pressure = 198.9755 bars 60 iterations

Pressures of gases in bars

H = 3.811698E-11 H2 = 173.8364 He = 24.35378 Ne = 4.581306E-02  
 O = 5.577271E-32 O2 = 4.952734E-37 OH = 4.869619E-16 H2O = 2.379484E-03  
 N = 2.157185E-30 N2 = 1.431473E-04 NH3 = 8.223923E-02 CN = 1.501928E-30  
 C = 0 CO = 8.875572E-12 CO2 = 5.147592E-16 CH4 = .6503657  
 S = 9.896119E-20 S2 = 1.461816E-16 HS = 6.45086E-12 H2S = 4.411634E-03  
 C2H6 = 4.007302E-08 C2H4 = 1.458255E-12 C2H2 = 6.585225E-20  
 CH3OH = 1.894221E-13 HCHO = 3.121904E-15 HCOOH = 4.985739E-20  
 CH3NH2 = 9.973546E-11 CH3CN = 3.895991E-18 C2N2 = 3.656833E-32  
 N2H4 = 7.872921E-18 HCN = 4.493508E-13 CH3SH = 2.429193E-10 SO2 =  
 4.001868E-25

INPUT ABUNDANCES OF ELEMENTS

H = 350.5348 O = 2.379484E-03 C = .6503658 N = 8.244355E-02 S =  
 4.411628E-03

Graphite activity = 1.537263E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 350.5346 O = 2.379484E-03 C = .6503658 N = 8.252553E-02 S =  
 4.411634E-03

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N  
 Temperature = 600 Pressure = 75.26708 bars 65 iterations

Pressures of gases in bars

H = 3.765497E-16 H2 = 66.63121 He = 9.334786 Ne = 1.756011E-02  
 O = 0 O2 = 0 OH = 9.555198E-21 H2O = 9.120544E-04  
 N = 0 N2 = 1.945015E-06 NH3 = .0316235 CN = 0  
 C = 0 CO = 3.714632E-16 CO2 = 0 CH4 = .2492847  
 S = 2.251161E-26 S2 = 2.696713E-17 HS = 8.494815E-16 H2S = 1.690976E-03  
 C2H6 = 3.237286E-10 C2H4 = 3.351467E-17 C2H2 = 4.243173E-18  
 CH3OH = 1.941765E-16 HCHO = 8.891815E-20 HCOOH = 0  
 CH3NH2 = 2.328658E-13 CH3CN = 2.517666E-24 C2N2 = 0  
 N2H4 = 1.923949E-22 HCN = 1.777272E-18 CH3SH = 1.976682E-12 SO2 = 0

INPUT ABUNDANCES OF ELEMENTS

H = 134.3597 O = 9.120544E-04 C = .2492847 N = 3.160055E-02 S =  
 1.690973E-03

Graphite activity = 5.60367E-07

CALCULATED ABUNDANCES OF ELEMENTS

H = 134.3596 O = 9.120544E-04 C = .2492847 N = 3.162739E-02 S =  
 1.690976E-03

Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

Temperature = 2000  
Use adiabat (1) or insert pressures (2)? 1

ORIGINAL PAGE IS  
OF POOR QUALITY

Water severely depleted; 4x solar C; 2x solar N  
Temperature = 2000 Pressure = 4222.55 bars 50 iterations

## Pressures of gases in bars

$H = 9.855925E-02$   $H_2 = 3692.267$   $He = 516.4525$   $Ne = .9715232$   
 $O = 2.184115E-15$   $O_2 = 1.082947E-23$   $OH = 1.161387E-10$   $H_2O = 4.21309E-03$   
 $N = 3.160501E-10$   $N_2 = .1106789$   $NH_3 = 1.527136$   $CN = 1.503344E-10$   
 $C = 6.91797E-14$   $CO = 2.005459E-07$   $CO_2 = 5.052815E-16$   $CH_4 = 11.02841$   
 $S = 1.189722E-07$   $S_2 = 1.693537E-09$   $HS = 2.583118E-04$   $H_2S = .0932951$   
 $C_2H_6 = 1.523396E-03$   $C_2H_4 = 6.548416E-04$   $C_2H_2 = 1.667547E-05$   
 $CH_3OH = 3.0544E-10$   $HCHO = 7.676327E-10$   $HCOOH = 3.040968E-18$   
 $CH_3NH_2 = 3.81249E-05$   $CH_3CN = 3.113846E-07$   $C_2N_2 = 7.810931E-13$   
 $N_2H_4 = 8.703143E-09$   $HCN = 6.159691E-04$   $CH_3SH = 6.280474E-07$   $SO_2 = 2.004583E-22$

## INPUT ABUNDANCES OF ELEMENTS

$H = 7433.529$   $O = 4.233217E-05$   $C = 11.03346$   $N = 1.748319$   $S = 9.355409E-02$   
Graphite activity = 2.070009E-03

## CALCULATED ABUNDANCES OF ELEMENTS

$H = 7433.527$   $O = 4.233217E-05$   $C = 11.03346$   $N = 1.749148$   $S = 9.355416E-02$   
Type 1 to continue, 0 to end:  
1LIST 2RUN 3LOAD 4SAVE 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 1800  
Use adiabat (1) or insert pressures (2)? 1

Water severely depleted; 4x solar C; 2x solar N  
Temperature = 1800 Pressure = 2971.98 bars 50 iterations

## Pressures of gases in bars

$H = 1.817377E-02$   $H_2 = 2598.79$   $He = 363.5012$   $Ne = .6837993$   
 $O = 6.18104E-17$   $O_2 = 3.782605E-25$   $OH = 6.907098E-11$   $H_2O = 2.976092E-05$   
 $N = 1.00508E-11$   $N_2 = 7.557689E-02$   $NH_3 = 1.079881$   $CN = 1.023402E-11$   
 $C = 2.403227E-16$   $CO = 3.4098E-08$   $CO_2 = 1.034465E-16$   $CH_4 = 7.765272$   
 $S = 1.501565E-08$   $S_2 = 5.093931E-10$   $HS = 3.620836E-05$   $H_2S = 6.580979E-02$   
 $C_2H_6 = 1.698058E-04$   $C_2H_4 = 4.067164E-05$   $C_2H_2 = 4.45051E-07$   
 $CH_3OH = 4.690456E-11$   $HCHO = 9.270799E-11$   $HCOOH = 3.968527E-19$   
 $CH_3NH_2 = 6.652883E-06$   $CH_3CN = 1.486181E-08$   $C_2N_2 = 1.148374E-14$   
 $N_2H_4 = 1.606815E-09$   $HCN = 1.136122E-04$   $CH_3SH = 1.373873E-06$   $SO_2 = 4.268621E-23$

## INPUT ABUNDANCES OF ELEMENTS

$H = 5232.033$   $O = 2.979518E-05$   $C = 7.765815$   $N = 1.230541$   $S = 6.584734E-02$   
Graphite activity = 8.567654E-04

## CALCULATED ABUNDANCES OF ELEMENTS

$H = 5232.031$   $O = 2.979518E-05$   $C = 7.765816$   $N = 1.231155$   $S = 6.584738E-02$   
Type 1 to continue, 0 to end:  
1LIST 2RUN 3LOAD 4SAVE 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 1500  
 Use adiabat (1) or insert pressures (2)? 1

ORIGINAL PAGE IS  
 OF POOR QUALITY

Water severely depleted; 4x solar C; 2x solar N  
 Temperature = 1500 Pressure = 1618.401 bars 70 iterations

Pressures of gases in bars

H = 6.598249E-04 H2 = 1415.136 He = 197.9564 Ne = .3723851  
 O = 8.715959E-20 O2 = 4.661363E-28 OH = 2.941738E-12 H2O = 1.622062E-05  
 N = 1.222253E-16 N2 = 4.05858E-06 NH3 = .670507 CN = 2.212856E-16  
 C = 1.564919E-20 CO = 5.294263E-09 CO2 = 2.366926E-17 CH4 = 4.228946  
 S = 2.397563E-10 S2 = 4.545866E-11 HS = 2.789974E-06 H2S = 3.585612E-02  
 C2H6 = 8.389894E-05 C2H4 = 5.66401E-06 C2H2 = 1.040984E-08  
 CH3OH = 7.325129E-12 HCHO = 8.174095E-12 HCOOH = 4.108515E-20  
 CH3NH2 = 1.500101E-08 CH3CN = 1.256317E-11 C2N2 = 8.543137E-21  
 N2H4 = 7.6027E-15 HCN = 1.011949E-07 CH3SH = 4.319208E-07 SO2 =  
 1.960185E-24

INPUT ABUNDANCES OF ELEMENTS

H = 2849.273 O = 1.622593E-05 C = 4.229126 N = .6701309 S = 3.585931E-02  
 Graphite activity = 8.029344E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 2849.272 O = 1.622593E-05 C = 4.229126 N = .6705152 S = 3.585934E-02  
 Type 1 to continue, 0 to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 1200  
 Use adiabat (1) or insert pressures (2)? 1

Water severely depleted; 4x solar C; 2x solar N  
 Temperature = 1200 Pressure = 769.2399 bars 55 iterations

Pressures of gases in bars

H = 5.091088E-06 H2 = 672.6368 He = 94.08841 Ne = .1769942  
 O = 3.600928E-24 O2 = 2.093047E-32 OH = 2.89253E-14 H2O = 7.712042E-06  
 N = 3.784081E-19 N2 = 8.129111E-03 NH3 = .3024626 CN = 5.926152E-19  
 C = 3.130446E-27 CO = 1.224926E-10 CO2 = 1.029291E-18 CH4 = 2.010084  
 S = 4.872674E-13 S2 = 1.201146E-12 HS = 6.247638E-08 H2S = 1.704379E-02  
 C2H6 = 7.190275E-06 C2H4 = 6.148713E-08 C2H2 = 6.554474E-12  
 CH3OH = 2.499719E-13 HCHO = 9.945679E-14 HCOOH = 6.480291E-22  
 CH3NH2 = 6.733833E-08 CH3CN = 4.210506E-12 C2N2 = 4.153263E-22  
 N2H4 = 5.579431E-13 HCN = 8.062643E-08 CH3SH = 4.08967E-08 SO2 =  
 2.011761E-26

INPUT ABUNDANCES OF ELEMENTS

H = 1354.256 O = 7.712164E-06 C = 2.010099 N = .3185124 S = 1.704388E-02  
 Graphite activity = 2.810095E-04

CALCULATED ABUNDANCES OF ELEMENTS

H = 1354.255 O = 7.712165E-06 C = 2.010098 N = .318721 S = .0170439  
 Type 1 to continue, 0 to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 1000  
Use adiabat (1) or insert pressures (2)? 1

ORIGINAL PAGE IS  
OF POOR QUALITY

Water severely depleted; 4x solar C; 2x solar N  
Temperature = 1000 Pressure = 418.907 bars 60 iterations

Pressures of gases in bars

H = 4.32347E-08 H2 = 366.2962 He = 51.23862 Ne = .0963874  
O = 1.551316E-28 O2 = 9.888972E-37 OH = 3.142237E-16 H2O = 4.199884E-06  
N = 1.23028E-23 N2 = 1.721704E-03 NH3 = .1701121 CN = 1.354106E-23  
C = 5.280051E-34 CO = 2.394E-12 CO2 = 3.96034E-20 CH4 = 1.094657  
S = 1.006242E-15 S2 = 3.202348E-14 HS = 1.526487E-09 H2S = 9.281751E-03  
C2H6 = 6.693206E-07 C2H4 = 6.307591E-10 C2H2 = 3.421414E-16  
CH3OH = 9.746829E-15 HCHO = 1.210671E-15 HCOOH = 1.048438E-23  
CH3NH2 = 3.717071E-09 CH3CN = 1.288063E-14 C2N2 = 1.545482E-25  
N2H4 = 5.682748E-15 HCN = 6.004549E-10 CH3SH = 4.345745E-09 SO2 =  
2.203204E-28

INPUT ABUNDANCES OF ELEMENTS

H = 737.5001 O = 4.199887E-06 C = 1.094659 N = .1734553 S = 9.28175E-03  
Graphite activity = 8.36775E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 737.4999 O = 4.199887E-06 C = 1.094659 N = .1735555 S = 9.281758E-03  
Type 1 to continue, 0 to end:  
1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

Temperature = 800  
Use adiabat (1) or insert pressures (2)? 1

Water severely depleted; 4x solar C; 2x solar N  
Temperature = 800 Pressure = 199.1056 bars 60 iterations

Pressures of gases in bars

H = 3.814574E-11 H2 = 174.0989 He = 24.35378 Ne = 4.581306E-02  
O = 4.671863E-35 O2 = 0 OH = 4.082169E-19 H2O = 1.996211E-06  
N = 2.152343E-30 N2 = 1.425054E-04 NH3 = 8.224059E-02 CN = 1.195233E-30  
C = 0 CO = 5.929854E-15 CO2 = 0 CH4 = .5202926  
S = 9.881199E-20 S2 = 1.457411E-16 HS = 6.445996E-12 H2S = 4.411634E-03  
C2H6 = 2.560806E-08 C2H4 = 9.304709E-13 C2H2 = 4.195508E-20  
CH3OH = 1.269372E-16 HCHO = 2.088923E-18 HCOOH = 0  
CH3NH2 = 7.966937E-11 CH3CN = 2.47847E-18 C2N2 = 2.315859E-32  
N2H4 = 7.86131E-18 HCN = 3.578628E-13 CH3SH = 1.940424E-10 SO2 = 0

INPUT ABUNDANCES OF ELEMENTS

H = 350.5348 O = 1.996212E-06 C = .5202926 N = 8.244354E-02 S =  
4.411628E-03

Graphite activity = 1.226104E-05

CALCULATED ABUNDANCES OF ELEMENTS

H = 350.5346 O = 1.996211E-06 C = .5202926 N = .0825256 S = 4.411634E-03  
Type 1 to continue, 0 to end:  
1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY OSCREEN

Use adiabat (1) or insert pressures (2)? 1

Water severely depleted; 4x solar C; 2x solar N

Temperature = 600 Pressure = 76.31692 bars 65 iterations

Pressures of gases in bars

H = 3.768339E-16 H2 = 66.73183 He = 9.334786 Ne = 1.756011E-02  
 O = 0 O2 = 0 OH = 8.010059E-24 H2O = 7.651463E-07  
 N = 0 N2 = 1.936232E-06 NH3 = 3.162351E-02 CN = 0  
 C = 0 CO = 2.481781E-19 CO2 = 0 CH4 = .1994278  
 S = 2.247766E-26 S2 = 2.688586E-17 HS = 8.488406E-16 H2S = 1.690976E-03  
 C2H6 = 2.068738E-10 C2H4 = 2.138475E-17 C2H2 = 2.703365E-18  
 CH3OH = 1.301233E-19 HCHO = 5.942987E-23 HCOOH = 0  
 CH3NH2 = 1.860118E-13 CH3CN = 1.60161E-24 C2N2 = 0  
 N2H4 = 1.921049E-22 HCN = 1.415396E-18 CH3SH = 1.578961E-12 SO2 = 0

INPUT ABUNDANCES OF ELEMENTS

H = 134.3597 O = 7.651463E-07 C = .1994277 N = 3.160054E-02 S =  
 1.690973E-03

Graphite activity = 4.469427E-07

CALCULATED ABUNDANCES OF ELEMENTS

H = 134.3596 O = 7.651463E-07 C = .1994278 N = 3.162738E-02 S =  
 1.690976E-03

Type 1 to continue, 0 to end:

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 10x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 2000 Pressure = 49.87719 bars 18 iterations

## Pressures of gases in bars

H = 1.070872E-02 H2 = 43.58863 He = 6.100001 Ne = .011475  
 O = 6.501539E-13 O2 = 9.595968E-19 OH = 3.756277E-09 H2O = 1.480545E-04  
 N = 9.068982E-11 N2 = 9.113181E-03 NH3 = 5.62083E-04 CN = 4.199784E-09  
 C = 6.73511E-12 CO = 5.811931E-03 CO2 = 4.35894E-09 CH4 = .1496371  
 S = 1.163633E-07 S2 = 1.620078E-09 HS = 2.74508E-05 H2S = 1.077233E-03  
 C2H6 = 2.375666E-05 C2H4 = 8.650247E-04 C2H2 = 1.865909E-03  
 CH3OH = 1.233653E-09 HCHO = 2.626276E-07 HCOOH = 3.096987E-13  
 CH3NH2 = 1.61279E-08 CH3CN = 1.086305E-06 C2N2 = 6.095932E-10  
 N2H4 = 9.987153E-14 HCN = 1.869681E-03 CH3SH = 8.334689E-09 SO2 =  
 1.737306E-17

## INPUT ABUNDANCES OF ELEMENTS

H = 87.8 O = .00596 C = .1629 N = .02065 S = .001105  
 Graphite activity = .2015293

## CALCULATED ABUNDANCES OF ELEMENTS

H = 87.79988 O = 5.960001E-03 C = .1628306 N = 2.065923E-02 S =  
 1.104812E-03

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 1800 Pressure = 44.88272 bars 18 iterations

## Pressures of gases in bars

H = 2.232697E-03 H2 = 39.22298 He = 5.49 Ne = .0103275  
 O = 1.044425E-13 O2 = 1.079997E-18 OH = 1.433825E-08 H2O = 7.589819E-04  
 N = 3.422217E-12 N2 = 8.761989E-03 NH3 = 6.817684E-04 CN = 2.785055E-10  
 C = 1.920771E-14 CO = 4.604954E-03 CO2 = 2.36063E-08 CH4 = .1413762  
 S = 1.496627E-08 S2 = 5.060485E-10 HS = 4.433667E-06 H2S = 9.899868E-04  
 C2H6 = 3.729257E-06 C2H4 = 5.918232E-05 C2H2 = 4.290822E-05  
 CH3OH = 1.442947E-09 HCHO = 1.889657E-07 HCOOH = 1.366019E-12  
 CH3NH2 = 5.066652E-09 CH3CN = 5.993686E-08 C2N2 = 8.504686E-12  
 N2H4 = 4.243445E-14 HCN = 3.79837E-04 CH3SH = 2.493076E-08 SO2 =  
 1.214754E-16

## INPUT ABUNDANCES OF ELEMENTS

H = 79.02001 O = 5.364001E-03 C = .14661 N = .018585 S = .0009945  
 Graphite activity = 6.847667E-02

## CALCULATED ABUNDANCES OF ELEMENTS

H = 79.01996 O = .005364 C = .146573 N = 1.858565E-02 S = 9.944614E-04

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 10x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 1500 Pressure = 37.38291 bars 60 iterations

## Pressures of gases in bars

H = 1.002367E-04 H2 = 32.65836 He = 4.575 Ne = 8.606251E-03  
 O = 5.930467E-16 O2 = 2.158044E-20 OH = 3.040716E-09 H2O = 2.547049E-03  
 N = 7.879071E-16 N2 = 1.686561E-04 NH3 = 1.515345E-02 CN = 7.614259E-14  
 C = 8.353191E-19 CO = 1.922825E-03 CO2 = 5.849149E-08 CH4 = .1202223  
 S = 2.399968E-10 S2 = 4.554991E-11 HS = 4.242616E-07 H2S = 8.283139E-04  
 C2H6 = 2.938098E-06 C2H4 = 8.594836E-06 C2H2 = 6.844807E-07  
 CH3OH = 1.416909E-09 HCHO = 6.851255E-08 HCOOH = 2.34309E-12  
 CH3NH2 = 4.176242E-10 CH3CN = 8.08963E-10 C2N2 = 1.011501E-15  
 N2H4 = 1.682628E-16 HCN = 5.2897E-06 CH3SH = 1.229115E-08 SO2 =  
 9.084054E-17

## INPUT ABUNDANCES OF ELEMENTS

H = 65.85001 O = .00447 C = .122175 N = .0154875 S = 8.2875E-04  
 Graphite activity = 4.285886E-02

## CALCULATED ABUNDANCES OF ELEMENTS

H = 65.84999 O = 4.469997E-03 C = .122175 N = 1.549606E-02 S = 8.287508E-04

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 1200 Pressure = 29.91357 bars 15 iterations

## Pressures of gases in bars

H = 1.003583E-06 H2 = 26.13762 He = 3.66 Ne = .006885  
 O = 4.241115E-20 O2 = 2.903423E-24 OH = 6.715621E-11 H2O = 3.529558E-03  
 N = 3.045723E-19 N2 = 5.266269E-03 NH3 = 1.864789E-03 CN = 1.535244E-17  
 C = 1.007583E-25 CO = 4.643548E-05 CO2 = 4.595619E-09 CH4 = 9.769209E-02  
 S = 4.877596E-13 S2 = 1.203574E-12 HS = 1.232812E-08 H2S = 6.629643E-04  
 C2H6 = 4.370697E-07 C2H4 = 9.618427E-08 C2H2 = 2.638593E-10  
 CH3OH = 1.430877E-10 HCHO = 1.465076E-09 HCOOH = 1.124309E-13  
 CH3NH2 = 5.192546E-10 CH3CN = 2.689314E-11 C2N2 = 2.787396E-19  
 N2H4 = 5.457837E-16 HCN = 4.117416E-07 CH3SH = 1.989629E-09 SO2 =  
 2.793484E-18

## INPUT ABUNDANCES OF ELEMENTS

H = 52.68 O = .003576 C = 9.774001E-02 N = .01239 S = 6.630001E-04  
 Graphite activity = 9.044731E-03

## CALCULATED ABUNDANCES OF ELEMENTS

H = 52.67999 O = 3.576003E-03 C = 9.774001E-02 N = 1.239774E-02 S =  
 6.629786E-04

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 100x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 2000 Pressure = 49.86685 bars 18 iterations

## Pressures of gases in bars

H = 1.070751E-02 H2 = 43.57882 He = 6.100001 Ne = .011475  
 O = 6.299005E-14 O2 = 9.007418E-21 OH = 3.638853E-10 H2O = 1.4341E-05  
 N = 9.054086E-11 N2 = 9.083266E-03 NH3 = 5.609702E-04 CN = 4.331166E-09  
 C = 6.957232E-12 CO = 5.816583E-04 CO2 = 4.226533E-11 CH4 = .1545025  
 S = 1.164111E-07 S2 = 1.62141E-09 HS = 2.745899E-05 H2S = 1.077433E-03  
 C2H6 = 2.533236E-05 C2H4 = 9.226067E-04 C2H2 = 1.990564E-03  
 CH3OH = 1.234085E-10 HCHO = 2.627787E-08 HCOOH = 3.002237E-15  
 CH3NH2 = 1.662307E-08 CH3CN = 1.156844E-06 C2N2 = 6.483294E-10  
 N2H4 = 9.949889E-14 HCN = 1.927954E-03 CH3SH = 8.609225E-09 SO2 =  
 1.631422E-19

## INPUT ABUNDANCES OF ELEMENTS

H = 87.8 O = .000596 C = .1629 N = .02065 S = .001105  
 Graphite activity = .2081757

## CALCULATED ABUNDANCES OF ELEMENTS

H = 87.80001 O = .000596 C = .1628915 N = 2.065663E-02 S = 1.10502E-03  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N: Lightning discharge  
 Temperature = 1800 Pressure = 44.8745 bars 15 iterations

## Pressures of gases in bars

H = 2.232483E-03 H2 = 39.21545 He = 5.49 Ne = .0103275  
 O = 1.018887E-14 O2 = 1.027826E-20 OH = 1.39863E-09 H2O = 7.40281E-05  
 N = 3.421168E-12 N2 = 8.75662E-03 NH3 = 6.813632E-04 CN = 2.86714E-10  
 C = 1.977989E-14 CO = 4.626175E-04 CO2 = 2.313521E-10 CH4 = .1455318  
 S = 1.496648E-08 S2 = 5.060624E-10 HS = 4.433303E-06 H2S = 9.898106E-04  
 C2H6 = 3.952471E-06 C2H4 = 6.273671E-05 C2H2 = 4.549396E-05  
 CH3OH = 1.44904E-10 HCHO = 1.898001E-08 HCOOH = 1.339285E-14  
 CH3NH2 = 5.21348E-09 CH3CN = 6.352321E-08 C2N2 = 9.0134E-12  
 N2H4 = 4.239216E-14 HCN = 3.909945E-04 CH3SH = 2.566392E-08 SO2 =  
 1.15609E-18

## INPUT ABUNDANCES OF ELEMENTS

H = 79.02001 O = .0005364 C = .14661 N = .018585 S = .0009945  
 Graphite activity = 7.051653E-02

## CALCULATED ABUNDANCES OF ELEMENTS

H = 79.02018 O = 5.366478E-04 C = .1466099 N = 1.858567E-02 S =  
 9.942854E-04  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 10x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 2000 Pressure = 9.980935 bars 33 iterations

## Pressures of gases in bars

H = 4.792545E-03 H2 = 8.730329 He = 1.22 Ne = .002295  
 O = 3.878059E-14 O2 = 3.414171E-21 OH = 1.002731E-10 H2O = 1.768794E-06  
 N = 3.650441E-11 N2 = 1.476534E-03 NH3 = 2.028025E-05 CN = 5.803986E-09  
 C = 2.31237E-11 CO = 1.190231E-03 CO2 = 5.324636E-11 CH4 = 2.060946E-02  
 S = 1.127089E-07 S2 = 1.519918E-09 HS = 1.189947E-05 H2S = 2.089821E-04  
 C2H6 = 2.250001E-06 C2H4 = 4.090426E-04 C2H2 = 4.405274E-03  
 CH3OH = 1.013488E-11 HCHO = 1.07723E-08 HCOOH = 7.577149E-16  
 CH3NH2 = 4.00148E-10 CH3CN = 4.620084E-07 C2N2 = 1.16423E-09  
 N2H4 = 6.491276E-16 HCN = 1.156367E-03 CH3SH = 1.111882E-09 SO2 =  
 5.987076E-20

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .691912

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = 3.259014E-02 N = 4.130187E-03 S = 2.209984E-04  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 1800 Pressure = 9.974824 bars 23 iterations

## Pressures of gases in bars

H = 1.052588E-03 H2 = 8.717609 He = 1.22 Ne = .002295  
 O = 6.128183E-15 O2 = 3.718188E-21 OH = 3.966239E-10 H2O = 9.897892E-06  
 N = 1.580012E-12 N2 = 1.867705E-03 NH3 = 3.298184E-05 CN = 5.625504E-10  
 C = 8.403311E-14 CO = 1.182101E-03 CO2 = 3.555591E-10 CH4 = 3.055377E-02  
 S = 1.488891E-08 S2 = 5.008308E-10 HS = 2.079414E-06 H2S = 2.18895E-04  
 C2H6 = 7.836869E-07 C2H4 = 5.595713E-05 C2H2 = 1.825354E-04  
 CH3OH = 1.829757E-11 HCHO = 1.078125E-08 HCOOH = 4.575636E-15  
 CH3NH2 = 2.383366E-10 CH3CN = 5.549858E-08 C2N2 = 3.469873E-11  
 N2H4 = 4.468252E-16 HCN = 3.61704E-04 CH3SH = 5.360108E-09 SO2 =  
 4.160512E-19

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .2995833

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = 3.257625E-02 N = 4.130151E-03 S = 2.209956E-04  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 1500 Pressure = 9.970698 bars 45 iterations

## Pressures of gases in bars

H = 5.17706E-05 H2 = 8.711785 He = 1.22 Ne = .002295  
 O = 9.122366E-17 O2 = 5.106196E-22 OH = 2.415744E-10 H2O = 1.045126E-04  
 N = 1.238705E-15 N2 = 4.168575E-04 NH3 = 3.282264E-03 CN = 4.401309E-13  
 C = 3.07124E-18 CO = 1.087476E-03 CO2 = 5.088518E-09 CH4 = 3.145372E-02  
 S = 2.398021E-10 S2 = 4.547602E-11 HS = 2.189462E-07 H2S = 2.207777E-04  
 C2H6 = 7.539234E-07 C2H4 = 8.267721E-06 C2H2 = 2.468293E-06  
 CH3OH = 5.702265E-11 HCHO = 1.033625E-08 HCOOH = 5.437514E-14  
 CH3NH2 = 8.872005E-11 CH3CN = 2.368718E-09 C2N2 = 3.379673E-14  
 N2H4 = 2.95937E-17 HCN = 1.579216E-05 CH3SH = 3.213119E-09 SO2 =  
 2.147655E-18

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .1575803

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = 1.191999E-03 C = 3.257999E-02 N = 4.131774E-03 S =  
 2.210002E-04

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 10x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 1200 Pressure = 9.971814 bars 15 iterations

## Pressures of gases in bars

H = 5.794502E-07 H2 = 8.713478 He = 1.22 Ne = .002295  
 O = 3.843064E-20 O2 = 2.383996E-24 OH = 3.513553E-11 H2O = 1.066212E-03  
 N = 1.855947E-19 N2 = 1.95548E-03 NH3 = 2.187224E-04 CN = 2.79641E-17  
 C = 3.011821E-25 CO = 1.257754E-04 CO2 = 1.127944E-08 CH4 = 3.245331E-02  
 S = 4.876982E-13 S2 = 1.203271E-12 HS = 7.117135E-09 H2S = 2.209841E-04  
 C2H6 = 1.446852E-07 C2H4 = 9.551067E-08 C2H2 = 7.259496E-10  
 CH3OH = 4.307241E-11 HCHO = 1.322914E-09 HCOOH = 9.199301E-14  
 CH3NH2 = 6.069008E-11 CH3CN = 2.818394E-11 C2N2 = 9.247952E-19  
 N2H4 = 2.252278E-17 HCN = 4.330231E-07 CH3SH = 6.608712E-10 SO2 =  
 2.293437E-18

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .0270361

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = 1.19201E-03 C = 3.258001E-02 N = 4.130115E-03 S =  
 2.209919E-04

Type 1 to continue, 0 to end:

Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

O depleted by 100x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 2000 Pressure = 9.979018 bars 33 iterations

## Pressures of gases in bars

H = 4.792168E-03 H2 = 8.728957 He = 1.22 Ne = .002295  
 O = 3.777804E-15 O2 = 3.239926E-23 OH = 9.767319E-12 H2O = 1.722796E-07  
 N = 3.63475E-11 N2 = 1.463868E-03 NH3 = 2.018832E-05 CN = 5.932633E-09  
 C = 2.373828E-11 CO = 1.190277E-04 CO2 = 5.187185E-13 CH4 = 2.115057E-02  
 S = 1.127254E-07 S2 = 1.520363E-09 HS = 1.190022E-05 H2S = 2.089798E-04  
 C2H6 = 2.370073E-06 C2H4 = 4.30939E-04 C2H2 = 4.641822E-03  
 CH3OH = 1.013209E-12 HCHO = 1.077103E-09 HCOOH = 7.380391E-18  
 CH3NH2 = 4.088567E-10 CH3CN = 4.846861E-07 C2N2 = 1.216413E-09  
 N2H4 = 6.433568E-16 HCN = 1.181905E-03 CH3SH = 1.141242E-09 SO2 =  
 5.682353E-22

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .0001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .7103015

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = .0001192 C = 3.260275E-02 N = 4.130323E-03 S = 2.20997E-04  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

O depleted by 100x; 5x solar C; 2x solar N; Lightning discharge  
 Temperature = 1800 Pressure = 9.972706 bars 18 iterations

## Pressures of gases in bars

H = 1.052463E-03 H2 = 8.715545 He = 1.22 Ne = .002295  
 O = 5.928473E-16 O2 = 3.479794E-23 OH = 3.836529E-11 H2O = 9.573064E-07  
 N = 1.577683E-12 N2 = 1.862203E-03 NH3 = 3.292153E-05 CN = 5.808037E-10  
 C = 8.688782E-14 CO = 1.182427E-04 CO2 = 3.440665E-12 CH4 = 3.157677E-02  
 S = 1.489276E-08 S2 = 5.010895E-10 HS = 2.079705E-06 H2S = 2.188996E-04  
 C2H6 = 8.372421E-07 C2H4 = 5.979526E-05 C2H2 = 1.951018E-04  
 CH3OH = 1.829394E-12 HCHO = 1.078167E-09 HCOOH = 4.426693E-17  
 CH3NH2 = 2.459244E-10 CH3CN = 5.922487E-08 C2N2 = 3.698703E-11  
 N2H4 = 4.45298E-16 HCN = 3.733961E-04 CH3SH = 5.541004E-09 SO2 =  
 3.894765E-21

## INPUT ABUNDANCES OF ELEMENTS

H = 17.56 O = .0001192 C = .03258 N = .00413 S = .000221  
 Graphite activity = .3097605

## CALCULATED ABUNDANCES OF ELEMENTS

H = 17.56 O = .0001192 C = .03258 N = 4.130784E-03 S = 2.210008E-04  
 Type 1 to continue, 0 to end:  
 Ok

1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

ORIGINAL PAGE IS  
 OF POOR QUALITY

ORIGINAL PAGE IS  
OF POOR QUALITY

II

Venus: Halide Cloud Condensation and  
Volatile Element Inventories

ORIGINAL PAGE IS  
OF POOR QUALITY

### Venus: Halide Cloud Condensation and Volatile Element Inventories

*Abstract. Several recently suggested Venus cloud condensates, including aluminum chloride and halides, oxides, and sulfides of arsenic and antimony, are assessed for their thermodynamic and geochemical plausibility. Aluminum chloride can confidently be ruled out, and condensation of arsenic sulfides on the surface will cause arsenic compounds to be too rare to produce the observed clouds. Antimony may be sufficiently volatile, but the expected molecular form is gaseous antimony sulfide, not the chloride. Arsenic and antimony compounds in the atmosphere will be regulated at very low levels by sulfide precipitation, irrespective of the planetary inventory of arsenic and antimony. Thus arguments for a volatile-deficient origin for Venus based on depletion of water and mercury (relative to the earth) cannot be tested by a search for atmospheric arsenic or antimony.*

Soviet spacecraft have analyzed cloud particles in the main Venus cloud layer by means of x-ray fluorescence (XRF) spectroscopy (1). On the basis of these analyses it is claimed that chlorine is present as a cloud constituent and sulfur is not detected. These results are in conflict with the weight of evidence from Earth-based studies (2) and from other spacecraft experiments (3), which strongly suggest that the dominant cloud constituent on Venus is H<sub>2</sub>SO<sub>4</sub> droplets. Satisfactory photochemical models for the production of an H<sub>2</sub>SO<sub>4</sub> aerosol from geochemically plausible primary gases, including COS, H<sub>2</sub>S, and SO<sub>2</sub>, are available (4). Direct evidence regarding the abundances of these species in the lower troposphere is lacking: copious production of COS and so on by reactions between sulfuric acid and the inlet system of the mass spectrometer on the Pioneer Venus large probe leads to masking of the atmospheric sulfur gases,

and abundances up to 100 ppm are possible for COS and H<sub>2</sub>S (5). No composition data are available for the bottom 22 km of the atmosphere, a region which contains not only the best-equilibrated gases, but also over 80 percent of the total atmospheric mass.


The source of a chlorine-bearing aerosol is less obvious. We have pointed out the high volatility of halides and sulfides of mercury, arsenic, and antimony (6) and have shown that the terrestrial crustal abundance of even the rarest of these elements, mercury, would suffice to produce substantial masses of halide cloud condensates on Venus: mercury is so volatile at the surface temperature of Venus that it would reside almost completely in the atmosphere.

More recently, in the context of a model for the formation of the planets in the presence of a steep gradient outward from the proto-sun, we have favored compositional models in which the vola-

tile elements are severely depleted in the accreting Venus relative to the earth. Combined condensation-accretion models with a variety of recent estimates of the accretion sampling functions of the terrestrial planets (7) show a primordial water content on Venus from 10<sup>-1</sup> to as low as 10<sup>-4</sup> of the terrestrial value. The failure of the 1978 Pioneer Venus mission to detect even a trace of mercury in the lower atmosphere (5) strongly implies that Venus is deficient in mercury relative to the earth. The severe depletion of the most abundant terrestrial volatile, H<sub>2</sub>O, on Venus is well known and is attributed to either a lack of water in preplanetary solids at the orbit of Venus (8) or massive loss of oceans of H<sub>2</sub> after differentiation and outgassing of the planet (9). Donahue (10) found an enhancement of the D/H ratio on Venus by about a factor of 100 over the terrestrial value, requiring either loss of hydrogen from 100 times the present water inventory or accumulation of deuterium-rich material on Venus. Such a late loss mechanism could not deplete mercury while leaving vast amounts of the lighter and more volatile species N<sub>2</sub> and CO<sub>2</sub>. Thus the observed severe depletion of mercury is more convincing evidence for a volatile-poor high-temperature origin of Venus than is the depletion of water. Other moderately volatile elements, such as arsenic, antimony, bismuth, and germanium, are also potential indicators of the overall volatile content of Venus. In addition, such species, if present in the hot lower atmosphere, would condense at intermediate altitudes to form solid halides, sulfides, and oxide cloud particles.

In the past year, Krasnopolsky and Parshev (11) suggested Al<sub>2</sub>Cl<sub>6</sub> as the major cloud layer constituent and Barsukov *et al.* (12) suggested arsenic and antimony halides and oxides. Mole fractions of at least 10 ppm of condensable gases are required to provide the observed cloud density (13). Are these species plausible cloud constituents? Can useful limits on the abundances of the volatile elements arsenic and antimony be derived from atmospheric measurements?

It is simple to estimate the volatility of aluminum compounds at the mean surface conditions of Venus. Consider coexisting anorthite (CaAl<sub>2</sub>Si<sub>2</sub>O<sub>8</sub>), quartz (SiO<sub>2</sub>), and calcite (CaCO<sub>3</sub>) at 750 K and 95 bars in contact with the atmosphere



ORIGINAL PAGE IS  
OF POOR QUALITY

**Venus: Halide Cloud Condensation and  
Volatile Element Inventories**

John S. Lewis and Bruce Fegley, Jr.

The equilibrium constant for this reaction at this temperature is  $10^{-26.1}$ . Assuming the spectroscopic HCl abundance in the clouds, about 1 ppm (14), and the most typical water vapor abundance figures for the lower atmosphere, about 100 ppm (15), an  $\text{AlCl}_3$  partial pressure of  $10^{-21.0}$  bar is calculated. With an extreme effort to bias the equilibrium in favor of  $\text{AlCl}_3$  production (raising HCl to 10 ppm and lowering  $\text{H}_2\text{O}$  to 10 ppm), we can force the  $\text{AlCl}_3$  pressure only as high as  $10^{-16.5}$  bar. This is still ten orders of magnitude too small to provide detectable amounts of condensate. Note that the presence of granitic rocks on Venus, with anorthite and quartz as common primary minerals, is expected both from consideration of the atmospheric composition (16, 17) and from surface passive gamma-ray spectroscopy (18). The surface of Venus lies, as accurately as can be determined, precisely on the calcite-quartz-wollastonite ( $\text{CaSiO}_3$ ) buffer



and calcite is therefore also a plausible surface mineral. Since weathering reactions at higher altitudes on Venus will preferentially tend to mobilize a fine calcium-rich dust, which can be transported readily by winds to the hot lowlands, this buffer may not be difficult to establish on Venus (19).

Thermodynamic treatment of arsenic and antimony volatilization can be carried out without a precise a priori knowledge of the minerals formed by these elements on the surface (20, 21). We will calculate the partial pressures of a number of arsenic and antimony gases at the surface as a function of the activities of arsenic and antimony: an activity of 1 means the pure element is present on the surface and  $10^{-4}$  means the pressure of the monatomic vapor of that element is  $10^{-4}$  times its abundance at saturation. We can then assess the stability of possible surface minerals containing these elements. Figure 1 presents the results for arsenic (22). For activities greater than about  $10^{-2}$  the dominant gas is  $\text{As}_4$ , with  $\text{As}_4\text{O}_6$  second. Based on recent thermochemical data of Johnson *et al.* (18), we calculate that liquid  $\text{As}_2\text{S}_3$  will precipitate if the elemental arsenic activity is greater than 0.13, thus ruling out higher arsenic activities. This, in turn, places firm upper limits on the pressures of arsenic-bearing gases, as indicated in Fig. 1.

The results of a similar calculation for antimony are given in Fig. 2. Precipitation of  $\text{Sb}_4\text{O}_6(\text{s})$  occurs for an elemental

antimony activity of 0.20. Thus for all possible antimony activities,  $\text{SbS}$  is the dominant gas. The best available data on  $\text{Sb}_2\text{S}_3$  (stibnite) from Johnson *et al.* (20) indicate that  $\text{Sb}_2\text{S}_3(\text{s})$  precipitates at an antimony activity of  $10^{-2.3}$ . Therefore we expect that stibnite precipitation on the Venus surface will regulate the antimony gas phase abundance.

We conclude that the mole fraction of all arsenic gases is below  $10^{-7}$ , probably making these species too rare to account for the clouds no matter what species condenses. We expect that the total mole fraction of antimony gases will be lower than for arsenic, but that the most stable




Fig. 1. Partial pressures of arsenic gases as a function of elemental arsenic activity at the Venus surface. Limitations of the arsenic activity imposed by precipitation of liquids of  $\text{As}_2\text{S}_3$  (orpiment) and  $\text{As}_4\text{S}_4$  (realgar) composition are indicated by vertical dashed lines. Heavy dots indicate the upper limits on the  $\text{As}_4$ ,  $\text{As}_4\text{O}_6$ , and  $\text{AsS}$  partial pressures. A condensable species with a partial pressure near  $10^{-4}$  bar is needed to provide the observed cloud density. The maximum total mole fraction of arsenic gases is  $\leq 0.1$  ppm.




Fig. 2. Partial pressure of antimony gases as a function of elemental antimony activity. The  $\text{Sb}_4\text{O}_6$  (valentinite) and  $\text{Sb}_2\text{S}_3$  (stibnite) precipitation points are indicated by vertical dashed lines. Firm upper limits on the antimony gas pressures are indicated by the dots. The maximum total mole fraction of antimony gases is about 0.03 ppm. An  $\text{SbS}$  gas abundance of 0.3 ppm is needed to make clouds.

gas is  $\text{SbS}$ , not a halide. Partial pressures as high as  $10^{-4}$  bar are conceivable for  $\text{SbS}$  and cannot be ruled out.

Observational constraints on the abundance of arsenic in the Venus atmosphere even down to the level of 0.1 ppm are therefore not sufficient to test whether arsenic is, like water and mercury, depleted in Venus relative to the earth: the stability of arsenic sulfides is great enough to preclude a larger abundance of gaseous arsenic compounds irrespective of the crustal abundance of arsenic. The same may be true of antimony, since the mineral  $\text{Sb}_2\text{S}_3$  seems to have low enough volatility to hide antimony in the lithosphere. We have also briefly considered bismuth, and find that  $\text{Bi}_2\text{S}_3$  is so stable that the most abundant bismuth-bearing gas,  $\text{BiS}$ , should have a mole fraction below  $10^{-12}$ .

In any event, we call into serious question the geochemical plausibility of all the species so far suggested as sources of chlorine-bearing clouds. We suggest that either the chlorine compound is a species which has not been considered or the XRF data used to deduce the presence of chlorine are in error.

JOHN S. LEWIS

Department of Earth and Planetary Sciences, Massachusetts Institute of Technology, Cambridge 02139, and Department of Planetary Sciences, University of Arizona, Tucson 85721

BRUCE FEGLEY, JR.

Harvard College Observatory, Cambridge 02138

#### References and Notes

1. Yu. A. Surkov *et al.*, *Geokhimiya* **1981** (No. 1), 3 (1981).
2. A. T. Young, *Icarus* **18**, 564 (1973); J. E. Hansen and J. W. Hovenier, *J. Atmos. Sci.* **31**, 1137 (1974); J. B. Pollack *et al.*, *Icarus* **23**, 8 (1974).
3. R. G. Knollenberg and D. M. Hunten, *Science* **205**, 70 (1979); B. Ragent and J. Blamont, *ibid.* **203**, 790 (1979).
4. R. G. Prinn, *ibid.* **182**, 1132 (1973); *J. Atmos. Sci.* **32**, 1237 (1975).
5. J. H. Hoffman, R. R. Hodges, Jr., M. B. McElroy, T. M. Donahue, M. Kolpin, *Science* **205**, 49 (1979); J. H. Hoffman, R. R. Hodges, T. M. Donahue, M. B. McElroy, *J. Geophys. Res.* **85**, 7882 (1980).
6. J. S. Lewis, *Icarus* **8**, 434 (1968); *ibid.* **11**, 367 (1969).
7. G. W. Wetherill, *Proc. 6th Lunar Sci. Conf.* **2**, 1539 (1975); W. K. Hartmann, *Icarus* **27**, 553 (1976); S. J. Weidenschilling, *ibid.* **22**, 426 (1974); *ibid.* **27**, 161 (1976); L. P. Cox and J. S. Lewis, *ibid.* **44**, 706 (1980).
8. J. S. Lewis, *Science* **186**, 440 (1974); S. S. Barshay, thesis, Massachusetts Institute of Technology (1981); H. D. Holland, in *The Origin and Evolution of Atmospheres and Oceans*, P. J. Brancazio and A. G. W. Cameron, Eds. (Wiley, New York, 1964), p. 86.
9. Y. Shimizu and T. Urabe, *Icarus* **9**, 498 (1968); S. I. Rasool, *J. Atmos. Sci.* **25**, 663 (1968); A. Palm, *Planet. Space Sci.* **17**, 1021 (1969); A. P. Ingersoll, *J. Atmos. Sci.* **26**, 1191 (1969); S. I. Rasool and C. deBergh, *Nature (London)* **226**, 1037 (1970); J. B. Pollack, *Icarus* **14**, 295 (1971); L. L. Smith and S. Gross, *J. Atmos. Sci.* **29**, 173 (1972); K. Turekian and S. P. Clark, *ibid.* **32**, 1257 (1975); J. C. G. Walker, *ibid.*, p. 1248.

10. T. M. Donahue, personal communication.
11. V. A. Krasnopolsky and V. A. Parshev, *Kosm. Issled.* **17**, 763 (1979).
12. V. L. Barsukov, I. L. Khodakovsky, V. P. Volkov, K. P. Florensky, *Space Res.* **20**, 197 (1980).
13. R. G. Knollenberg and D. M. Hunten, *J. Geophys. Res.* **85**, 8039 (1980).
14. P. Connes, J. Connes, W. S. Benedict, L. D. Kaplan, *Astrophys. J.* **147**, 1230 (1967); L. D. G. Young, *Icarus* **17**, 632 (1972).
15. V. I. Oyama, G. C. Carle, F. Woeller, *Science* **208**, 399 (1980).
16. J. S. Lewis and F. A. Kreimendahl, *Icarus* **42**, 330 (1980).
17. I. L. Khodakovsky, V. P. Volkov, Yu. I. Sidorov, V. A. Dorofeeva, M. V. Borisov, *Geokhimiya* **1979** (No. 12), 1747 (1979).
18. A. P. Vinogradov, Yu. A. Surkov, F. F. Kirnozov, *Icarus* **20**, 253 (1973); Yu. A. Surkov, F. F. Kirnozov, V. M. Glazov, G. A. Feodoseyev, *Space Res.* **17**, 659 (1976).
19. S. Nozette and J. S. Lewis, *Science* **216**, 181 (1982).
20. The calculations follow the method of Lewis and Kreimendahl (16). Thermodynamic data are from *JANAF Thermochemical Tables* (National Bureau of Standards, Washington, D.C., 1971); R. Robie, B. S. Hemingway, J. R. Fisher, *U.S. Geol. Surv. Bull.* **1452** (1979); K. C. Mills, *Thermodynamic Data for Inorganic Sulphides, Selenides and Tellurides* (Butterworth, London, 1974); G. K. Johnson, G. N. Papatheodorou, C. E. Johnson, *J. Chem. Thermodyn.* **13**, 745 (1981); *ibid.* **12**, 545 (1980); D. D. Wagman, W. H. Evans, V. B. Parker, R. H. Schumm, R. L. Nuttall, *Natl. Bur. Stand. (U.S.) Tech. Note* **270-8** (1981); and references listed in B. Fegley and J. S. Lewis, *Icarus* **38**, 166 (1979).
21. The pressure, temperature, composition conditions assumed in these calculations are 95 bars total pressure, 750 K surface temperature,  $P_{O_2} = 10^{-22.5}$  bar,  $P_{HCl} = 10^{-4.0}$  bar,  $P_{S_2} = 10^{-4.0}$  bar, and  $P_{HF} = 10^{-5.2}$  bar.
22. Other As- and Sb-bearing gases included in the calculations but not abundant enough to be graphed are As, AsO, AsH<sub>3</sub>, AsF, AsF<sub>2</sub>, AsF<sub>3</sub>, AsF<sub>4</sub>, AsCl, AsCl<sub>2</sub>, AsCl<sub>3</sub>, Sb, SbO, SbH<sub>3</sub>, SbN, and SbF<sub>3</sub>. The solid and liquid arsenic oxides As<sub>2</sub>O<sub>4</sub>, As<sub>2</sub>O<sub>5</sub>, and As<sub>4</sub>O<sub>6</sub> are unstable and do not precipitate as pure condensates on the surface of Venus. Formation of more complex minerals, such as As-bearing apatites and complex Sb oxides, for which thermodynamic data are unavailable, may result in lower gas phase abundances for As and Sb. Thus our results are firm upper limits.
23. We thank the Planetary Atmospheres and Planetary Geophysics and Geochemistry (NGR-22-007-269 to A. G. W. Cameron) program offices of NASA for support of this work through grants to J. S. Lewis Associates, Inc., and to A. G. W. Cameron, Harvard College Observatory. We thank G. K. Johnson for providing his Sb<sub>2</sub>S<sub>3</sub> data before publication. B.F. also thanks A. G. W. Cameron for support and facilities.

23 November 1981; revised 30 March 1982

ORIGINAL PAGE IS  
OF POOR QUALITY

ORIGINAL PAGE IS  
OF POOR QUALITY

III

Shock-Wave Processing of Interstellar  
Cloud Materials

Prior to the discovery of the short-lived radionuclide aluminum-26 in meteorite inclusions by Lee, et al. (1973), the evolution of the presolar nebula was commonly assumed to have taken place in isolation from astrophysical disturbing forces. However, the necessity of having active nucleosynthesis occurring virtually simultaneously with the collapse of the presolar nebula strongly suggests that there is more than mere coincidence at work here. It is, in fact, entirely plausible to believe that the collapse of the nebula was triggered by a nearby supernova explosion. It has become well established that most supernovae occur in star-forming regions: indeed, it is hard to see how it could be otherwise, since the lifetimes of the massive stars responsible for supernova explosions are comparable to or shorter than the lifetimes of the dense molecular cloud complexes that give rise to stars. It is obviously important to understand how the chemical and physical evolution of dense clouds are influenced by nearby supernova explosions.

We consider here the effects of brief, intense shock events on interstellar clouds with pressures of  $10^{-10}$  b or higher; that is, in the densest molecular cloud regions. These regions are most vulnerable to external disturbances, and most likely to be triggered into rapid collapse by a shock event. We take the initial temperature to be 90K, well below the saturation point of ice. Shock heating is relatively ineffectual at heating the solid ice grains, and the release of water vapor into the gas will be incomplete and inhomogeneous at best. The models reported herein assume a  $10^4$ -fold depletion of oxygen relative to carbon in the

ORIGINAL PAGE IS  
OF POOR QUALITY

gas; however, very similar results are found for runs in which the oxygen depletion is only a factor of two.

The results for a number of interesting products are shown in Fig. 1. Note that the abundances of most of the complex molecular products grow rapidly with increasing pressure along the 2000K isotherm. This is not solely a pressure effect, however, since cooling the low-pressure runs to 1800 or 1600 K also gives a large yield of products. It is irrelevant whether the peak shock temperature is too high for optimum yields of complex products (as it clearly is in the low-pressure 2000K runs), since post-shock cooling through more benign temperatures will form these products without any memory of prior, higher-temperature conditions.

A number of features of these results are of interest. First, the dominant carbon compound at high temperatures and low pressures, atomic C (Lewis, et al., 1979) forms HCN, C<sub>2</sub>H and C<sub>2</sub>H<sub>2</sub> very efficiently at slightly lower temperatures. This was documented for C novae by Lewis and Ney (1979).

Note also that, in any C-rich system, oxygen is essentially completely tied up as CO. The multiple-bonded molecules CN, C<sub>2</sub>, CO, CS and N<sub>2</sub> are quite abundant. In analogy with the behavior of oxygen, sulfur is essentially all carried by CS.

Quenching and further slow cooling of the gas mixtures described herein would surely lead to spontaneous polymerization of HCN, C<sub>2</sub>H and C<sub>2</sub>H<sub>2</sub> on grain surfaces. Because of the paucity of oxygen in the gas (due to the very cold initial temperatures of interstellar dense clouds), essentially no water-soluble organics (alcohols, acids, ketones, aldehydes, esters, amino acids, etc.)


ORIGINAL PAGE IS  
OF POOR QUALITY

Caption to Fig. 1.

Mole fractions of gases in an interstellar cloud shocked to 2000K, as a function of the peak shock pressure. For lower temperatures the general behavior is similar, but the abundance curves are shifted to the left.

ORIGINAL PAGE IS  
OF POOR QUALITY

Fig. 1



ORIGINAL PAGE IS  
OF POOR QUALITY

are formed.

Throughout the regime explored, ammonia is a very minor constituent. Thus ammonia, its salts, and organic amines are very minor products from shock heating events.

Conditions conducive to the formation of complex polymeric organic matter, essentially devoid of soluble organic species, should be found in C-nova ejecta, in shocked interstellar clouds, and in shocked regions of the solar nebula in which water is largely condensed prior to the shock.

It will be easy to follow up on these results by modeling the chemical consequences of shock processing of solar nebular material originating both above and below the ice saturation point. Plausible sources of shock energy include supersonic accretion of interstellar material onto the nebular disc, bow shocks in front of massive accreting planetesimals that have attained Keplerian speeds, and lightning produced by charge-separation effects in the turbulent, dusty nebular gas. Cameron's time-dependent accretion disc models permit a reasonable approximate quantification of the first effect, and the latter, based on an old suggestion by Whipple, can be reasonably treated using recent results on the efficiency of lightning production by convective energy fluxes on Venus, Earth, and Jupiter.

The continuation of this project at the University of Arizona, which will be based on computer programs now fully developed under the present grant, will be able to produce publishable work on isolated-grain condensation, on the shock-

wave chemistry of star-forming regions, and on the shock chemistry of the solar nebula. Much of the computer work for these projects has already been done.

```

1 REM *
2 REM best first-guess; 35/65; 470 ITER>12
10 READ BH, PHE, BO, BC, BN, BSU, BNE, BAR
20 FOR NT = 1 TO 10
25 READ T, KH, KH20, KCO, KCH4, KCO2
26 READ T, KOH, KO, KC, KH2S, KHS
27 READ T, KCN, KHCN, KNH3, KSO2, KS
33 READ T, KC2, KC2H, KC2H2, KC3H4AC, KN
34 READ T, KC3H4AL, KC3H6, KC3H8, KC2H6, KC2H4
35 READ T, KC4, KC4H, KC4H2, KCH3NH2, KCH3SH
36 READ T, KCH3CN, KC2N2, KCH, KCH2
37 READ T, KCH3, KCHO, KCH2O, KNH
38 READ T, KNH2, KCS, KCO6, KCS2, KHCNO
39 READ T, KCH3OH, KHCOOH, KN2H4
50 FOR NP = 1 TO 10
59 F = 10^(NP-13)
60 ABH = F*BH
61 ABHE = F*PHE
62 ABO = F*BO
63 ABC = F*BC
64 ABN = F*BN
65 ABSU = F*BSU
66 ABNE = F*BNE
72 ABAR = F*BAR
80 REM INITIAL GUESS OF ELEMENTAL ACTIVITIES
81 RH = (-KH+SQR(KH^2+8*ABH))/4
82 RN = (-KN+SQR(KN^2+8*ABN))/4
83 PH2 = RH^2
84 RO = (2*ABO)/(3*KH20*PH2)
85 GR = ABC/((KCO*RO)+(KCH4*PH2^2))
86 RS = ABSU/((KSO2*RO^2)+(KH2S*PH2))
87 IF ABC>ABO THEN 88 ELSE 99
88 A = KC2H*RH + KC2H2*PH2
89 B = KC + KCH4*PH2^2
91 GR = (-B+SQR(B^2+4*A*ABC))/(2*A)
92 RO = ABO/(KCO*GR+KH20*PH20)
93 RS = ABSU/(KSO2*RO^2+KH2S*PH2+KCS*GR)
94 BB = KHCN*GR*RH + KN
95 RN = (-BB+SQR(BB^2+4*ABN))/2
99 INDEX = 0
100 PRINT "ITER", "INDEX", "ABO/SO", "ABN/SN", "Graphite"
200 FOR ITER = 1 TO 150
201 INDEX = INDEX+1
202 PH = KH*RH
203 PH2 = RH^2
204 PHE = ABHE
205 PNE = ABNE
206 PAR = ABAR
207 PC2 = KC2*GR^2
208 PC2H = KC2H*RH*GR^2
209 PC2H2 = KC2H2*PH2*GR^2
210 PCO = KCO*GR*RO
211 PCH4 = KCH4*GR*PH2^2
212 PCO2 = KCO2*RO*GR*RO
213 PC = KC*GR
214 PC3H4AC = KC3H4AC*PH2^2*GR^3
215 PC3H4AL = KC3H4AL*PH2^2*GR^3
216 PC3H6 = KC3H6*(PH2*GR)^3
217 PC3H8 = KC3H8*PH2*(PH2*GR)^3
218 PC4 = KC4*GR^4
219 PC4H = KC4H*RH*GR^4
220 POH = KOH*RH*RO
221 PH20 = KH20*PH2*RO
222 PO = KO*RO

```

ORIGINAL PAGE IS  
OF POOR QUALITY

```

224 PC4H2 = KC4H2*PH2*GR^4
225 PGH3CN = KCH3CN*RN*GR^2*PH2*RH
226 PC2N2 = KC2N2*PN2*GR^2
227 PCH = KCH*GR*RH
228 PCH2 = KCH2*GR*PH2
229 PCH3 = KCH3*GR*PH2*RH
230 PCN = KCN*GR*RN
231 PHCN = KHCN*GR*RH*RN
232 PNH3 = KNH3*RN*PH2*RH
233 PN2 = RN^2
234 PN = KN*RN
235 PCH0 = KCH0*GR*RH*RO
236 PCH20 = KCH20*GR*PH2^RO
237 PNH = KNH*RN*RH
238 PNH2 = KNH2*RN*PH2
239 PCS = KCS*GR*RS
240 PH2S = KH2S*RS*PH2
241 PS02 = KSO2*RS*PO2
242 PHS = KHS*RH*RS
243 PS = KS*RS
244 PS2 = RS^2
245 PCOS = KCOs*GR*RO*RS
246 PCS2 = KCS2*GR*PS2
247 PHCNO = KHCNO*RH*GR*RN*RO
248 PC2H6 = KC2H6*GR^2*PH2^3
249 PC2H4 = KC2H4*GR^2*PH2^2
250 PCH3NH2 = KCH3NH2*GR*RN*RH*PH2^2
251 PCH3SH = KCH3SH*GR*RS*PH2^2
252 PCH3OH = KCH3OH*GR*RO*PH2^2
253 PHCOOH = KHCOOH*GR*PO2*PH2
254 PN2H4 = KN2H4*PN2*PH2^2
400 REM CALCULATE ELEMENTAL SUMS
401 SH = PH+2*PH2+2*PH20+4*PCH4+3*PNH3+POH+PHS+PHCN+2*PH2S
402 SC = FCO+PCO2+PCH4+PC+PHCN+PCN+PHCNO+PCS2+PCOS+PCS+PCH0+PCH20+PCH+PCH2+PCH3+
2*(PC2N2+PCH3CN+PC2+PC2H+PC2H2)+3*(PC3H4AC+PC3H4AL+PC3H6+PC3H8)+4*(PC4+PC4H+PC4H
2)
403 SO = PH20+2*PO2+2*PCO2+PCO+POH+2*PSO2+PO+PCH0+PCH20+PCOS+PHCNO
404 SN = PN+2*PN2+PNH3+PCN+PHCN+PCH3CN+PNH+PNH2+PHCNO+2*PC2N2
405 SS = PS+PHS+PSO2+PH2S+2*PS2+PCS+PCOS+PCS2
419 P = PH2+PH+PH20+PCO+PCO2+PCH4+PN2+PNH3+POH+PH2S+PHE+PO+PNE+PHS+PAR
420 EH = ABS(SH-ABH)/ABH
421 EO = ABS(SO-ABO)/ABO
422 EC = ABS(SC-ABC)/ABC
423 EN = ABS(SN-ABN)/ABN
424 ES = ABS(SS-ABSU)/ABSU
429 PRINT ITER, INDEX, ABO/SO; ABN/SN; GR
430 IF EH<.001 THEN IF EO<.001 THEN IF EC<.001 THEN IF EN<.001 THEN IF ES<.001 T
HEN GOTO 501 ELSE 431
431 RHNEW = RH
432 RONEW = RO
433 RNNEW = RN
434 RSNEW = RS
435 GRNEW = GR
440 IF INDEX = 1 THEN 444 ELSE 455
444 IF PH>PH2 THEN 445 ELSE 452
445 RHNEW = RH*(ABH/SH)
446 GOTO 470
452 RHNEW = RH*SQR(ABH/SH)
453 GOTO 470
455 IF INDEX = 2 THEN 456 ELSE 462
456 RNNEW = RN*SQR(ABN/SN)
457 RONEW = RO*(ABO/SO)
459 GOTO 470
462 GRNEW = GR*(ABC/SC)
465 RSNEW = RS*(ABSU/SS)
466 INDEX = 0

```

ORIGINAL PAGE IS  
OF POOR QUALITY

```
470 IF ITER<12 THEN 471 ELSE 481
471 RH = RHNEW
472 RO = RONEW
473 RN = RNNEW
474 GR = GRNEW
475 RS = RSNEW
476 GOTO 500
481 RH = .35*RH+.65*RHNEW
482 RO = .35*RO+.65*RONEW
483 RN = .35*RN+.65*RNNEW
484 RS = .35*RS+.65*RSNEW
485 GR = .35*GR+.65*GRNEW
500 NEXT ITER
```

ORIGINAL PAGE IS  
OF POOR QUALITY

```
501 REM          OUTPUT RESULTS OF ITERATION
502 BEEP
505 PRINT "Temperature = ";T, "Pressure = ";P, ITER;"Iterations"
507 PRINT "          Pressures of Gases in Bars"
508 PRINT "H =";PH;" H2 =";PH2;" He =";PHE;" Ne =";PNE
509 PRINT "O =";PO;" O2 =";PO2;" OH =";POH;" H2O =";PH2O
510 PRINT "N =";PN;" N2 =";PN2;" NH3 =";PNH3;" CN =";PCN;" N2H4 =";PN2H4
511 PRINT "C =";PC;" CO =";PCO;" CO2 =";PCO2;" CH4 =";PCH4;" HCN =";PHCN;" SO2 =";
PSO2;" Ar =";PAR
512 PRINT "S =";PS;" S2 =";PS2;" HS =";PHS;" H2S =";PH2S
515 PRINT "C2 =";PC2;" C2H =";PC2H;" C2H2 =";PC2H2;" CH2CCH2 =";PC3H4AL;" CH3CCH
=";PC3H4AC;" C3H8 =";PC3H8;" C4 =";PC4
516 PRINT "C4H =";PC4H;" C4H2 =";PC4H2;" CH3C2H =";PCH3C2H;" CH3CN =";PCH3CN;" C
2N2 =";PC2N2;" CH =";PCH;" CH2 =";PCH2
517 PRINT " CH3 =";PCH3;" CH0 =";PCH0;" CH2O =";PCH2O;" NH =";PNH;" NH2 =";PNH2;"
" CS =";PCS;" CH3OH =";PCH3OH
518 PRINT "CS2 =";PCS2;" HCN0 =";PHCN0;" C2H6 =";PC2H6;" C2H4 =";PC2H4;" CH3NH2
=";PCH3NH2;" CH3SH =";PCH3SH;" HCOOH =";PHCOOH
535 PRINT "          Input Abundances of Elements"
536 PRINT "H =";ABH;" O =";ABO;" C =";ABC;" N =";ABN;" S =";ABSU
538 PRINT "          Calculated Abundances of Elements          Graphite Activity =";GR
```

```
539 PRINT "H =";SH;" O =";SO;" C =";SC;" N =";SN;" S =";SS
545 INPUT "Input (1) to go on; (0) to end: ",GOON
546 IF GOON = 1 THEN 590 ELSE 650
590 NEXT NP
600 NEXT NT
```

```
610 DATA 1756.,122.,1.192e-4,0.6516,0.2065,0.0221,0.2295,6.222E-3
625 DATA 2000, 1.622E-3, 3467.4, 2.944E7, 3.908E-4, 2.254E10
626 DATA 2000, 0.5808, 6.637E-4, 3.342E-11, 0.614, 0.1033
627 DATA 2000, 2.183E-7, 1.472E-2, 2.046E-5, 4.498E5, 2.871E-3
633 DATA 2000, 1.611E-12, 1.714E-6, 8.81E-4, 5.026E-8, 9.5E-10
634 DATA 2000, 2E-8, 2.884E-10, 5.75E-14, 7.063E-9, 1.121E-5
635 DATA 2000, 2.838E-15, 3E-9, 1.6E-6, 6.683E-11, 5.408E-7
636 DATA 2000, 9.736E-7, 1.61E-6, 2.01E-10, 9.099E-9
637 DATA 2000, 1.556E-5, 442.6, 19.28, 2.446E-8
638 DATA 2000, 1.706E-7, 2.286, 6.592E5, 4.227, 15.07
639 DATA 2000, 3.289E-3, 3.674E4, 5.768E-15
650 END
```

```

22      1      .9998351 .9999986 1.875164E-02
Temperature = 2000      Pressure = 1.877169E-09      22 Iterations
      Pressures of Gases in Bars
H = 1.753662E-09 H2 = 1.168934E-12 He = 1.22E-10 Ne = 2.295E-13
O = 1.433318E-25 O2 = 0 OH = 1.356102E-28 H2O = 8.753159E-31
N = 3.039932E-16 N2 = 1.023955E-13 NH3 = 8.274296E-30 CN = 1.309885E-15
N2H4 = 0
C = 6.266797E-13 CO = 1.192197E-16 CO2 = 1.971218E-35 CH4 = 1.001322E-29
HCN = 9.549538E-17 SO2 = 0 Ar = 6.222E-15
S = 1.396518E-15 S2 = 2.333441E-25 HS = 5.395029E-20 H2S = 3.467025E-25
C2 = 5.66466E-16 C2H = 6.516047E-16 C2H2 = 3.621131E-19 CH2CCH2 =
1.801887E-37 CH3CCH = 4.528143E-37 C3H8 = 0 C4 = 3.508885E-22
C4H = 4.010264E-22 C4H2 = 2.31242E-25 CH3C2H = 0 CH3CN = 1.384472E-34 C2N2 =
5.796754E-23 CH = 4.075025E-18 CH2 = 1.994449E-22
CH3 = 3.687515E-25 CHO = 1.937834E-27 CH2O = 0 NH = 8.462368E-21 NH2 =
6.381304E-26 CS = 2.070684E-14 CH3OH = 0
CS2 = 1.84956E-26 HCN0 = 2.111342E-35 C2H6 = 0 C2H4 = 5.385972E-33 CH3NH2 =
0 CH3SH = 0 HCOOH = 0
      Input Abundances of Elements
H = 1.756E-09 O = 1.192E-16 C = 6.516E-13 N = 2.065E-13 S = 2.21E-14
      Calculated Abundances of Elements      Graphite Activity = 1.875164E-02
H = 1.756E-09 O = 1.192197E-16 C = 6.51352E-13 N = 2.065003E-13 S =
2.210341E-14
Input (1) to go on; (0) to end:
1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

```

```

30      3      .9995598 .9999872 .1570265
Temperature = 2000      Pressure = 1.866916E-08      30 Iterations
      Pressures of Gases in Bars
H = 1.733162E-08 H2 = 1.141765E-10 He = 1.22E-09 Ne = 2.295E-12
O = 1.712098E-25 O2 = 0 OH = 1.600928E-27 H2O = 1.021263E-28
N = 9.511923E-16 N2 = 1.002512E-12 NH3 = 2.499283E-26 CN = 3.43219E-14
N2H4 = 0
C = 5.247824E-12 CO = 1.192525E-15 CO2 = 2.355268E-34 CH4 = 7.999832E-25
HCN = 2.472941E-14 SO2 = 0 Ar = 6.222E-14
S = 1.766626E-15 S2 = 3.734162E-25 HS = 6.745051E-19 H2S = 4.283924E-23
C2 = 3.972293E-14 C2H = 4.515906E-13 C2H2 = 2.480267E-15 CH2CCH2 =
1.00949E-30 CH3CCH = 2.536848E-30 C3H8 = 0 C4 = 1.725456E-18
C4H = 1.948952E-17 C4H2 = 1.110678E-19 CH3C2H = 0 CH3CN = 2.932487E-29
C2N2 = 3.979949E-20 CH = 3.372541E-16 CH2 = 1.631335E-19
CH3 = 2.980901E-21 CHO = 1.915709E-25 CH2O = 0 NH = 2.616915E-19 NH2 =
1.950296E-23 CS = 2.193541E-13 CH3OH = 0
CS2 = 2.478554E-25 HCN0 = 6.530945E-33 C2H6 = 0 C2H4 = 3.603337E-27 CH3NH2 =
0 CH3SH = 0 HCOOH = 0
      Input Abundances of Elements
H = 1.756E-09 O = 1.192E-15 C = 6.516E-12 N = 2.065E-12 S = 2.21E-13
      Calculated Abundances of Elements      Graphite Activity = .1570265
H = 1.756E-08 O = 1.192525E-15 C = 6.515433E-12 N = 2.065026E-12 S =
2.211214E-13
Input (1) to go on; (0) to end:
1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

```

ORIGINAL PAGE IS  
OF POOR QUALITY

```

40          1          1.000858  1.000078  .3718258
Temperature = 2000          Pressure = 1.784758E-07          40 Iterations
          Pressures of Gases in Bars
H = 1.568871E-07  H2 = 9.355629E-09  He = 1.22E-08  Ne = 2.295E-11
O = 7.221016E-25  O2 = 0  OH = 6.112088E-26  H2O = 3.529421E-26
N = 2.910689E-15  N2 = 9.387379E-12  NH3 = 5.672667E-23  CN = 2.486941E-13
N2H4 = 0
C = 1.242642E-11  CO = 1.190978E-14  CO2 = 9.920794E-33  CH4 = 1.271862E-20
HCN = 1.622019E-12  SO2 = 0  Ar = 6.222E-13
S = 7.48362E-15  S2 = 6.70081E-24  HS = 2.586428E-17  H2S = 1.48698E-20
C2 = 2.227279E-13  C2H = 2.292062E-11  C2H2 = 1.139536E-12  CH2CCH2 =
8.999006E-26  CH3CCH = 2.261451E-25  C3H8 = 0  C4 = 5.424635E-17
C4H = 5.54646E-15  C4H2 = 2.861219E-16  CH3C2H = 0  CH3CN = 3.731997E-25  C2N2 =
2.089533E-18  CH = 7.228898E-15  CH2 = 3.165237E-17
CH3 = 5.235505E-18  CHO = 1.731865E-23  CH2O = 0  NH = 7.248784E-18  NH2 =
4.890173E-21  CS = 2.200287E-12  CH3OH = 0
CS2 = 1.053172E-23  HCNO = 1.806708E-30  C2H6 = 7.996268E-34  C2H4 =
1.356534E-22  CH3NH2 = 6.445635E-37  CH3SH = 4.556023E-35  HCOOH = 0
          Input Abundances of Elements
H = 1.756E-07  O = 1.192E-14  C = 6.516001E-11  N = 2.065E-11  S = 2.21E-12
          Calculated Abundances of Elements          Graphite Activity = .3718258
H = 1.756E-07  O = 1.190978E-14  C = 6.510593E-11  N = 2.064839E-11  S =
2.207797E-12
Input (1) to go on; (0) to end:
1LIST  2RUN  3LOAD"  4SAVE"  5CONT 6,"LPT1 7TRON 8TROFF9KEY  OSCREEN

```

```

49          1          .9992615  .9998802  .4549451
Temperature = 2000          Pressure = 1.49946E-06          49 Iterations
          Pressures of Gases in Bars
H = 9.983176E-07  H2 = 3.788225E-07  He = 1.22E-07  Ne = 2.295E-10
O = 5.911151E-24  O2 = 0  OH = 3.183792E-24  H2O = 1.169876E-23
N = 8.702916E-15  N2 = 8.392328E-11  NH3 = 4.370191E-20  CN = 9.098168E-13
N2H4 = 6.946709E-38
C = 1.520427E-11  CO = 1.192881E-13  CO2 = 8.134173E-31  CH4 = 2.551438E-17
HCN = 3.775947E-11  SO2 = 0  Ar = 6.222E-12
S = 6.130428E-14  S2 = 4.496612E-22  HS = 1.34822E-15  H2S = 4.93227E-18
C2 = 3.334368E-13  C2H = 2.183467E-10  C2H2 = 6.90764E-11  CH2CCH2 =
2.702579E-22  CH3CCH = 6.791581E-22  C3H8 = 0  C4 = 1.215762E-16
C4H = 7.909975E-14  C4H2 = 2.59652E-14  CH3C2H = 0  CH3CN = 4.304208E-22  C2N2 =
2.796574E-17  CH = 5.628244E-14  CH2 = 1.568153E-15
CH3 = 1.650525E-15  CHO = 1.103795E-21  CH2O = 0  NH = 1.379163E-16  NH2 =
5.920467E-19  CS = 2.205352E-11  CH3OH = 1.91247E-36
CS2 = 8.647224E-22  HCNO = 3.442956E-28  C2H6 = 7.947204E-29  C2H4 =
3.329623E-19  CH3NH2 = 2.460146E-32  CH3SH = 7.487034E-31  HCOOH = 0
          Input Abundances of Elements
H = 1.756E-06  O = 1.192E-13  C = 6.516E-10  N = 2.065E-10  S = 2.21E-11
          Calculated Abundances of Elements          Graphite Activity = .4549451
H = 1.756E-06  O = 1.192881E-13  C = 6.520398E-10  N = 2.065247E-10  S =
2.211618E-11
Input (1) to go on; (0) to end:
1LIST  2RUN  3LOAD"  4SAVE"  5CONT 6,"LPT1 7TRON 8TROFF9KEY  OSCREEN

```

ORIGINAL PAGE IS  
OF POOR QUALITY

ORIGINAL PAGE IS  
OF POOR QUALITY

49 1 9994402 .9998721 .5264224  
 Temperature = 2000 Pressure = 1.209942E-05 49 Iterations  
 Pressures of Gases in Bars  
 H = 4.193159E-06 H2 = 6.683144E-06 He = 1.22E-06 Ne = 2.295E-09  
 O = 5.107626E-23 O2 = 5.922356E-39 OH = 1.155485E-22 H2O = 1.783331E-21  
 N = 2.61154E-14 N2 = 7.556942E-10 NH3 = 9.717395E-18 CN = 3.159084E-12  
 N2H4 = 1.946852E-34  
 C = 1.759304E-11 CO = 1.192668E-12 CO2 = 7.027207E-29 CH4 = 9.188626E-15  
 HCN = 5.50688E-10 SO2 = 0 Ar = 6.222E-11  
 S = 5.298232E-13 S2 = 3.358657E-20 HS = 4.894109E-14 H2S = 7.520246E-16  
 C2 = 4.464412E-13 C2H = 1.22792E-09 C2H2 = 1.631644E-09 CH2CCH2 =  
 1.303151E-19 CH3CCH = 3.274818E-19 C3H8 = 1.673378E-35 C4 = 2.179465E-16  
 C4H = 5.955922E-13 C4H2 = 8.211797E-13 CH3C2H = 0 CH3CN = 1.281426E-19  
 C2N2 = 3.371636E-16 CH = 2.7354E-13 CH2 = 3.20117E-14  
 CH3 = 1.415191E-13 CHO = 4.63536E-20 CH2O = 0 NH = 1.738281E-15 NH2 =  
 3.134245E-17 CS = 2.205431E-10 CH3OH = 5.951235E-33  
 CS2 = 7.473643E-20 HCNO = 4.338687E-26 C2H6 = 5.842522E-25 C2H4 =  
 1.387509E-16 CH3NH2 = 1.116686E-28 CH3SH = 2.33032E-27 HCOOH = 0  
 Input Abundances of Elements  
 H = 1.756E-05 O = 1.192E-12 C = 6.516E-09 N = 2.065E-09 S = 2.21E-10  
 Calculated Abundances of Elements Graphite Activity = .5264224  
 H = 1.756E-05 O = 1.192668E-12 C = 6.519321E-09 N = 2.065264E-09 S =  
 2.211226E-10  
 Input (1) to go on; (0) to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN

Temperature = 2000 Pressure = 1.073048E-04 43 Iterations  
 Pressures of Gases in Bars  
 H = 1.455464E-05 H2 = 8.051949E-05 He = .0000122 Ne = 2.295E-08  
 O = 4.701451E-22 O2 = 5.017879E-37 OH = 3.691792E-21 H2O = 1.977721E-19  
 N = 8.022649E-14 N2 = 7.131623E-09 NH3 = 1.248393E-15 CN = 1.05473E-11  
 N2H4 = 2.666955E-31  
 C = 1.912052E-11 CO = 1.19314E-11 CO2 = 6.470941E-27 CH4 = 1.449606E-12 HCN =  
 6.381839E-09 SO2 = 0 Ar = 6.222E-10  
 S = 4.875208E-12 S2 = 2.843741E-18 HS = 1.563133E-12 H2S = 8.337089E-14  
 C2 = 5.273297E-13 C2H = 5.034403E-09 C2H2 = 2.322007E-08 CH2CCH2 =  
 2.428354E-17 CH3CCH = 6.102453E-17 C3H8 = 4.526389E-31 C4 = 3.040785E-16  
 C4H = 2.884331E-12 C4H2 = 1.380366E-11 CH3C2H = 0 CH3CN = 1.944522E-17  
 C2N2 = 3.758382E-15 CH = 1.031905E-12 CH2 = 4.191679E-13  
 CH3 = 6.432124E-12 CHO = 1.60959E-18 CH2O = 0 NH = 1.853537E-14 NH2 =  
 1.160043E-15 CS = 2.205538E-09 CH3OH = 8.6421E-30  
 CS2 = 6.877263E-16 HCNO = 4.628192E-24 C2H6 = 1.206921E-21 C2H4 =  
 2.378999E-14 CH3NH2 = 1.878501E-25 CH3SH = 3.382805E-24 HCOOH =  
 8.492854E-37  
 Input Abundances of Elements  
 H = .0001756 O = 1.192E-11 C = 6.516E-08 N = 2.065E-08 S = 2.21E-09  
 Calculated Abundances of Elements Graphite Activity = .5721282  
 H = .0001756 O = 1.19314E-11 C = 6.521508E-08 N = 2.065574E-08 S =  
 2.21206E-09  
 Input (1) to go on; (0) to end:  
 1LIST 2RUN 3LOAD" 4SAVE" 5CONT 6,"LPT1 7TRON 8TROFF9KEY 0SCREEN