

NASA Contractor Report 181956

ICASE Report No. 89-77

ICASE

A NOTE CONCERNING THE ONSET OF THREE DIMENSIONALITY AND TIME DEPENDENCE IN GÖRTLER VORTICES

Andrew P. Bassom
Sharon O. Seddougui

Contract No. NAS1-18605
October 1989

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, Virginia 23665-5225

Operated by the Universities Space Research Association

(NASA-CR-181956) A NOTE CONCERNING THE
ONSET OF THREE DIMENSIONALITY AND TIME
DEPENDENCE IN GÖRTLER VORTICES Final Report
(ICASE) 29 p

CSC 01A

N90-14193

Unclass
0252153

G3/02



National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23665-5225



A NOTE CONCERNING THE ONSET OF THREE DIMENSIONALITY AND TIME DEPENDENCE IN GÖRTLER VORTICES

Andrew P. Bassom¹

Department of Mathematics

North Park Road

Exeter, Devon EX4 4QE

U.K.

and

Sharon O. Seddougui

ICASE

NASA Langley Research Center


Hampton, VA 23665

U.S.A.

ABSTRACT

Recently Hall & Seddougui (1989) considered the secondary instability of large amplitude Görtler vortices in a growing boundary layer evolving into a three-dimensional flow with wavy vortex boundaries. They obtained a pair of coupled, linear ordinary differential equations for this instability which constituted an eigenproblem for the wavelength and frequency of this wavy mode. Investigations into the nonlinear version of this problem by Seddougui & Bassom have revealed several omissions in the numerical work of Hall & Seddougui; these issues are addressed in this note. In particular, we find that many neutrally stable modes are possible; we derive the properties of such modes in a high wavenumber limit and show that the combination of the results of Hall & Seddougui and our modifications lead to conclusions which are consistent with the available experimental observations.

¹This research was supported by the National Aeronautics and Space Administration under NASA Contract No. NAS1-18605 while the authors were in residence at the Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, VA 23665.



§1. Introduction.

The purpose of this article is to repeat and improve the calculation of Hall & Seddougui (1989) (hereafter referred to as HS), concerning an asymptotic description of the three-dimensional breakdown of steady, spanwise periodic Görtler vortices. These authors noted that in the experiments of Bippes & Görtler (1972) and of Aihara & Koyama (1981) this breakdown led to a time-periodic flow with wavy vortex boundaries similar to those which occur in the Taylor problem. In order to investigate this phenomenon theoretically, HS superimposed small spanwise periodic travelling waves on the Görtler vortices and monitored their development.

The first analytical work on Görtler vortices concentrated on the linear stability of external flows over concave walls. However Hall (1982*a, b*, 1983) showed that much of this early work was fundamentally flawed for it invoked the parallel flow approximation; but Hall demonstrated that this assumption is unjustifiable except in the limit of small vortex wavelength. Moreover, in this limit, the Görtler instability may be described by an asymptotic structure which accounts for boundary layer growth in a rational manner. This asymptotic structure was obtained by Hall (1982*a*) for the case of infinitesimal amplitude vortices and this was used by Hall (1982*b*) to determine the modified structure in the case of weakly nonlinear vortices. Subsequently, Hall & Lakin (1988) used this latter work to deduce the flow configuration for fully nonlinear, high wavenumber vortices, at which point the mean flow correction generated by the presence of the vortices becomes as large as the basic (undisturbed) flow itself. These fully nonlinear vortices are of the type whose stability to travelling wave disturbances were considered by HS. Hall & Lakin (1988) demonstrated that for these large amplitude vortices the flow structure consists of essentially three distinct regions. The main vortex activity is restricted to a central 'core' region which is bounded by two thin shear layers. The vortices decay exponentially within these shear regions and outside these zones the mean flow is governed by the usual boundary layer equations.

Hall & Seddougui (1989) imposed infinitesimal secondary instabilities upon the flow within the shear layers; these modes took the form of short wavelength, high frequency travelling waves which were $\pi/2$ radians out of phase with the fundamental in the spanwise direction so that any instabilities which occurred produced locally wavy vortex boundaries in the shear layers. It was shown in HS that the governing

equations for these secondary modes take the forms

$$\frac{d^2v}{d\eta^2} - \left(1 + \frac{2i\Omega}{3}\right)\eta v - \frac{2i}{3}Kv - V^2v + \frac{2\sqrt{6}}{3}Vw = 0, \quad (1.1a)$$

$$\frac{d^2w}{d\eta^2} - i(\Omega\eta + K)w + 2V^2w - \frac{i\sqrt{6}}{3}(\Omega\eta + K)Vv = 0. \quad (1.1b)$$

In this pair of coupled ordinary differential equations for the functions $v(\eta)$ and $w(\eta)$, η is an $O(1)$ co-ordinate based upon the thickness of the shear layers, (v, w) are the normal and spanwise components of the velocity of the travelling wave disturbance, and K and Ω are the (dimensionless) wavenumber and frequency of the imposed perturbation. Further, the function $V(\eta)$ satisfies the Painlevé equation,

$$\frac{d^2V}{d\eta^2} - \eta V = V^3, \quad (1.1c)$$

with $V \sim (-\eta)^{\frac{1}{2}}$ as $\eta \rightarrow -\infty$ and $V \sim \sqrt{2}Ai(\eta)$ as $\eta \rightarrow \infty$, see Hastings & McLeod (1980). For a complete description of the derivation of (1.1), together with an extended account of the previous theoretical and practical work relating to Görtler vortices, see HS.

Hall & Seddougui (1989) concentrated on locating solutions of (1.1) for which the flow is neutrally stable, or, in other words, on finding solutions of (1.1) for which K and Ω are real. To ensure that the travelling waves were confined within the shear layers, it was necessary to impose the boundary conditions

$$v, w \rightarrow 0 \quad \text{as} \quad \eta \rightarrow \pm\infty. \quad (1.1d)$$

This then implies that as $\eta \rightarrow \infty$, (1.1a, b) can be written as

$$\frac{d^2v}{d\eta^2} - \left(1 + \frac{2i\Omega}{3}\right)\eta v - \frac{2iK}{3}v = 0, \quad (1.2a)$$

and

$$\frac{d^2w}{d\eta^2} - i(\Omega\eta + K)w = 0, \quad (1.2b)$$

so that in this limit two independent solutions for v and w can be found in terms of the Airy function, Ai . For $\eta \rightarrow -\infty$, HS demonstrated that (1.1a, b) assume the forms

$$\frac{d^2v}{d\eta^2} - \frac{2i\Omega}{3}\eta v - \frac{2iK}{3}v = -\frac{2\sqrt{6}}{3}(-\eta)^{\frac{1}{2}}w, \quad (1.3a)$$

$$\frac{d^2 w}{d\eta^2} - (2 + i\Omega)\eta w - iKw = \frac{i\sqrt{6}}{3}\Omega\eta(-\eta)^{\frac{1}{2}}v + \frac{i\sqrt{6}}{3}K(-\eta)^{\frac{1}{2}}v. \quad (1.3b)$$

Then the appropriate expansions within (1.3) take the form

$$v = (v_{01} + \dots) \exp\left(-\phi|\eta|^{\frac{3}{2}}\right), \quad w = |\eta|^{\frac{1}{2}}(w_{01} + \dots) \exp\left(-\phi|\eta|^{\frac{3}{2}}\right),$$

where

$$243\phi^4 + 36\phi^2(6 + 5i\Omega) - 32\Omega^2 = 0,$$

and the two roots of this equation with positive real part were used in order to generate two independent solutions of (1.3) with $v, w \rightarrow 0$, $\eta \rightarrow -\infty$.

These asymptotic solutions for v and w as $|\eta| \rightarrow \infty$ were taken as initial values in the numerical integration scheme used to solve (1.1). These equations were written as a system of four first order differential equations which was solved using a standard fourth order Runge-Kutta method. The integration procedure was started at $\eta = -\infty$ and at $\eta = \infty$ and continued to $\eta = 0$, finding two independent solutions from each direction. At $\eta = 0$ the continuity of a linear combination of the independent solutions from each direction produced a problem of the form

$$\mathbf{A}\mathbf{x} = \mathbf{0}, \quad (1.4)$$

where $\mathbf{A}(K, \Omega)$ is a 4×4 complex valued matrix and \mathbf{x} is a vector containing the coefficients of the independent solutions from $\eta = \pm\infty$. Clearly there is only a non-trivial solution of (1.4) if $\det(\mathbf{A}) = 0$. HS found real values of K and Ω for which $\det(\mathbf{A}) = 0$ (and hence for which there are non-trivial neutrally stable travelling waves) by employing a Newton-Raphson iteration scheme for two variables.

Hall & Seddougui (1989) focussed their attention on the region $K, \Omega > 0$ and found only one eigenvalue pair, which was located at

$$(K, \Omega) = (4.156, 0.742). \quad (1.5)$$

They speculated that other eigenpairs might possibly exist at higher values of K and Ω , although none were found. Additionally, HS compared these theoretical findings with the results of the experimental observations by Kohama (1987) and Peerhossaini & Wesfreid (1988). They reported good qualitative agreement with these practical results, although the lack of details given by these papers concerning the specific

experimental configurations used prevented a detailed quantitative comparison being made.

Recent work by Seddougui & Bassom (1990) concerns with the extension of HS to the nonlinear regime. Seddougui & Bassom have shown that at the point at which the secondary wavy mode becomes nonlinear, the steady vortex flow of HS is affected by self-interactions of the wavy mode and the problem is then governed by an eighth order coupled set of ordinary differential equations. In the course of this work we reconsidered the linear problem of HS and discovered that the numerical results quoted in that paper are incomplete. In particular, we found eigenvalue pairs lower than (1.5), have shown that there are (plausibly) an infinite number of real valued solution pairs of (1.1), and have obtained an asymptotic description of the solution of this equation for $K \gg 1$.

The procedure for the remainder of this note is as follows. In section 2 we present a revised solution of (1.1) for $O(1)$ values of K and Ω , in section 3 we consider the case $K \gg 1$ and finally we draw some conclusions.

§2. The numerical solution of (1.1)–(1.3).

To obtain real eigenvalues of the system (1.1)–(1.3) we employed the numerical method briefly described in the previous section with one modification. Instead of using the double Newton–Raphson iteration scheme to locate the neutral modes, we considered a whole range of real pairs (K, Ω) and for each chosen pair we computed the matrix \mathbf{A} defined by (1.4). Then, using a standard package, we constructed the contours in the (K, Ω) plane on which $\text{Re}(\det \mathbf{A}) = 0$ or $\text{Im}(\det \mathbf{A}) = 0$. These lines, for the region $0.05 \leq \Omega \leq 3$, $0.1 \leq K \leq 7.9$ are illustrated in Figure (1), where solid lines denote contours on which $\text{Re}(\det \mathbf{A}) = 0$ and broken lines those on which $\text{Im}(\det \mathbf{A}) = 0$. Plainly, where these contours cross we have a solution for infinitesimal, neutrally stable wavy modes. We remark that the apparent multi-crossings of the contours on the left hand side of this figure are not real features and are a direct result of the combination of very small absolute values of the determinant of \mathbf{A} in this area and of the coarseness of the grid from which the contours were interpolated. By repeatedly refining this grid we can show that in reality there are no neutral pairs (K, Ω) in this region.

Having identified approximate values of possible wavenumbers and frequencies for

this secondary mode using this contouring technique, we then applied the Newton-Raphson iteration scheme to obtain more accurate values for these parameters. Overall, we found eight eigenpairs, namely

$$\begin{aligned} (0.690, 0.372), & \quad (2.900, 0.659), & \quad (4.156, 0.742), & \quad (5.435, 0.795), \\ (7.53, 1.00), & \quad (9.60, 1.17), & \quad (11.4, 1.27) & \quad \text{and} & \quad (15.7, 1.60), \end{aligned} \quad (2.1)$$

of which the lowest five are marked on figure (1). We found no more eigenpairs within the size of grid used ($0 < K \leq 16$, $0 < \Omega \leq 3$), but there is no reason to doubt that if this area were suitably enlarged more neutral pairs could be identified. Importantly, we deduce from (2.1) that there are neutral pairs lower than (1.5). In a practical setting, if the frequency of the imposed perturbations on a steady vortex flow were gradually increased from zero, then the mode with the lowest frequency eigenvalue would be expected to be the most dangerous as it would occur first. The eigenfunctions corresponding to this mode are shown in Figure (2) and the asymmetry of these functions is noticeable. Also, for the case plotted here, $\text{Re}(v) > 0$ across the whole of the dominant part of the mode, whereas the spanwise velocity component w has a much more oscillatory nature.

For the higher eigenvalues it was found that it became increasingly more difficult to locate neutral modes accurately and although we have given only eight pairs in (2.1), we believe that there is indeed an infinite sequence of neutral modes. Inspection of the eigenfunctions of Figure (2); of those corresponding to the pair (4.156, 0.742) (presented in HS); and of those corresponding to (9.60, 1.17), illustrated in Figure (3), suggests some definite trends. In particular, the eigenfunctions are effectively confined to a region in η space centred somewhere to the left of $\eta = 0$. Furthermore, for the higher modes, the majority of the disturbance shifts to increasingly more negative values of η , and becomes ever more symmetrical in appearance. The ratio of the spanwise and normal velocity components, $|w|/|v|$, increases. These behaviours led us to an analytical consideration of the problem (1.1)-(1.3) in the limit $K \gg 1$ in an attempt to elucidate the governing behaviour of neutral modes in this high wavenumber limit. This work is the subject of the following section.

§3. The high wavenumber solution of (1.1)-(1.3).

Here we consider the solution of (1.1)–(1.3) for $K \gg 1$. It is found convenient to seek neutral modes in which the wavenumber K and the frequency Ω take the forms

$$K = \left(\beta + \beta^{\frac{2}{3}} K_1 + \dots \right), \quad \Omega = \Omega_0 \beta^{\frac{1}{3}}, \quad (3.1)$$

where $\beta \gg 1$ and Ω_0, K_1 are $O(1)$ constants. Further, we suppose that the mode is concentrated within a region of thickness $O(\beta^{-\frac{1}{3}})$ centred at a point $O(\beta^{\frac{2}{3}})$ from the origin $\eta = 0$. If we then write

$$\eta + \frac{K}{\Omega} = -\beta^{\frac{2}{3}} \eta_c + \beta^{-\frac{1}{3}} y, \quad (3.2a)$$

and note that since for large negative η , $V \sim (-\eta)^{\frac{1}{2}} + \dots$, we also have

$$V = \beta^{\frac{1}{3}} \sqrt{\eta_c + \frac{1}{\Omega_0}} \left[1 - \frac{\beta^{-\frac{2}{3}} \left(y - \frac{K_1}{\Omega_0} \right)}{2 \left(\eta_c + \frac{1}{\Omega_0} \right)} + O(\beta^{-\frac{4}{3}}) \right]. \quad (3.2b)$$

These scalings suggest that

$$v = v_0 + \beta^{-\frac{2}{3}} v_1 + \dots, \quad w = \beta^{\frac{1}{3}} \left(w_0 + \beta^{-\frac{2}{3}} w_1 + \dots \right), \quad (3.3a)$$

and inserting these expansions in (1.1) and equating coefficients of leading powers of β we find that $\eta_c = 0$. At next order we obtain the coupled equations for the functions v_0 and w_0 ,

$$\frac{d^2 v_0}{dy^2} - \frac{2i\Omega_0}{3} y v_0 + \frac{2\sqrt{6}}{3\sqrt{\Omega_0}} w_0 = 0, \quad (3.4a)$$

$$\frac{d^2 w_0}{dy^2} + \left(\frac{2}{\Omega_0} - i\Omega_0 y \right) w_0 - \frac{i\sqrt{6\Omega_0}}{3} y v_0 = 0, \quad (3.4b)$$

with boundary conditions

$$v_0, w_0 \longrightarrow 0 \quad \text{as} \quad y \longrightarrow \pm\infty. \quad (3.4c)$$

By eliminating w_0 between (3.4a, b) we can derive a fourth order differential equation for v_0 which may be solved numerically using standard Runge–Kutta ideas. However, careful inspection and analysis of (3.4) reveals that a solution of the system for which $v_0, w_0 \longrightarrow 0$ as $|\eta| \longrightarrow \infty$ is possible for all non-zero real Ω_0 . Without loss of generality, we can find solutions of (3.4) such that $\text{Re}(v_0)$ and $\text{Re}(w_0)$ are even functions about $y = 0$ whilst $\text{Im}(v_0)$ and $\text{Im}(w_0)$ are odd functions. As Ω_0 is not

determined at this stage, we are forced to consider the next order equations. These are found to be

$$\frac{d^2 v_1}{dy^2} - \frac{2i\Omega_0}{3} y v_1 + \frac{2\sqrt{6}}{3\sqrt{\Omega_0}} w_1 = \frac{\sqrt{6\Omega_0}}{3} \left(y - \frac{K_1}{\Omega_0} \right) w_0, \quad (3.5a)$$

$$\frac{d^2 w_1}{dy^2} + \left(\frac{2}{\Omega_0} - i\Omega_0 y \right) w_1 - \frac{i\sqrt{6\Omega_0}}{3} y v_1 = \left(2w_0 - \frac{i\Omega_0^{\frac{3}{2}} y v_0}{\sqrt{6}} \right) \left(y - \frac{K_1}{\Omega_0} \right), \quad (3.5b)$$

with

$$v_1, w_1 \longrightarrow 0 \quad \text{as} \quad y \longrightarrow \pm\infty. \quad (3.5c)$$

Now as the homogeneous forms of (3.5) are precisely those equations given in (3.4), the set (3.5) has a solution only if a certain compatibility condition holds. To derive this compatibility criterion we consider the system adjoint to (3.4) which in this case is formed by the functions $(g_1(y), g_2(y))$ where

$$\frac{d^2 g_1}{dy^2} - \frac{2i\Omega_0}{3} y g_1 - \frac{i\sqrt{6\Omega_0}}{3} y g_2 = 0, \quad (3.6a)$$

$$\frac{d^2 g_2}{dy^2} + \frac{2\sqrt{6}}{3\sqrt{\Omega_0}} g_1 + \left(\frac{2}{\Omega_0} - i\Omega_0 y \right) g_2 = 0, \quad (3.6b)$$

with

$$g_1, g_2 \longrightarrow 0 \quad \text{as} \quad y \longrightarrow \pm\infty. \quad (3.6c)$$

As before, we can obtain solutions of (3.6) in which the real parts g_1 and g_2 are even functions whilst the respective imaginary parts are odd functions. We multiply (3.5a) by g_1 , (3.5b) by g_2 , add the resulting equations and integrate by parts over the range $(-\infty, \infty)$. Then on comparing imaginary parts of the resulting eigenrelation we find that (3.5) has a solution only if

$$\frac{\sqrt{6\Omega_0}}{3} \int_0^\infty (y \text{Im}(g_1 w_0)) dy + 2 \int_0^\infty (y \text{Im}(g_2 w_0)) dy - \frac{\Omega_0^{\frac{3}{2}}}{\sqrt{6}} \int_0^\infty (y^2 \text{Re}(g_2 v_0)) dy = 0, \quad (3.7)$$

where the evenness and oddness of the various functions about $y = 0$ has been utilised. Incidentally, we note that on equating real parts of the eigenrelation we obtain $K_1 = 0$. Numerically, we found that (3.7) is satisfied for $\Omega_0 \sim 0.26$ and the respective disturbance functions v_0 and w_0 for this case are presented in Figure (4).

This then suggests that high frequency, high wavenumber modes are indeed possible; in particular if $K \gg 1$ we have $\Omega \sim 0.26 K^{\frac{2}{3}}$. Additionally, the position of the

disturbance within the shear layer moves towards the core region of vortex activity, the perturbation is confined within a thin, $O(K^{-\frac{1}{2}})$ sized zone relative to the depth of the shear layer, and the spanwise velocity component of the wavy mode is $O(K^{\frac{1}{2}})$ times that of the normal component. These asymptotic results provide satisfactory agreement with the higher numerical values calculated in (2.1) and it is observed that there is resemblance between the asymptotic eigenfunctions of Figure (4) and those corresponding to the eigenpair (9.60, 1.17) shown in Figure (3). We also remark that here we have not specified K and have only assumed that this is some large quantity. In practice the actual possible values for $K \gg 1$ would be determined by a higher order problem but we do not pursue such a consideration here. Instead, we feel that the principal result of this analysis is that, almost certainly, there is an infinity of neutral wavy modes for this flow structure. This confirms the conjecture presented by HS, who postulated that neutral wavy modes may possibly exist at wavenumbers higher than that for the mode with eigenvalues given by (1.5).

§4. Conclusions.

In this article, which should be read in conjunction with HS, we have demonstrated that there are many solutions for infinitesimal, neutrally stable wavy modes confined to the shear layers of a fully nonlinear, high wavenumber Görtler vortex. This is at variance with the result of Hall & Seddougui (1989) who found only one such mode: although their conclusion, that such a large amplitude vortex is unstable to modes trapped within the shear layers, remains unaltered. Our results are improved from those of HS due to the use of the contouring method described in section 2. This ensured that all possible neutral mode solutions within the region $0 < K \leq 16$, $0 < \Omega \leq 3$ should be identified. This is in contrast to the scheme employed by HS, who used the Newton-Raphson technique alone and relied on finding all possible neutral solutions by using a sufficiently wide variety of initial estimates to which the iteration procedure was applied.

Since the wavy modes are stable when their non-dimensional frequency $\Omega = 0$, HS noted that as Ω is increased the mode with the lowest frequency is potentially more dangerous than the other modes as it occurs first. However, at large values of Ω it is unclear as to which is the most important mode as we would need to identify

the mode with the largest growth rate. The contour plot in Figure (1) shows that $(K, \Omega) = (0.690, 0.372)$ is the lowest mode within the region $K > 0, \Omega > 0$ and that, plausibly, an indefinite number of neutral modes exist. In particular there are modes with $K \gg \Omega \gg 1$ whose asymptotic structure has been discussed in section 3.

The physical and experimental implications of our revised analysis follow similar lines to those given in HS. Rather than repeat the discussion given there and to which the reader is referred, we merely emphasise that our results concerning the possibility of wavy modes in Görtler vortices are consistent with the available experimental accounts on the subject, although it is very difficult to obtain quantitative comparisons between our theory and the practical observations.

Our conclusions concerning the existence of many possible neutral linear modes raises questions of which of these modes is likely to be the most important in practice. This may be resolved by pursuing weakly nonlinear and, ultimately, fully nonlinear analyses of these wavy modes — a task currently being tackled by Seddougui & Basom. When concrete conclusions become available from this study, the full implications of our work for this wavy vortex problem should become clearer.

Acknowledgements.

The authors would like to express their thanks to Professor P. Hall and Dr. P. Blennerhassett for useful discussions. Additionally, we are grateful to Andrew Coward of the University of Exeter for his assistance with the graphics for this article. This work was undertaken whilst the first author was in receipt of financial assistance from SERC, Contract No. XG-10176.

REFERENCES

Aihara, Y. & Koyama, H. 1981 Secondary instability of Görtler vortices: formation of periodic three-dimensional coherent structures. *Trans. Japan Soc. Aero. Space Sci.*, **24**, 78-94.

Bippes, H. & Görtler, H. 1972 Dreidimensionales Störungen in der Grenzschicht an einer Konkaven Wand. *Acta. Mech.*, **14**, 251-267.

Hall, P. 1982a Taylor-Görtler vortices in fully developed or boundary layer flows: linear theory. *J. Fluid Mech.*, **124**, 475-494.

Hall, P. 1982b On the nonlinear evolution of Görtler vortices in non-parallel boundary layers. *IMA J. Appl. Maths.* **29**, 173-196.

Hall, P. 1983 The linear development of Görtler vortices in growing boundary layers. *J. Fluid Mech.*, **130**, 41-58.

Hall, P. & Lakin, W.D. 1988 The fully nonlinear development of Görtler vortices in growing boundary layers. *Proc. R. Soc. Lond. A*, **415**, 421-444.

Hall, P. & Seddougui, S. 1989 On the onset of three-dimensionality and time-dependence in Görtler vortices. *J. Fluid Mech.*, **204**, 405-420.

Hastings, S.P. & McLeod, J.B. 1980 A boundary value problem associated with the Second Painlevé Transcendent and the Korteweg-de Vries equation. *Arch. Ration. Mech. Anal.*, **73**, 31-51.

Kohama, Y. 1987 Three-dimensional boundary layer transition on a convex-concave wall. In *Proc. Bangalore IUTAM Symp. on Turbulence Management and Relaminarisation* (ed. H. Liepmann & R. Narasimha). Springer.

Peerhossaini, H. & Wesfreid, J.E. 1988 On the inner structure of streamwise Görtler rolls. *Intl. J. Heat Fluid Flow*, **9**, 12-18.

Seddougui, S.O. & Bassom, A.P. 1990 On the instability of Görtler vortices to nonlinear travelling waves. *To become ICASE Report 90-1.*

FIGURE CAPTIONS

Figure (1). Sketches of the contours in (K, Ω) space on which the real and imaginary parts of $\det(\mathbf{A})$, defined by (1.4), vanish. On solid lines $\text{Re}(\det \mathbf{A})=0$, on broken lines $\text{Im}(\det \mathbf{A})=0$. We have a solution for neutral wavy modes wherever these contours meet and five such locations are ringed, see (2.1). Here $0.05 \leq \Omega \leq 3$, and (a) $0.1 \leq K \leq 4$, (b) $4 \leq K \leq 7.9$.

Figure (2). Neutral eigenfunctions $v(\eta)$ and $w(\eta)$ for $(K, \Omega) = (0.690, 0.372)$. a) $\text{Re}(v)$, b) $\text{Im}(v)$, c) $\text{Re}(w)$, d) $\text{Im}(w)$.

Figure (3). Neutral eigenfunctions $v(\eta)$ and $w(\eta)$ for $(K, \Omega) = (9.60, 1.17)$. a) $\text{Re}(v)$, b) $\text{Im}(v)$, c) $\text{Re}(w)$, d) $\text{Im}(w)$.

Figure (4). Asymptotic eigenfunctions $v_0(y)$ and $w_0(y)$ (defined by (3.3)) in the limit $K \gg 1$ with $\Omega_0 = 0.26$.




Figure 1(a)




Figure 1(b)




Figure 2(a)




Figure 2(b)




Figure 2(c)




Figure 2(d)




Figure 3(a)




Figure 3(b)




Figure 3(c)




Figure 3(d)




Figure 4(a)




Figure 4(b)




Figure 4(c)




Figure 4(d)



Report Documentation Page

1. Report No. NASA CR-181956 ICASE Report No. 89-77		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle A NOTE CONCERNING THE ONSET OF THREE DIMENSIONALITY AND TIME DEPENDENCE IN GÖRTLER VORTICES				5. Report Date October 1989	
				6. Performing Organization Code	
7. Author(s) Andrew P. Bassom Sharon O. Seddougui				8. Performing Organization Report No. 89-77	
				10. Work Unit No. 505-90-21-01	
9. Performing Organization Name and Address Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23665-5225				11. Contract or Grant No. NAS1-18605	
				13. Type of Report and Period Covered Contractor Report	
				14. Sponsoring Agency Code	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Langley Research Center Hampton, VA 23665-5225					
15. Supplementary Notes Langley Technical Monitor: Richard W. Barnwell Final Report Submitted to Journal of Fluid Mechanics					
16. Abstract Recently Hall & Seddougui(1989) considered the secondary instability of large amplitude Görtler vortices in a growing boundary layer evolving into a three-dimensional flow with wavy vortex boundaries. They obtained a pair of coupled, linear ordinary differential equations for this instability which constituted an eigenproblem for the wavelength and frequency of this wavy mode. Investigations into the nonlinear version of this problem by Seddougui & Bassom have revealed several omissions in the numerical work by Hall & Seddougui; these issues are addressed in this note. In particular, we find that many neutrally stable modes are possible; we derive the properties of such modes in a high wave-number limit and show that the combination of the results of Hall & Seddougui and our modifications lead to conclusions which are consistent with the available experimental observations.					
17. Key Words (Suggested by Author(s)) Görtler vortices, wavy vortices, instability, stability, boundary layers			18. Distribution Statement 02 - Aerodynamics 34 - Fluid Mechanics and Heat Transfer Unclassified - Unlimited		
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of pages 28	22. Price A03

