

NASA SP-7011 (401)
May 1995

P-66

AEROSPACE MEDICINE AND BIOLOGY

(NASA-SP-7011(401)) AEROSPACE
MEDICINE AND BIOLOGY: A CONTINUING
BIBLIOGRAPHY WITH INDEXES
(SUPPLEMENT 401) (NASA) 66 p

N95-25800

Unclas

00/52 0048297

A CONTINUING BIBLIOGRAPHY WITH INDEXES



National Aeronautics and
Space Administration

Scientific and Technical
Information Office

The NASA STI Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Office plays a key part in helping NASA maintain this important role.

The NASA STI Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Office is also NASA's institutional mechanism for disseminating the results of its research and development activities.

Specialized services that help round out the Office's diverse offerings include creating custom thesauri, translating material to or from 34 foreign languages, building customized databases, organizing and publishing research results ... even providing videos.

For more information about the NASA STI Office, you can:

- **Phone** the NASA Access Help Desk at (301) 621-0390
- **Fax** your question to the NASA Access Help Desk at (301) 621-0134
- **E-mail** your question via the **Internet** to help@sti.nasa.gov
- **Write** to:

NASA Access Help Desk
NASA Center for AeroSpace Information
800 Elkridge Landing Road
Linthicum Heights, MD 21090-2934

NASA SP-7011 (401)
May 1995

AEROSPACE MEDICINE AND BIOLOGY

A CONTINUING BIBLIOGRAPHY WITH INDEXES



National Aeronautics and Space Administration
Scientific and Technical Information Office
Washington, DC

1995

This publication was prepared by the NASA Center for Aerospace Information,
800 Elkridge Landing Road, Linthicum Heights, MD 21090-2934, (301) 621-0390.

INTRODUCTION

This issue of *Aerospace Medicine and Biology* (NASA SP-7011) lists 140 reports, articles, and other documents recently announced in the NASA STI Database. The first issue of *Aerospace Medicine and Biology* was published in July 1964.

Accession numbers cited in this issue include:

<i>Scientific and Technical Aerospace Reports (STAR)</i> (N-10000 Series)	N95-19506 — N95-19882
Open Literature (A-60000 Series)	A95-65816 — A95-69893

In its subject coverage, *Aerospace Medicine and Biology* concentrates on the biological, physiological, psychological, and environmental effects to which humans are subjected during and following simulated or actual flight in the Earth's atmosphere or in interplanetary space. References describing similar effects on biological organisms of lower order are also included. Such related topics as sanitary problems, pharmacology, toxicology, safety and survival, life support systems, exobiology, and personnel factors receive appropriate attention. Applied research receives the most emphasis, but references to fundamental studies and theoretical principles related to experimental development also qualify for inclusion.

Each entry in the publication consists of a standard bibliographic citation accompanied, in most cases, by an abstract. The listing of the entries is arranged by *STAR* categories 51 through 55, the Life Sciences division. The citations include the original accession numbers from the NASA STI Database.

Seven indexes—subject, personal author, corporate source, foreign technology, contract number, report number, and accession number—are included.

A cumulative index for 1995 will be published in early 1996.

The NASA CASI price code table, addresses of organizations, and document availability information are located at the back of this issue.



SCAN Goes Electronic!

If you have NASA Mail or if you can access the Internet, you can get biweekly issues of *SCAN* delivered to your desk—top absolutely free!

Electronic SCAN takes advantage of computer technology to alert you to the latest aerospace-related, worldwide scientific and technical information that has been published.

No more waiting while the paper copy is printed and mailed to you. You can review *Electronic SCAN* the same day it is released! And you get all 191—or any combination of—subject areas of announcements with abstracts to browse at your leisure. When you locate a publication of interest, you can print the announcement or electronically add it to your publication order list.

Electronic SCAN
Timely
Flexible
Complete
Free!

For instant access via Internet:

[ftp.sti.nasa.gov](ftp://sti.nasa.gov)
gopher.sti.nasa.gov
listserv@sti.nasa.gov

For additional information:

e-mail: help@sti.nasa.gov
scan@sti.nasa.gov

(Enter this address on the "To" line. Leave the subject line blank and send. You will receive an automatic reply with instructions in minutes.)

Phone: (301) 621-0390 Fax: (301) 621-0134

Write: NASA Access Help Desk
NASA STI Office
NASA Center for AeroSpace Information
800 Elkridge Landing Road
Linthicum Heights, MD 21090-2934



National Aeronautics and
Space Administration
Scientific and Technical
Information Office

TABLE OF CONTENTS

Category 51	Life Sciences (General)	189
Category 52	Aerospace Medicine Includes physiological factors; biological effects of radiation; and effects of weightlessness on man and animals.	209
Category 53	Behavioral Sciences Includes psychological factors; individual and group behavior; crew training and evaluation; and psychiatric research.	211
Category 54	Man/System Technology and Life Support Includes human engineering; biotechnology; and space suits and protective clothing.	213
Category 55	Space Biology Includes exobiology; planetary biology; and extraterrestrial life.	214
Subject Index		A-1
Personal Author Index		B-1
Corporate Source Index		C-1
Foreign Technology Index		D-1
Contract Number Index		E-1
Report Number Index		F-1
Accession Number Index		G-1
Appendix		APP-1

TYPICAL REPORT CITATION AND ABSTRACT

NASA SPONSORED

↓
ON MICROFICHE

ACCESSION NUMBER → N95-10863*# National Aeronautics and Space Administration. ← **CORPORATE SOURCE**
Ames Research Center, Moffett Field, CA.

TITLE → **BIOTELEMETRY IMPLANT VOLUME AND WEIGHT IN RATS:
A PILOT STUDY REPORT**

AUTHOR → CHRIS J. SOMPS May 1994 19 p ← **PUBLICATION DATE**

CONTRACT NUMBER → (Contract RTOP 545-20-01)

REPORT NUMBERS → (NASA-TM-108812; A-94059; NAS 1.15:108812) Avail: CASI HC ← **AVAILABILITY AND PRICE CODE**
A03/MF A01

This paper reports the results of a pilot study in which a 240-gram rat was implanted for 41 days with biotelemetry devices weighing a total of 36 gm (18 cc). The implanted animal showed no differences in weight gain, food and water consumption, and postnecropsy organ weights when compared to both an unoperated control animal and an animal that underwent surgery but did not receive an implant. The implanted animal also had temperature and activity rhythms similar to those reported using much smaller implants. Thus, this pilot study showed that a 240-gm rat could be implanted with biotelemetry devices weighing nearly 15 percent of body weight without significant changes in health or behavior. A larger study involving more animals and similar implant sizes is recommended.

Author

TYPICAL JOURNAL ARTICLE CITATION AND ABSTRACT

NASA SPONSORED

↓

ACCESSION NUMBER → A95-63745* National Aeronautics and Space Administration. John F. ← **CORPORATE SOURCE**
Kennedy Space Center, Cocoa Beach, FL.

TITLE → **THE ORIGIN AND EARLY EVOLUTION OF ISSOL**

AUTHOR → RICHARD S. YOUNG NASA. Kennedy Space Center, Cocoa Beach, ← **AUTHORS' AFFILIATION**
FL, US ISSOL Meeting, 7th, Barcelona, Spain, July 4-9, 1993.

PRIMARY DOCUMENT → A95-63744 Origins of Life and Evolution of the Biosphere (ISSN 0169- ← **JOURNAL TITLE**
6149) vol. 24, no. 2-4. June 1994 p. 83 ← **PUBLICATION DATE**
Copyright

This is a discussion of the beginnings of the International Society for the Study of the Origin of Life (ISSOL)—how it came to be and the people responsible for it. It will include the early meetings on the subject of the Origin of Life which led to the formation of the Society. It will discuss the genesis of the interest of NASA in such a program and how the Exobiology Program got started, leading up to the Viking Program and the early exploration of Mars. Photographs of early meetings and the scientists involved will be included.

Author (Hemer)

AEROSPACE MEDICINE AND BIOLOGY

A Continuing Bibliography (Suppl. 401)

May 1995

51

LIFE SCIENCES (GENERAL)

A95-66753

LABORATORY ASTROPHYSICS AND MOLECULAR ASTRONOMY OF PURE CARBON MOLECULES

P. F. BERNATH Univ. of Waterloo, Waterloo, Ontario, Canada
Life sciences and space research 24 (4): Planetary biology and
origins of life; Topical Meeting of the COSPAR Interdisciplinary
Scientific Commission F (Meeting F3) of the COSPAR Plenary
Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751
Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3
March 1995 p. 15-23

Copyright

The pure carbon molecules C(n) are currently of great experimental and theoretical interest. Our work in this area begins with detection of the SiC molecule, which is isovalent with C2. New infrared electronic transitions of C2 and C3 were discovered by emission spectroscopy of hydrocarbon discharges. The C3 and C5 molecules were found by infrared vibration-rotation spectroscopy of the prototypical obscured carbon star, IRC+10216. C7 and C9 were searched for in the same source, but not found. The laboratory infrared emission spectrum of C60 was recorded to aid in a search for C60 in extraterrestrial sources. Author (Herner)

A95-66754* National Aeronautics and Space Administration, Washington, DC.

TOWARD THE DETECTION OF PURE CARBON CLUSTERS IN THE INTERSTELLAR MEDIUM (ISM)

J. R. HEATH T. J. Watson Research Center, Yorktown Heights, NY, US, A. VAN ORDEN Univ. of California, Berkeley, CA, US, H. J. HWANG Univ. of California, Berkeley, CA, US, E. W. KUO Univ. of California, Berkeley, CA, US, K. TANAKA Kyushu Univ., Higashi-ku, Fukuoka, Japan, and R. J. SAYKALLY Univ. of California, Berkeley, CA, US
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 25-33 Research sponsored by the NSF (Contract(s)/Grant(s): NAGW-2991; NAGW-2763)

Copyright

Determination of the form and distribution of carbon in the universe is critical to understanding the origin of life on Earth and elsewhere. Two potentially large reservoirs of carbon in the interstellar medium (ISM) remain unexplored. These are polycyclic aromatic hydrocarbons (PAH) and pure carbon clusters. Little information exists on the structures, properties, and transition frequencies of pure carbon clusters. The work described is designed to provide a specific inventory of laboratory frequencies and physical properties of this carbon clusters so that efforts can be made to detect them in cold interstellar sources by far-infrared astronomy. Data is given from infrared laser spectroscopy determination of the structure of C3, C4, C5, C6, C7, and C9. Herner

A95-66755* National Aeronautics and Space Administration, Washington, DC.

ORGANIC MOLECULES IN THE GAS PHASE OF DENSE INTERSTELLAR CLOUDS

W. M. IRVINE Univ. of Massachusetts, Amherst, MA, US
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 35-43

(Contract(s)/Grant(s): NAGW-436; NSF AST-91-15721)

Copyright

Since a previous Committee on Space Research (COSPAR) review on this subject, the number of molecular species identified by astronomers in dense interstellar clouds or in the envelopes expelled by evolved stars has grown from about eighty to approximately one hundred. Recent detections in stellar envelopes include the radical CP, the second phosphorus-containing astronomical molecule; SiN, the first astronomical molecule with a Si-N bond; and the HCCN radical. In the dense interstellar clouds recent detections or verifications of previous possible identifications include the H3O(+) ion, which is a critical intermediary in the production of H2O and O2; the CCO radical, which is isoelectronic with HCCN; the SO(+) ion, which appears to be diagnostic of shock chemistry; two new isomers of cyanoacetylene, HCCNC and CCCNH; and the two cumulenes H2C3 and H2C4. Some recent work is also described on the mapping of interstellar clouds in multiple molecular transitions in order to separate variations in chemical abundance from gradients in physical parameters. Author (Herner)

A95-66757

COMETARY ORIGIN OF THE BIOSPHERE: A PROGRESS REPORT

A. H. DELSEMME Univ. of Toledo, Toledo, OH, US
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 49-57

Copyright

Empirical evidence of the accretion temperature for undifferentiated meteorites coming from the asteroid belt, combined with any reasonable temperature gradient extending from the asteroid belt to the Earth's zone, suggests that the Earth accreted from very hot dust grains that were degassed from all volatile elements and depleted in labile compounds. Isotopic evidence from the atmospheric noble gases also shows that no primary atmosphere has survived on the Earth. The only possible source for the atmosphere and the oceans is therefore the cometary bombardment that is predicted as the inescapable consequence of the formation of the giant planets. This implies that comets are the only source of organic carbon, nitrogen and water, hence of the total biosphere of the Earth. Author (Herner)

ABSTRACTS

A95-66758
MASS-SPECTROMETRIC IN SITU STUDIES OF COMETARY ORGANICS FOR P/HALLEY AND OPTIONS FOR THE FUTURE

J. KISSEL Max Planck Institut fuer Kernphysik, Heidelberg, Germany and F. R. KRUEGER Ing.-Bureau Krueger, Darmstadt, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 59-63
 Copyright

When the VEGA and GIOTTO spacecrafts flew by comet p/Halley in 1986 the mass-spectrometers Puma and Particulate Impact Analyzer (PIA) measured the composition of cometary dust particles impacting at speeds of well above 65 km/s. Ion formation upon impact lead to mostly atomic ions. However, a small fractions of the ions measured could be related to molecules. A sophisticated analysis allowed for the first time to point to the chemical nature of cometary organics based on actual mass spectra. With the instrument Cometary Matter Analyzer (CoMA) for the NASA-BMFT mission Comet Rendezvous Asteroid Flyby (CRAF) much higher mass-resolution and molecule masses become accessible for in situ measurement, and will yield complementary information to the gas chromatograph CIDEX also onboard CRAF. Author (Herner)

A95-66759
POSSIBLE EVIDENCE OF HYDROCARBONS RELEASED BY THE NUCLEUS OF HALLEY'S COMET

G. MOREELS Observatoire de Besancon, Besancon, France, J. CLAIREMIDI Observatoire de Besancon, Besancon, France, and P. ROUSSELOT Observatoire de Besancon, Besancon, France Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 65-70
 Copyright

After subtracting the intense dust-scattered continuum from the original spectra transmitted by the Vega 2 three-channel spectrometer, a broad-band emission emerges in the 342-375 nm spectral range when the cometocentric projected distance p is smaller than 5000 km. This newly detected emission varies as $p(\exp -1)$, which implies that the involved molecule(s) has a parent-type behavior. The emission band presents four peaks at 347, 356, 364, and 373 nm. It is tentatively identified as being due to phenanthrene, a three-cycle aromatic condensed hydrocarbon. A determination of the gQ product, where g is the fluorescence quantum efficiency and Q the production rate gives $gQ = 1.2 \times 10(\exp 25)$. If $g = 0.012$, it comes $Q = 1 \times 10(\exp 27)/s$. The detection of phenanthrene in Halley's inner coma is an important argument in favor of a similarity of composition between cometary material and interstellar matter. It supports the hypothesis that comets have kept trace of the interstellar composition through the solar system formation epoch. Author (Herner)

A95-66760
ORGANIC ANALYSIS OF HYDROGEN CYANIDE POLYMERS: PREBIOTIC AND EXTRATERRESTRIAL CHEMISTRY

S. A. LIEBMAN CCS Instrument Systems, Inc., Avondale, PA, US, R. A. PESCE-RODRIGUEZ U.S. Army Research Lab., Aberdeen, MD, US, and C. N. MATTHEWS Univ. of Illinois at Chicago, Chicago, IL, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 71-80
 Copyright

Hydrogen cyanide polymerizes readily to a black solid from

which a yellow-brown powder can be extracted by water and further hydrolyzed to alpha-amino acids. These macromolecules could be major components of the dark matter observed on many bodies in the outer solar system, including comets and asteroids. Primitive Earth might therefore have been covered with HCN polymers through bolide bombardment or by terrestrial synthesis. Several instrumental methods were used for the separation and identification of these intriguing polymeric materials, including photoacoustic Fourier transform infrared spectroscopy, supercritical fluid extraction chromatography and pyrolysis mass spectrometry. Our integrated analytical approach revealed fragmentation patterns and chemical functionalities consistent with the presence of polymeric peptide precursors both in HCN polymers and in the Murchison meteorite. Author (Herner)

A95-66761*
COMETS AND LIFE IN THE UNIVERSE

J. ORO Univ. of Houston, Houston, TX, US, T. MILLS Univ. of Houston, Houston, TX, US, and A. LAZCANO Facultad de Ciencias-UNAM, Mexico, Mexico Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 81-90 Research sponsored by NASA
 Copyright

The notion that comets supplied the primitive Earth with the requisite chemical species for the process of chemical evolution, which is widely held to have led to the origin of life on Earth, has now gained considerable intellectual momentum since its first formulation in 1961. The role of comets in the Earth's biogenesis has been thoroughly addressed in the literature. At this time, in light of a few recent findings, we present here a concise review of this topic together with a brief discussion of the possible role of cometary material in the origin of life elsewhere in the Universe. Author (Herner)

A95-66762* National Aeronautics and Space Administration, Washington, DC.

CHARACTERISTICS AND FORMATION OF AMINO ACIDS AND HYDROXY ACIDS OF THE MURCHISON METEORITE

J. R. CRONIN Arizona State Univ., Tempe, AZ, US, G. W. COOPER Arizona State Univ., Tempe, AZ, US, and S. PIZZARELLO Arizona State Univ., Tempe, AZ, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 91-97
 (Contract(s)/Grant(s): NAGW-1899; NGT-70100)
 Copyright

Eight characteristics of the unique suite of amino acids and hydroxy acids found in the Murchison meteorite can be recognized on the basis of detailed molecular and isotopic analyses. The marked structural correspondence between the alpha-amino acids and alpha-hydroxy acids and the high deuterium/hydrogen ratio argue persuasively for their formation by aqueous phase Strecker reactions in the meteorite parent body from presolar, i.e., interstellar, aldehydes, ketones, ammonia, and hydrogen cyanide. The characteristics of the meteoritic suite of amino acids and hydroxy acids are briefly enumerated and discussed with regard to their consonance with this interstellar-parent body formation hypothesis. The hypothesis has interesting implications for the organic composition of both the primitive parent body and the presolar nebula. Author (Herner)

A95-66763* National Aeronautics and Space Administration, Washington, DC.

STABLE ISOTOPE ANALYSIS AT THE MOLECULAR LEVEL: A NEW APPROACH FOR DETERMINING THE ORIGINS OF AMINO ACIDS IN THE MURCHISON METEORITE

M. H. ENGEL Univ. of Oklahoma, Norman, OK, US, S. A. MACKO Univ. of Virginia, Charlottesville, VA, US, Y. QIAN Univ. of

Oklahoma, Norman, OK, US, and J. A. SILFER Univ. of Oklahoma, Norman, OK, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 99-106
(Contract(s)/Grant(s): NAGW-2765)
Copyright

A combined gas chromatography/isotope ratio mass spectrometry (GC/IRMS) method has been developed that permits the direct stable carbon isotope analysis of N(O)-trifluoroacetyl-isopropyl esters of individual amino acids and their respective enantiomers at nanomole abundances. Calculation of the original delta C-13 values of the amino acids is accomplished via a correction for the carbon introduced during the derivatization process. Previous GC/IRMS analyses of individual amino acids in the non-hydrolyzed water extract of an interior sample of a Murchison meteorite stone revealed an enrichment in C-13 relative to terrestrial organic matter, in agreement with previous findings for bulk extracts. The range of amino acid delta C-13 values (+5 to +30 per mill, PDB) suggests possible kinetic effects during synthesis. In this study, an apparent kinetic isotope effect was also observed for the amino acid products of a spark discharge experiment. These preliminary results are supportive of a similar mechanism for the abiotic synthesis of amino acids in the Murchison meteorite. Author (Herner)

A95-66765
WERE MICROMETEORITES A SOURCE OF PREBIOTIC MOLECULES ON THE EARLY EARTH?

M. MAURETTE Centre de Spectrometrie Nucleaire, Orsay-Campus, France, A. BRACK Centre de Biophysique Moleculaire du CNRS, Orleans, France, G. KURAT Naturhistorisches Museum, Wien, Austria, M. PERREAU ONERA, Chatillon, France, and C. ENGRAND Centre de Spectrometrie Nucleaire, Orsay-Campus, France Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 113-126 Research sponsored by the Austrian FWF (Contract(s)/Grant(s): EEC-SC1-CT-91-0618)
Copyright

Interplanetary Dust Particles (IDPs) with sizes approximately 10 micrometer collected in the stratosphere (IDPs), as well as much larger 'giant' micrometeorites retrieved from Antarctic ice melt water (AMMs), are mostly composed of unequilibrated assemblages of minerals, thus being related to primitive unequilibrated meteorites. Two independent evaluations of the mass flux of micrometeorites measuring approximately 50 micrometer to approximately 200 micrometer, recovered from either the Greenland or the Antarctic ice sheets have been reported (approximately 20,000 tons/a). A comparison with recent evaluation of the flux of meteorites reaching the Earth's surface (up to masses of 10,000 tons), indicates that micrometeorites represent about 99.5% of the extraterrestrial material falling on the Earth's surface each year. As they show carbon concentrations exceeding that of the most C-rich meteorite (Orgueil), they are the major contributors of extraterrestrial C-rich matter accreting to the Earth today. Moreover they are complex microstructured aggregates of grains. They contain not only a variety of C-rich matter, such as a new 'dirty' magnetite phase enriched in P, S, and minor elements, but also a diversity of potential catalysts (hydrous silicates, oxides, sulfides and metal grains of Fe/Ni composition, etc.). They could have individually functioned on the early Earth, as 'micro-chondritic-reactors' for the processing of prebiotic organic molecules in liquid water. Future progress requires the challenging development of meaningful laboratory simulation experiments, and a better understanding of the partial reprocessing of micrometeorites in the atmosphere. Author (Herner)

A95-66767
RADIATION AND THERMAL STABILITIES OF ADENINE NUCLEOTIDES

V. V. DEMIDOV Russian Academy of Sciences, Kurchatov Square, Moscow, Russia, V. N. POTAMAN Russian Academy of Sciences, Kurchatov Square, Moscow, Russia, I. P. SOLYANINA Scientific Industrial Association Biotechnology, Moscow, Russia, and V. I. TROFIMOV Scientific Industrial Association Biotechnology, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 131-134
Copyright

We have investigated in detail radiation and thermal stabilities and transformations of adenosine mono- and triphosphates in liquid and frozen solid aqueous solutions within a wide range of absorbed radiation dose (up to 75 kGy) and temperature (up to 160 C). Dephosphorylation is the main pathway of high temperature hydrolysis of adenine nucleotides. Basic thermodynamic and kinetic parameters of this process have been determined. Radiolysis of investigated compounds at room temperature results in scission of N-glycosidic bond with a radiation yield about of 1 mol/100 eV. Solution freezing significantly enhances radiation stability of nucleotides as well as other biomolecules. This circumstance is essential in the discussion of panspermia concepts. Author (Herner)

A95-66769
EXTINCTION OF DINOSAURS: A POSSIBLE NOVEL CAUSE

S. RAMADURAI Tata Inst. of Fundamental Research, Bombay, India, DAVID LLOYD Univ. of Wales, Cardiff, UK, MAX WALLIS Univ. of Wales, Cardiff, UK, and N. C. WICKRAMASINGHE Univ. of Wales, Cardiff, UK Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 139-146 Research sponsored by INDO-US, the Univ. of Wales, and Queen Mary and Westfield College, London, UK
Copyright

A novel cause of mass extinction of fauna close to the Cretaceous/Tertiary (K/T) Boundary is suggested. A large amount of non-protein amino acids alpha-amino isobutyric acid (AIB) and isovaline (ISOVAL) has been observed close to this event. It is speculated that these amino acids may be toxic and are responsible for the extinction. The toxicity level is estimated for this suggestion to be true and experimentalists are encouraged to test this level of toxicity for the amino acids. Author (Herner)

A95-66777* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

A PRELIMINARY COMPARISON OF TWO PERENNIALY ICE-COVERED LAKES IN ANTARCTICA: ANALOGS OF PAST MARTIAN LACUSTRINE ENVIRONMENTS

D. T. ANDERSEN SETI Inst., Mt. View, CA, US, P. DORAN Desert Research Inst., Reno, NV, US, D. BOLSHIYANOV Inst. of Arctic and Antarctic Research, St. Petersburg, Russia, J. RICE Arizona State Univ., Tempe, AZ, US, V. GALCHENKO Russian Academy of Sciences, Moscow, Russia, N. CHERYCH Russian Academy of Sciences, Moscow, Russia, R. A. WHARTON Desert Research Inst., Reno, NV, US, C. P. MCKAY NASA. Ames Research Center, Moffett Field, CA, US, M. MEYER Lockheed Engineering and Sciences Co., Washington, DC, US, and V. GARSHNEK Lockheed Engineering and Sciences Co., Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 199-202

Copyright

Perennially ice-covered lakes in the Antarctic have been suggested as analogs to lakes which may have existed on the surface of Mars 3.5 billion years ago. During the 1991-1992 austral summer, a joint Russian/American research effort was directed at studies of ice-covered lakes in the Bunger Hills Oasis, Antarctica (66 deg S, 100 deg E). The primary objective of the expedition was to investigate this ice-free area for features analogous to ancient martian environments that may have been capable of supporting life and to compare the ice-covered lakes of the Bunger Hills with those in the McMurdo Dry Valleys of southern Victoria Land (77 deg S, 166 deg E) as part of the continuing studies of Antarctic-Mars analogs. Author (Herner)

A95-66778

BIOCHEMICAL CONSTRAINTS FOR SURVIVAL UNDER MARTIAN CONDITIONS

K. DOSE J. Gutenberg-Universitaet, Mainz, Germany, C. STRIDDE J. Gutenberg-Universitaet, Mainz, Germany, R. DILLMANN J. Gutenberg-Universitaet, Mainz, Germany, S. RISI J. Gutenberg-Universitaet, Mainz, Germany, and A. BIEGER-DOSE J. Gutenberg-Universitaet, Mainz, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 203-210

Copyright

A wide variety of terrestrial organisms, the so-called 'anhydrobiotes,' have learned to survive in a state of extreme dehydration in dry environments. Strategies for survival include the accumulation of certain polyols and nonreducing saccharides, which help to prevent damage to membranes and proteins, but at low water partial pressure DNA is also progressively damaged by various lesions, including strand breaks and cross-linking to proteins. These lesions, if they are not too numerous, can be repaired before the first replication step after rehydration, but long-term exposure to dry conditions finally diminishes the chances of survival as these lesions accumulate. If an organism has no chance to repair the accumulated DNA damage during intermittent periods of active life, survival will not exceed a few decades. The restriction of survival by dryness-induced DNA lesions is corroborated by new data on conidia of *Aspergillus* and the free plasmid pBR 322. Our results will be discussed with respect to the chance of finding dormant life or biochemical fossils on the surface of Mars. Author (Herner)

A95-66779

STUDIES IN THE SEARCH FOR LIFE ON MARS

J. KOIKE Tokyo Inst. of Tech., Yokohama, Japan, T. OSHIMA Tokyo Inst. of Tech., Yokohama, Japan, K. KOBAYASHI Yokohama National Univ., Yokohama, Japan, and Y. KAWASAKI Mitsubishi-kasei Inst. of Life, Machida, Tokyo, Japan Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 211-214 Research sponsored by the Ministry of Education of Japan

Copyright

The ability of living organisms to survive extraterrestrial conditions has implications for the origins of life in the solar system. We have therefore studied the survival of viruses, bacteria, yeast, and fungi under simulated Martian conditions. The environment on Mars was simulated by low temperature, proton irradiation, ultraviolet irradiation, and simulated Martian atmosphere (CO₂ 95.46%, N₂ 2.7%, water vapor 0.03%) in a special cryostat. After exposure to these conditions, tobacco mosaic virus and spores of *Bacillus*, *Aspergillus*, *Clostridium*, and some species of coccus showed significant survival. Author (Herner)

A95-66780

BIOGEOCHEMICAL EVIDENCE OF MICROBIAL ACTIVITY**ON MARS**

M. V. IVANOV Russian Academy of Science, Moscow, Russia and A. YU. LEIN Russian Academy of Science, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 215-221

Copyright

We suggest a new interpretation of the data on and chassignites (SNC meteorites) and delta C-13 values of the calcium carbonate minerals and organic matter discovered in them. The delta C-13 value of calcite (up to 15 ppt) is accounted for by the microbial reaction CO₂ + H₂ yields CH₄ + H₂O. Methane-forming bacteria also synthesize organic carbon (in the form of biomass) from CO₂, and this process is accompanied by C-12 fractionation. Therefore, the organic carbon of SNC meteorites is enriched with C-12 (delta C-13 as low as -35 ppt). The environmental conditions under which the calcite of SNC meteorites was formed were favorable for the activity of methanogens. Author (Herner)

A95-66781* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

A 'CRYTIC' MICROBIAL MAT: A NEW MODEL ECOSYSTEM FOR EXTANT LIFE ON MARS

L. J. ROTHSCHILD NASA. Ames Research Center, Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 223-228

Copyright

If life were present on Mars today, it would face potentially lethal environmental conditions such as a lack of water, frigid temperatures, ultraviolet radiation, and soil oxidants. In addition, the Viking missions did not detect near-surface organic carbon available for assimilation. Autotrophic organisms that lived under a protective layer of sand or gravel would be able to circumvent the ultraviolet radiation and lack of fixed carbon. Two terrestrial photosynthetic near-surface microbial communities have been identified, one in the inter- and supertidal of Laguna Ojo de Liebere (Baja California Sur, Mexico) and one in the acidic gravel near several small geysers in Yellowstone National Park (Wyoming, U.S.A.). Both communities have been studied with respect to their ability to fix carbon under different conditions, including elevated levels of inorganic carbon. Although these sand communities have not been exposed to the entire suite of Martian environmental conditions simultaneously, such communities can provide a useful model ecosystem for a potential extant Martian biota. Author (Herner)

A95-66783

PRESERVATION OF CELL STRUCTURES IN PERMAFROST: A MODEL FOR EXO BIOLOGY

V. S. SOINA Moscow State Univ., Moscow, Russia, E. A. VOROBIOVA Moscow State Univ., Moscow, Russia, D. G. ZVYAGINTSEV Moscow State Univ., Moscow, Russia, and D. A. GILICHINSKY Russian Academy of Sciences, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 237-242

Copyright

The present report is the first contribution toward a comprehensive fine-structural study of microbial cells from permafrost. Prokaryotes with a variety of cell wall types demonstrate high stability of cell structure after long-term cryopreservation in frozen soils and sediments of the Arctic. The surface capsular layers that were a salient

feature of the cells both in situ and on nutrient media may be an adaptation to low temperature. To the extent that permafrost regions on Earth approximate Martian conditions, preservation of cell structure there can serve as the basis for predictions about preservation in Martian permafrost sediments. Author (Herner)

A95-66784

A PRIMITIVE CYANOBACTERIUM AS PIONEER MICROORGANISM FOR TERRAFORMING MARS

E. IMRE FRIEDMANN Florida State Univ., Tallahassee, FL, US and R. OCAMPO-FRIEDMANN Florida State Univ., Tallahassee, FL, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 243-246* Copyright

The primitive characteristics of the cyanobacterium *Chroococcidiopsis* suggest that it represents a very ancient type of this group. Its morphology is simple but shows a wide range of variability, and it resembles certain Proterozoic microfossils. *Chroococcidiopsis* is probably the most desiccation-resistant cyanobacterium, the sole photosynthetic organism in extreme arid habitats. It is also present in a wide range of other extreme environments, from Antarctic rocks to thermal springs and hypersaline habitats, but it is unable to compete with more specialized organisms. Genetic evidence suggests that all forms belong to a single species. Its remarkable tolerance of environmental extremes makes *Chroococcidiopsis* a prime candidate for use as a pioneer photosynthetic microorganism for terraforming of Mars. The hypolithic microbial growth form (which lives under stones of a desert pavement) could be used as a model for development of technologies for large-scale Martian farming. Author (Herner)

A95-66793* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

SURFACE-ATMOSPHERE INTERACTIONS ON TITAN COMPARED WITH THOSE ON THE PRE-BIOTIC EARTH

J. I. LUNINE Univ. of Arizona, Tucson, AZ, US and C. P. MCKAY NASA. Ames Research Center, Moffett Field, CA, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 303-311* Research sponsored by NASA Copyright

The surface and atmosphere of Titan constitute a system which is potentially as complex as that of the Earth, with the possibility of precipitation, surface erosion due to liquids, chemistry in large surface or subsurface hydrocarbon reservoirs, surface expressions of internal activity, and occasional major impacts leading to crustal melting. While none of the above have been observed as yet, the composition, density and thermal properties of Titan's atmosphere make it uniquely suited in the outer solar system as a place where such processes may occur. The one attribute of the Earth not expected on Titan is biological activity, which has had a profound effect on the evolution of the Earth's surface-atmosphere system. The earliest environment of Titan could have been warm enough for liquid ammonia-water solutions to exist on or near surface; pre-biotic organic processes may have taken place in such an environment. After a few hundred million years surface ammonia-water would have disappeared. Therefore, study of Titan through Cassini/Huygens mission, planned for launch in 1997, primarily affords the opportunity to understand planet-side surface-atmosphere interactions in the presence of fluids but in the absence of life. More speculative is the possibility that endogenic and exogenic heating continue to provide short-lived environments on Titan wherein pre-biotic organic processes in the presence of water happen. Author (Herner)

Author (Herner)

A95-66794* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

THE PHYSICAL NATURE OF TITAN'S AEROSOLS: LABORATORY SIMULATIONS

T. W. SCATTERGOOD NASA. Ames Research Center, Moffett Field, CA, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 313-316* Copyright

The atmosphere of Titan is known to contain aerosols, as evidenced by the Voyager observations of at least three haze layers. Such aerosols can have significant effects on the reflection spectrum of Titan and on the chemistry and thermal structure of its atmosphere. To investigate some of these effects laboratory simulations of the chemistry of Titan's atmosphere have been done. The results of these studies show that photolysis of acetylene, ethylene, and hydrogen cyanide, known constituents of Titan's atmosphere, yields sub-micron sized spheres, with mean diameters ranging from 0.4 to 0.8 micron, depending on the pressures of the reactant gases. Most of the spheres are contained in near-linear aggregates. The formation of the aggregates is consistent with models of Titan's reflection spectrum and polarization, which are best fit with non-spherical particles. At room temperature, the particles are very sticky, but their properties at low temperatures on Titan are presently not known. Author (Herner)

Author (Herner)

A95-66797* National Aeronautics and Space Administration, Washington, DC.

ANALYTICAL PYROLYSIS EXPERIMENTS OF TITAN AEROSOL ANALOGUES IN PREPARATION FOR THE CASSINI HUYGENS MISSION

P. EHRENFREUND Service d'Aeronomie, Verrieres, France, J. J. BOON FOM-Inst., Amsterdam, Holland, J. COMMANDEUR FOM-Inst., Amsterdam, Holland, C. SAGAN Cornell Univ., Ithaca, NY, US, W. R. THOMPSON Cornell Univ., Ithaca, NY, US, and B. KHARE Cornell Univ., Ithaca, NY, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 335-342* Research sponsored by SRON, and NWO (Contract(s)/Grant(s): NAGW-1896) Copyright

Comparative pyrolysis mass spectrometric data of Titan aerosol analogs, called 'tholins', are presented. The Titan tholins were produced in the laboratory at Cornell by irradiation of simulated Titan atmospheres with high energy electrons in plasma discharge. Mass spectrometry measurements were performed at FOM of the solid phase of various tholins by Curie-point pyrolysis Gas-Chromatography/Mass-Spectrometry (GCMS) and by temperature resolved in-source Pyrolysis Mass-Spectrometry to reveal the composition and evolution temperature of the dissociation products. The results presented here are used to further define the ACP (Aerosol Collector Pyrolyser)-GCMS experiment and provide a basis for modelling of aerosol composition on Titan and for the interpretation of Titan atmosphere data from the Huygens probe in the future. Author (Herner)

Author (Herner)

A95-66798* National Aeronautics and Space Administration, Washington, DC.

ON THE LEVELS OF ENZYMATIC SUBSTRATE SPECIFICITY: IMPLICATIONS FOR THE EARLY EVOLUTION OF METABOLIC PATHWAYS

A. LAZCANO UNAM, Mexico, Mexico, E. DIAZ-VILLAGOMEZ UNAM, Mexico, Mexico, T. MILLS Univ. of Houston, Houston, TX, US, and J. ORO Univ. of Houston, Houston, TX, US *Life sciences and space research 24 (4): Planetary biology and origins of life;*

Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 *Advances in Space Research* (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 345-356
 Research sponsored by UNAM
 (Contract(s)/Grant(s): NAGW-2788)
 Copyright

The most frequently invoked explanation for the origin of metabolic pathways is the retrograde evolution hypothesis. In contrast, according to the so-called 'patchwork' theory, metabolism evolved by the recruitment of relatively inefficient small enzymes of broad specificity that could react with a wide range of chemically related substrates. In this paper it is argued that both sequence comparisons and experimental results on enzyme substrate specificity support the patchwork assembly theory. The available evidence supports previous suggestions that gene duplication events followed by a gradual neoDarwinian accumulation of mutations and other minute genetic changes lead to the narrowing and modification of enzyme function in at least some primordial metabolic pathways.

Author (Herner)

A95-66799

ROLE OF TRACE METAL IONS IN CHEMICAL EVOLUTION. THE CASE OF FREE-RADICAL REACTIONS

R. NAVARRO-GONZALEZ Universidad Nacional Autonoma de Mexico, Mexico, Mexico and C. PONNAMPERUMA Univ. of Maryland, College Park, MD, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 357-364*
 Copyright

We have studied the effect of iron in the free-radical oligomerization of hydrogen cyanide and acetic acid, and found that iron(II) and iron(III) readily reduces or oxidizes free radicals, respectively. The transient species produce by these reactions do not induce a chain oligomerization process and, therefore, they protect the solute molecules from degradation. Analysis of the available kinetic data for the reactions of a variety of transition metal ions with free radicals indicate that transition metal ions behave similarly to iron. Since Fe, Zn and Mo are essential to all living organisms, and there seems to be no apparent difference in chemical reactivity among transition metal ions towards free radicals, we suggest that these metal ions probably protected the biomolecules from degradation induced by free-radical reactions in the later stages of chemical evolution.

Author (Herner)

A95-66800

DIRECT INTERACTION BETWEEN AMINO ACIDS AND NUCLEOTIDES AS A POSSIBLE PHYSICO-CHEMICAL BASIS FOR THE ORIGIN OF THE GENETIC CODE

M. K. HOBISH Vectrix Corp., Baltimore, MD, US, N. S. M. D. WICKRAMASINGHE Univ. of Alabama at Birmingham, Birmingham, AL, US, and C. PONNAMPERUMA Univ. of Maryland, College Park, MD, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 365-382*
 Copyright

A study of the association of homocodonic amino acids and selected heterocodonic amino acids with selected nucleotides in aqueous solution was undertaken to examine a possible physical basis for the origin of codon assignments. These interactions were studied using ¹H nuclear magnetic resonance spectroscopy (NMR). Association constants for the various interactions were determined by fitting the changes in the chemical shifts of the anomeric and ring protons of the nucleoside moieties as a function of amino acid concentration to an isotherm which described the binding interac-

tion. The strongest association of all homocodonic amino acids were with their respective anticodonic nucleotide sequences. The strength of association was seen to increase with increase in the chain length of the anticodonic nucleotide. The association of these amino acids with different phosphate esters of nucleotides suggests that a definite isomeric structure is required for association with a specified amino acid; the 5'-mononucleotides and (3'-5')-linked dinucleotides are the favored geometries for strong associations. Use of heterocodonic amino acids and nonprotein amino acids supports these findings. We conclude that there is at least a physicochemical, anticodonic contribution to the origin of the genetic code.

Author (Herner)

A95-66802* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

VERY LOW TEMPERATURE FORMALDEHYDE REACTIONS AND THE BUILD-UP OF ORGANIC MOLECULES IN COMETS AND INTERSTELLAR ICES

W. A. SCHUTTE NASA. Ames Research Center, Moffett Field, CA, US, L. J. ALLAMANDOLA NASA. Ames Research Center, Moffett Field, CA, US, and S. A. SANDFORD NASA. Ames Research Center, Moffett Field, CA, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 401-406*
 Copyright

We have investigated thermally promoted reactions of formaldehyde (H₂CO) in very low temperature ices. No such reactions occurred in ices of pure formaldehyde. However, addition of trace amounts of ammonia (NH₃) were sufficient to catalyze reactions at temperatures as low as 40 K. Similar reactions could take place in interstellar ices and in Comets and produce considerable amounts of organic molecules.

Author (Herner)

A95-66804* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

POLYCYCLIC AROMATIC HYDROCARBON IONS AND THE DIFFUSE INTERSTELLAR BANDS

F. SALAMA NASA. Ames Research Center, Moffett Field, CA, US and L. J. ALLAMANDOLA NASA. Ames Research Center, Moffett Field, CA, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 413-422*
 Copyright

Neutral naphthalene (C₁₀H₈), phenanthrene (C₁₄H₁₀), and pyrene (C₁₆H₁₀) absorb strongly in the ultraviolet and may contribute to the extinction curve. High abundances are required to produce detectable structures. The cations of these Polycyclic Aromatic Hydrocarbons (PAHs) absorb in the visible. C₁₀H₈(+) has 12 discrete absorption bands which fall between 6800 and 5000 Å. The strongest band at 6741 Å falls close to the weak 6742 Å diffuse interstellar band (DIB). Five other weaker bands also match DIBs. The possibility that C₁₀H₈(+) is responsible for some of the DIBs can be tested by searching for new DIBs at 6520, 6151, and 5965 Å, other moderately strong naphthalene cation band positions. If C₁₀H₈(+) is indeed responsible for the 6742 Å feature, it accounts for 0.3% of the cosmic carbon. The spectrum of C₁₆H₁₀(+) is dominated by a strong band at 4435 Å in an Ar matrix and 4395 Å in a Ne matrix, a position which falls very close to the strongest DIB, that at 4430 Å. If C₁₆H₁₀(+), or a closely related pyrene-like ion is indeed responsible for the 4430 Å feature, it accounts for 0.2% of the cosmic carbon. We also report an intense, very broad UV-to-visible continuum which is associated with both ions and could explain how PAHs convert interstellar UV and visible radiation into IR.

Author (Herner)

A95-66805* National Aeronautics and Space Administration. Goddard Space Flight Center, Greenbelt, MD.
COSMIC DUST ANALOG SIMULATION IN A MICROGRAVITY ENVIRONMENT: THE STARDUST PROGRAM

F. FERGUSON Univ. of Virginia, Charlottesville, VA, US, L. U. LILLELEHT Univ. of Virginia, Charlottesville, VA, US, J. NUTH NASA. Goddard Space Flight Center, Greenbelt, MD, US, J. R. STEPHENS Matrix, Santa Fe, NM, US, E. BUSSOLETTI Istituto Universitario Navale, Cassino, Italy, L. CAROTENUTO MARS Center, Naples, Italy, L. COLANGELI Università di Cassino, Cassino, Italy, P. DELL'AVERSANA MARS Center, Naples, Italy, F. MELE MAR Center, Naples, Italy, V. MENNELLA Osservatorio Astronomico di Capodimonte, Naples, Italy et al. Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 423-426

Copyright

We have undertaken a project called STARDUST which is a collaboration with Italian and American investigators. The goals of this program are to study the condensation and coagulation of refractory materials from the vapor and to study the properties of the resulting grains as analogs to cosmic dust particles. To reduce thermal convective currents and to develop valuable experience in designing an experiment for the Gas-Grain Simulation Facility aboard Space Station, Freedom we have built and flown a new chamber to study these processes under periods of microgravity available on NASA's KC-135 Research Aircraft. Preliminary results from flights with magnesium and zinc are discussed.

Author (Herner)

A95-66807
ISOTOPE FRACTIONATIONS IN THE TERRESTRIAL CARBON CYCLE: A BRIEF OVERVIEW

M. SCHIDLowski Max-Planck-Institut fuer Chemie, Mainz, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 441-449

Copyright

The bias in favor of isotopically light carbon inherent in biological carbon fixation has brought about an isotopic disproportionation of primordial (mantle-derived) carbon on a global scale, causing an enrichment of C-12 in reduced (biogenic) carbon and a concomitant accumulation of the heavy complement (C-13) in the residual oxidized (inorganic) carbon pool. As a result, the terrestrial carbon cycle has gone bipartite, comprising an organic branch of isotopically light carbon, and an inorganic branch made up of C-13-enriched carbon (mostly in the form of carbonate). The isotopic disparity between the two principal terrestrial carbon species can be traced back over 3.8 Gyr of Earth history, attesting to a biological modulation of the carbon cycle since the time of formation of the oldest sediments.

Author (Herner)

A95-67426
INTELLIGENT MATERIALS PROPERTIES OF DNA AND STRATEGIES FOR ITS INCORPORATION INTO ELECTROACTIVE POLYMERIC THIN FILM SYSTEMS

KENNETH A. MARX Univ. of Massachusetts Lowell, Lowell, MA, JEONG OK LIM, DANIEL MINEHAN, RAJIV PANDE, MANJUNATH KAMATH, and SUKANT K. TRIPATHY Journal of Intelligent Material Systems and Structures (ISSN 1045-389X) vol. 5, no. 4 July 1994 p. 447-454 refs

(BTN-94-EIX94501433007) Copyright

We propose to create a novel class of intelligent materials by integrating two separate classes of intelligent materials - one biological and the other a thin film conducting polymer. The first class, DNA

possesses superior intelligent material properties designed over evolutionary time to function specifically and efficiently in integrated macromolecular arrays in cells called chromosomes. The second material is the polymeric thin film or two-dimensional Langmuir Blodgett (LB) monolayer film. In our approach, films will be comprised of electroactive alkylated conducting polymeric materials, such as polyalkylpyrrole and polyalkylthiophene, that are derivatized with biotin. Steptavidin conjugated DNA will be attached directly or biotinylated DNA will be stably attached to this film via a bridging streptavidin protein. To date, the bulk of our work has centered on characterizing the DNA binding to thick films of conducting polymers. A near term aim is to incorporate this signal transduction system into fiber optic biosensors for specifically detecting nucleic acid analyte. Our ultimate aim is to create novel ordered structures possessing unique integrated intelligent functions which respond to their environment and provide signal transduction approaches via their electronic and optical functions.

Author (EI)

A95-68775
THE USE OF COSMIC RAYS FOR CONTINUOUS MONITORING AND PREDICTION OF SOME DANGEROUS PHENOMENA FOR THE EARTH'S CIVILIZATION

L. I. DORMAN Inst. of Terrestrial Magnetism, Troitsk, Moscow, Russia, N. IUCCI Univ. 'La Sapienza', Rome, Italy, and G. VILLORESI Univ. 'La Sapienza', Rome, Italy Astrophysics and Space Science (ISSN 0004-640X) vol. 208, no. 1 October 1993 p. 55-68

(HTN-95-60524) Copyright

The main idea of the International Cosmic Ray Service (ICRS) is to combine satellite and spaceprobe cosmic rays, magnetic and plasma data with groundbased cosmic ray data (exchanged in real time) for obtaining continuous information on the electromagnetic and radiation situation in the interplanetary space and Earth's magnetosphere: prediction of great geomagnetic storms, big increases of radiation hazards and other dangerous phenomena in space and on the Earth for people and technology. ICRS can predict not only geomagnetic storms and unfavorable days in the environment (especially important for old people and people with some disease), but, in combination with astrophysical methods, can predict big increases of radiation hazards very dangerous for the Earth's civilization and big changes in the environment due to extremely powerful solar flares and local supernova explosions. We hope that, after some additional investigation of high energy cosmic-ray distribution function outside the heliosphere, it could be possible to solve by ICRS more complicated problems: to determine in combination with astrophysical methods the location and velocity of nearest dust-molecular galactic clouds with frozen-in magnetic fields and predict the expected time of the Sun capturing by some clouds with possible changes of Earth's global climate. The foundation of ICRS could bring a new possibility of development to the cosmic ray observatories, release scientists from a lot of routine work and increase the fundamental and applied research efficiency.

Author (Herner)

A95-68885
MYOSIN ISOFORMS IN MAMMALIAN SKELETAL MUSCLE

STEFANO SCHIAFFINO Padova Univ., Padua, Italy and CARLO REGGIANI Pavia Univ., Pavia, Italy Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 493-501 Research sponsored by Ministero dell'Università e delle Ricerche Scientifica e Tecnologica of Italy, Agenzia Spaziale Italiana, and Telethon-Italia

(HTN-95-A0102) Copyright

Skeletal muscles of different mammalian species contain four major myosin heavy-chain (MHC) isoforms: the 'slow' or beta-MHC and the three 'fast' Ila-, IIX-, and IIb-MHCs; and three major myosin light-chain (MLC) isoforms, the 'slow' MLC1s and the two 'fast' MLC1f and MLC3f. The differential distribution of the MHCs defines four major fiber types containing a single MHC isoform and a number of intermediate hybrid fiber populations containing both beta slow- and Ila-MHC, Ila- and IIX-MHC, or IIX- and IIb-MHC. The Ila-, IIX-,

51 LIFE SCIENCES (GENERAL)

and IIB-MHCs were first detected in neonatal muscles, and their expression in developing and adult muscle is regulated by neural, hormonal, and mechanical factors. The transcriptional mechanisms responsible for the fiber type-specific regulation of MHC and MLC gene expression are not known and are presently being explored by *in vivo* transfection experiments. The functional role of MHC isoforms has been in part clarified by correlated biochemical-physiological studies on single skinned fibers: these studies, in agreement with results from *in vitro* motility assays, indicate that both MHC and MLC isoforms determine the maximum velocity of shortening of skeletal muscle fibers.

Author (Hemer)

A95-68886

CAROTID BAROREFLEX CONTROL OF BLOOD PRESSURE AND HEART RATE IN MEN DURING DYNAMIC EXERCISE

Y. PAPELIER Paris-Sud Univ., Orsay, France, P. ESCOURROU Hopital Antoine Beclere, Clamart, France, J. P. GAUTHIER Centre National de la Recherche Scientifique, Gif-sur-Yvette, France, and L. B. ROWELL Washington Univ. School of Medicine, Seattle, WA, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 502-506 Research sponsored by Inst. de Recherches Internationales Servier (Contract(s)/Grant(s): NIH-HL-16910) (HTN-95-A0103) Copyright

The degree of control of blood pressure (BP) and heart rate (HR) by arterial baroreflex during exercise is still controversial. We studied baroreflex control of BP and HR in seven normal young men by a noninvasive procedure employing a neck suction chamber that delivers pulsatile positive and negative pressures to the carotid sinus (CS). Pressures applied to the CS ranged from -80 to +60 Torr in steps of 20 Torr. Pressure stimuli were triggered by electrocardiogram R wave, and each pressure step was maintained for 20 s in a continuous sequence. One baroreflex-response curve was obtained during the last 3 min of each 6-min period of exercise. The four levels of upright (cycle) exercise were 60, 120, 180, and 240 W, the highest requiring approximately 75% of maximal O₂ uptake. The sensitivity of the HR baroreflex response assessed by linear regression of HR vs. CS pressure (CSP) did not significantly decrease from rest (-0.09 ± 0.053 beat/Torr) to 240 W (-0.06 ± 0.025 beat/Torr). The BP above or below which CSP was increased or decreased by neck collar pressure was significantly increased from rest (76 ± 6.5 Torr) to 240 W (111.2 ± 4.0 Torr). The sensitivity of baroreflex response was assessed by linear regression of BP vs. CSP and was not significantly different from rest (-0.29 ± 0.054 Torr/Torr) up to exercise at 240 W (-0.29 ± 0.048 Torr/Torr). We conclude that mild to severe exercise does not reduce the gain of the CS reflex below resting values. CS function is upregulated by exercise so that at any CSP, systemic BP is higher and the change shift in the CS stimulus-response curve is increased as work rate increases.

Author (Hemer)

A95-68887

DEVELOPMENTAL DIFFERENCES IN VASCULAR RESPONSES TO HYPOXIA IN LUNGS OF RABBITS

CANDICE D. FIKE Utah Univ. Medical Center, Salt Lake City, UT, US and MARK R. KAPLOWITZ Utah Univ. Medical Center, Salt Lake City, UT, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 507-516 (Contract(s)/Grant(s): NIH-R-29-HL-42883) (HTN-95-A0104) Copyright

Our purpose was to determine whether postnatal age and prostaglandins influence the sites of hypoxic vasoconstriction in lungs of rabbits. To do this, we used the micropuncture technique to measure pressures in 20- to 80-micrometer-diam subpleural arterioles and venules during sequential inflation of lungs of newborn and adult rabbits with normoxic (21% O₂-7 - 10% CO₂-69 - 72% N₂) and hypoxic (90 - 93% N₂-7 - 10% CO₂) gas mixtures. Indomethacin (40 microgram/ml) was added to the perfusate of some lungs of each age group. During hypoxia in untreated lungs of newborn rabbits, both pulmonary arterial and 20- to 80-micrometer-diam arteriolar pressure increased by 5% whereas 20- to 80-micrometer-diam venular pressure remained the same. In contrast, during hypoxia in

untreated lungs of adult rabbits, pulmonary arterial pressure increased by 48%, whereas 20- to 80-micrometer-diam arteriolar pressure decreased slightly and 20- to 80-micrometer-diam venular pressure did not change. Regardless of the presence of indomethacin, location of vessels used for micropuncture, or level of left atrial pressure, pulmonary arterial pressure was the only measured vascular pressure that increased with hypoxia in adult lungs. Thus, in adult lungs, the site of hypoxia-induced vasoconstriction was limited to arteries greater than 80 micrometer diam, whereas in newborn lungs the site of hypoxia-induced vasoconstriction included vessels both larger and smaller than 20- to 80-micrometer-diam arteries. This age-related difference in the sites of hypoxia-induced vasoconstriction was not found in indomethacin-treated lungs.

Author (Hemer)

A95-68888

IMPAIRED PLASMA FFA OXIDATION IMPOSED BY EXTREME CHO DEFICIENCY IN CONTRACTING RAT SKELETAL MUSCLE

LORRAINE P. TURCOTTE Southern California Univ., Los Angeles, CA, US, PETER J. L. HESPEL Catholic Univ. of Leuven, Louvain, Belgium, TERRY E. GRAHAM Guelph Univ., Ontario, Canada, and ERIK A. RICHTER Copenhagen Univ., Copenhagen, Denmark Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 517-525 Research sponsored by Danish Medical Research Council, Fonds de la Recherche en Santa du Quebec, Belgian National Fonds voor Wetenschappelijk Onderzoek, and Natural Sciences and Engineering Research Council of Canada (HTN-95-A0105) Copyright

The extent to which carbohydrate (CHO) availability affects free fatty acid (FFA) metabolism in contracting skeletal muscle is not well characterized. To study this question, rats were depleted of glycogen by swimming exercise and lard feeding 24 h before perfusion of their isolated hindquarters. After 20 min of preperfusion with a medium containing no glucose, palmitate (600 or 2,000 micro M), and (1 - C-14) palmitate, flow was restricted to one hindlimb, which was electrically stimulated for 2 min to further deplete muscles of glycogen. After 2 min of recovery, glucose was added to the perfusate at final concentrations of 0, 6, or 20 mM, and after another 3 min muscles were stimulated for 30 min. At 6 and 2,000 micro M palmitate, glucose uptake after 30 min of stimulation averaged 23.5 ± 9.3 and 45.9 ± 10.6 micromol/g/h with 6 and 20 mM glucose, respectively. At 6 and 2,000 micro M palmitate, palmitate uptake was lower (30 - 37%, P less than 0.05) with 0 than with 6 or 20 mM glucose. At 600 micro M palmitate, percent palmitate oxidation was higher (27%, P less than 0.05) with 0 than with 6 or 20 mM glucose, resulting in similar total palmitate oxidation with the three glucose concentrations. At 2,000 micro M palmitate, percent palmitate oxidation was not significantly different among glucose concentrations, resulting in a significantly lower rate of palmitate oxidation with 0 than with 6 or 20 mM glucose. Muscle levels of malate and citrate and ammonia release from contracting muscle were not significantly different among glucose concentrations. It is concluded that, at a constant FFA concentration, contracting skeletal muscle does compensate for low CHO availability by increasing uptake and oxidation of FFA. At high FFA concentration, low CHO availability is accompanied by a decrease in FFA oxidation that is not associated with a decrease in the level of tricarboxylic acid cycle intermediates.

Author (Hemer)

A95-68889

NITRIC OXIDE-INDEPENDENT RESPONSE TO ACETYLCHOLINE BY TERMINAL ARTERIOLES IN RAT CREMASTER MUSCLE

ERIC VICAUT Hopital Fernand Widal, Paris, France, NATHALIE BAUDRY Hopital Fernand Widal, Paris, France, and XIN HOU Hopital Fernand Widal, Paris, France Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 526-533 Research sponsored by Paris VII Univ. and ASRM (HTN-95-A0106) Copyright

The aim of the present study was to establish whether, in terminal arterioles from the rat cremaster, acetylcholine (ACh) elicits nitric oxide (NO)-independent dilation corresponding to the transient

ACh-induced endothelium-dependent hyperpolarization described in arteries. For this purpose, The responses of terminal arterioles (mean diam 15.0 +/- 0.4 (SE) micrometers) were studied by intravital microscopy in rat cremaster muscle. During 15 min of superfusion by 10(exp -5) M ACh, the response was characterized by an initial maximal dilation (peak time less than 3 min) followed by a more sustained dilation that slightly decreased with time. Inhibition of NO synthesis by 2 x 10(exp -4) M N(sup omega)-nitro-L-arginine (L-NNA) significantly reduced, but did not eliminate, both the peak and sustained responses. Simultaneous administration of 2 x 10(exp -4) M L-NNA and 2 x 10(exp -5) M mefenamic acid, an inhibitor of prostaglandin synthesis, did not induce a significantly different response from that observed with L-NNA alone. Procaine (10 (exp -3) M), which is known to inhibit completely ACh-induced hyperpolarization in carotid artery, drastically reduced the initial part of the ACh-induced dilation but not the sustained response. Simultaneous administration of procaine and L-NNA almost completely inhibited the peak response to ACh. Similar results were obtained when L-NNA was combined with a superfusion bath containing 20 mM KCl, a concentration known to reduce hyperpolarization in arteries. Simultaneous administration of L-NNA and 10(exp -5) M ouabain, which inhibits Na(+)-K(+); 3 x 10(exp -6) M glibenclamide, which inhibits ATP sensitive K(+) channels; or 10(exp -8) M iberitoxin, which inhibits Ca(2+)-dependent K(+) channels, did not cause any additional inhibition to that resulting from L-NNA alone. We conclude that the ACh-induced dilation in terminal arterioles simultaneously involves NO- dependent and -independent mechanisms. The latter displayed characteristics very similar to those demonstrated for ACh-induced hyperpolarization in arteries. They were transient, chiefly occurred during the initial part of the response, and were abolished by procaine. In terminal arterioles, these NO-independent mechanisms were not mediated by either activation of the Na(+)-K(+) adenosinetriphosphatase or ATP-sensitive or Ca(2+)-dependent K(+) channels. +)

Author (Herner)

A95-68890

EFFECTS OF ACUTE RUNNING EXERCISE ON WHOLE BODY INSULIN ACTION IN OBESE MALE SHHF/MCC-FA(CP) RATS

J. GAO Ohio State Univ., Columbus, OH, US, W. M. SHERMAN, Ohio State Univ., Columbus, OH, US, S. A. MCCUNE Ohio State Univ., Columbus, OH, US, and K. OSEI Ohio State Univ., Columbus, OH, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 534-541 (HTN-95-A0107) Copyright

This study utilized the obese male spontaneously hypersensitive heart failure rat (SHHF/Mcc-fa(cp)), which has metabolic features very similar to human non-insulin-dependent diabetes mellitus. The purpose of this study was to assess the insulin sensitivity and responsiveness of whole body glucose disposal and insulin suppressability of hepatic glucose production with use of the euglycemic-hyperinsulinemic clamp procedure in 12- to 15-wk-old SHHF/Mcc-fa(cp) rats at rest (OS) and 2.5 h after a single session of acute exercise (OE). Lean male SHHF/Mcc-fa(cp) rats were sedentary (LS) control animals. At least three clamps producing different insulin-simulated responses were performed on each animal in a randomized order. At this age the obese animals are normotensive and have not developed congestive heart failure. Compared with LS, OS were significantly hyperglycemic and hyperinsulinemic and insulin sensitivity and responsiveness of whole body glucose uptake and insulin suppressability of hepatic glucose production were significantly decreased. Compared with LS and OS, acute exercise significantly decreased resting plasma glucose but did not alter plasma insulin. Compared with OS, acute exercise significantly increased the insulin responsiveness of whole body glucose disposal but did not affect the sensitivity of whole body glucose or insulin suppressability of hepatic glucose production. Compared with LS, however, acute exercise did not 'normalize' the insulin responsiveness of whole body glucose disposal. Thus a single acute exercise session improves but does not normalize whole body insulin resistance in the SHHF/MCC-fa(cp) rat.

Author (Herner)

A95-68891

TISSUE OXYGEN AND CARBON DIOXIDE STORES AND BREATH-HOLD DIVING IN HUMANS

MATS H. LINER Karolinska Inst., Stockholm, Sweden and DAG LINNARSSON Karolinska Inst., Stockholm, Sweden Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 542-547 Research sponsored by Swedish Medical Research Council and Swedish Space Board (HTN-95-A0108) Copyright

Alveolar gas exchange was studied in 11 submerged subjects during and after 75-s breath holds with or without a transient increase of ambient pressure to 3 ATA (20 msw). During surface breath holds (SBH) cardiac index fell to 73% of eupneic control but was partially restored at depth to 88% of control during breath-hold dives (BHD). O₂ uptake fell to 84% of control during SBH and was restored to control level during BHD. The turnover of O₂ stores was much slower during SBH than during the ensuing recovery. Carbon dioxide store dynamics were markedly slowed after BHD. We conclude that SBH and BHD are associated with large shifts in tissue O₂ and CO₂ stores and that much of these shifts in turnover rate for tissue O₂ and CO₂ stores could not be explained by the cardiac index changes alone but were compatible with peripheralization of venous blood volume and preferential peripheral vasoconstriction induced by apnea with elevated intrathoracic pressure during SBH. The transient compression during BHD reversed these central and peripheral circulatory changes by counteracting the increase in intrathoracic pressure.

Author (Herner)

A95-68892

HYPERTHERMIA STIMULATES NITRIC OXIDE FORMATION: ELECTRON PARAMAGNETIC RESONANCE DETECTION OF (DOT)NO-HEME IN BLOOD

DAVID M. HALL Iowa Univ., Iowa City, IA, US, GARRY R. BUETTNER Iowa Univ., Iowa City, IA, US, RONALD D. MATTHES Iowa Univ., Iowa City, IA, US, and CARL V. GISOLFI Iowa Univ., Iowa City, IA, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 548-553 (HTN-95-A0109) Copyright

Previous experiments from our laboratory have demonstrated that severe hyperthermia results in a selective loss of splanchnic vasoconstriction. Using electron paramagnetic resonance spectroscopy to scan whole blood samples collected in vivo from the portal vein and femoral artery of conscious unrestrained rats, we observed an increase in the concentration of spectroscopy-detectable species in portal venous blood of all heat-stressed animals. These spectra consisted of at least three distinct species: one with a broad feature having an effective g factor for the unpaired electron (g) of 2.06 assigned to the copper-binding acute phase protein ceruloplasmin, and two with narrower features that evolved at core temperature greater than 39 C representing a semiquinone radical and (dot)NO-heme. This heat-induced signal displays the classic nitrogen triplet hyperfine structure (nitrogen hyperfine splitting constant = 17.5 gauss, centered at g = 2.012) that is consistent with a five-coordinate heme complex and is characteristic of an unpaired electron coupled to nitrogen in the ferrous (dot)NO-heme adduct ((alpha(2+)NO)beta(3+))(sub 2). The intensity of this signal increased approximately twofold as core temperature rose to greater than 39 C, peaking 1 h post-heat exposure at greater than threefold basal concentration. This species was not seen in corresponding arterial blood samples. This is the first demonstration that whole body hyperthermia produces increased concentrations of radicals and metal binding proteins in the venous blood of the rat and suggests that severe hyperthermia stimulates an enhanced local release of (dot)NO within the splanchnic circulation.

Author (Herner)

A95-68893

SERIAL DISTRIBUTION OF AIRWAY MECHANICAL PROPERTIES IN DOGS: EFFECTS OF HISTAMINE

ROBERT H. HABIB, BELA SUKI, JASON H. T. BATES, and ANDREW C. JACKSON Journal of Applied Physiology (ISSN 8750-

7587) vol. 77, no. 2 August 1994 p. 554-566 Research sponsored by Whitaker Foundation, Medical Research Council of Canada, J. T. Costello Memorial Research Fund, and Respiratory Health Network of Centres of Excellence (Contract(s)/Grant(s): NIH-HL-31248) (HTN-95-A0110) Copyright

We measured respiratory input impedance (Z_{in} ; 8 - 2,048 Hz) in five dogs (anesthetized, tracheostomized, vagotomized, and mechanically ventilated) during 80 s of apnea after a bolus intravenous injection of saline or histamine (5.0 mg). In the control case, three antiresonances in Z_{in} were found in four of the dogs, whereas in the remaining dog only two were found. The magnitude and frequency of these antiresonances were significantly altered after bronchoconstriction. To interpret Z_{in} , a model incorporating detailed airway geometry, asymmetrical branching, and nonrigid airway walls was developed. The model fit both the saline and histamine Z_{in} data well and predicted a serial distribution of bronchoconstriction consistent with known effects of histamine; i.e., the diameters of the most peripheral airways were reduced (26% of their control values), whereas tracheal diameters were not significantly affected. The model provided estimates of tracheal diameters that were well correlated ($r = 0.92$) with direct measurements. Control estimates of soft tissue viscosity and Young's modulus compared closely with values in the literature. These results indicate that bronchoconstriction induced by histamine results in significant changes in Z_{in} over this frequency range and that by using this data analysis approach definitive physiological parameters relative to airway geometry and wall mechanical properties can be obtained from measurements made at the airway opening. Author (Herner)

A95-68894

REGIONAL LUNG HEMATOCRIT VARIATION AND ASSESSMENT OF ACUTE LUNG INJURY

MINORU KANAZAWA Keio Univ., Tokyo, Japan, NOKI HASEGAWA Keio Univ., Tokyo, Japan, TESTUYA URANO Keio Univ., Tokyo, Japan, KOICHI SAYAMA Keio Univ., Tokyo, Japan, SADATOMO TASAKA Keio Univ., Tokyo, Japan, FUMIO SAKAMAKI Keio Univ., Tokyo, Japan, HIDETOSHI NAKAMURA Keio Univ., Tokyo, Japan, YASUHIRO WAKI Keio Univ., Tokyo, Japan, TAKESHI TERASHIMA Keio Univ., Tokyo, Japan, SEITARO FUJISHIMA Keio Univ., Tokyo, Japan et al. Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 567-573 (HTN-95-A0111) Copyright

Estimating blood content in the lung remains a key step in calculating lung water volume and microvascular permeability. We studied the effect of regional lung hematocrit (Hct) variation on assessment of acute lung injury. *Escherichia coli* endotoxin was administered in guinea pigs intravenously. Lung injury was evaluated by measuring the wet-to-dry weight ratio (W/D) and transvascular I-125-labeled albumin leakage for 3 h (tissue-to-plasma I-125-albumin ratio (T/P)) in five tissue samples from each animal. Residual blood content was corrected using either Cr-51-red blood cells as a blood cell marker, (99m)Tc-albumin as a plasma marker, or both, injected 10 min before the guinea pigs were killed. Lung Hct, estimated from the marker counts of lung and peripheral blood samples, was lower than peripheral blood Hct; intraindividual variation, represented by the standard deviation in each subject, was 0.024 ± 0.015 for the control group (coefficient of variation $8.0 \pm 5.1\%$) and 0.026 ± 0.013 for the endotoxin group (coefficient of variation $8.5 \pm 4.1\%$). Uncorrected W/D for residual blood content was greater than the corrected W/D. (99m)Tc-albumin correction gave values closer to the W/D corrected by both markers. T/P corrected by (99m)Tc-albumin showed smaller data variations than the values obtained with Cr-51-red blood cell correction, which was affected by variations in lung Hct. We recommend using a plasma marker to correct for blood content in assessing acute lung injury by W/D and T/P. Author (Herner)

A95-68895

LONGITUDINAL DISTRIBUTION OF OZONE ABSORPTION IN THE LUNG: EFFECTS OF RESPIRATORY FLOW

SHU-CHIEH HU Pennsylvania State Univ., University Park, PA, US, ABDELLAZIZ BEN-JEBRIA Pennsylvania State Univ., University Park, PA, US, and JAMES S. ULTMAN Pennsylvania State Univ., University Park, PA, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 574-583 (Contract(s)/Grant(s): EPA-X-812059; NIH-ES-06075) (HTN-95-A0112) Copyright

In our previous work, we developed a bolus inhalation apparatus and measured the longitudinal distribution of ozone (O_3) uptake in intact human lungs at a quiet respiratory flow of 250 ml/min. The objective of the present study was to determine the effect of alternative respiratory flows between 150 and 1,000 ml/s. Uptake was expressed as the O_3 absorbed during a single breath relative to the amount of O_3 in the inhaled bolus Λ . Measurements of Λ were correlated with the penetration volume of the bolus into the respiratory tract (V_p). V_p in the range of 20 - 70 ml was considered to indicate upper airways (UA), the V_p interval of 70 - 180 ml was identified as lower conducting airways (CA), and V_p greater than 180 ml was associated with the respiratory air spaces (RA). During quiet oral breathing at 250 ml/s, Λ increased smoothly as V_p increased, with 50% of the inhaled O_3 absorbed in the UAs and the remainder absorbed within the CAs such that no O_3 reached the RAs. The effect of increasing the respiratory flow was to shift the Λ - V_p distribution distally such that significantly less O_3 was absorbed in the UAs and CAs and some O_3 reached the RAs. For example, at 1,000 ml/s, only 10% of the inhaled O_3 was absorbed in UAs and 65% was absorbed in the CAs such that 25% reached the RAs. Further analysis of these data with a mathematical diffusion model suggested that the reaction between O_3 and biochemical substrates is so fast (i.e., estimated 1st-order rate constant was on the order of $10(\exp 6)/s$) that O_3 is decomposed close to the gas-mucus interface in the CAs. Author (Herner)

A95-68896* National Aeronautics and Space Administration, Washington, DC.

INTERLEUKIN-2 THERAPY REVERSES SOME IMMUNOSUPPRESSIVE EFFECTS OF SKELETAL UNLOADING

JASON W. ARMSTRONG Kansas State Univ., Manhattan, KS, US, SIGNE BALCH Kansas State Univ., Manhattan, KS, US, and STEPHEN K. CHAPES Kansas State Univ., Manhattan, KS, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 584-589 (Contract(s)/Grant(s): NAGW-1197; NAGW-2328; DAMD17-89-Z-9039) (HTN-95-A0113) Copyright

Using antiorthostatic suspension, we characterized hematopoietic changes that may be responsible for the detrimental effect of skeletal unloading on macrophage development. Skeletally unloaded mice had suppressed macrophage development in unloaded and loaded bones, which indicated a systemic effect. Bone marrow cells from unloaded mice secreted less macrophage colony-stimulating factor and interleukin-6 than control mice. Additionally, T-lymphocyte proliferation was reduced after skeletal unloading. We show that polyethylene glycol-interleukin-2 therapy reversed the effects of skeletal unloading on macrophage development and cell proliferation. Author (Herner)

A95-68897

SURFACE ACTIVITY OF RABBIT PULMONARY SURFACTANT SUBFRACTIONS AT DIFFERENT CONCENTRATIONS IN A CAPTIVE BUBBLE

G. PUTZ California Univ., San Francisco, CA, US, J. GOERKE California Univ., San Francisco, CA, US, and J. A. CLEMENTS California Univ., San Francisco, CA, US Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 597-605 Research sponsored by Erwin-Schroedinger-Auslandsstipendium (Contract(s)/Grant(s): NIH-HL-24075; NIH-HL-40666) (HTN-95-A0114) Copyright

This study investigates the surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a new pressure-driven captive bubble surfactometer, which provides more

rigorous testing conditions than heretofore applied to this material. Subfractions were separated by centrifugation of lavage return into a third (P3; 1,000 x average g, 20min), a fourth (P4; 60,000 x average g, 60 min), and a fifth (P5; 100,000 x average g, 16 h) pellet. At 2.0 mg phospholipid/ml, P3 and P4 demonstrated more rapid adsorption, lower minimum surface tensions on first and subsequent compressions, and lower monolayer compressibilities than did P5. This surface activity of P3 and P4 increased with concentration between 0.1 and 2.0 mg phospholipid/ml. Adsorption rate constants were approximately 10,000 times faster than desorption constants. We conclude that, in a normal lung, components of P3 and P4 determine alveolar surface tension. We speculate that under special circumstances even less surface active P5 could have an important influence at the air-water interface. Author (Herner)

A95-68898

GENDER-RELATED DIFFERENCES IN THE BARORECEPTOR REFLEX CONTROL OF HEART RATE IN NORMOTENSIVE HUMANS

A. R. A. ABDEL-RAHMAN East Carolina Univ., Greenville, NC, US, R. H. MERRILL East Carolina Univ., Greenville, NC, US, and W. R. WOOLLES East Carolina Univ., Greenville, NC, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 606-613 (HTN-95-A0115) Copyright

The present study investigates the baroreceptor reflex control of heart rate (HR) of normotensive male and female human volunteers under two conditions: bolus- and infusion-evoked elevations of blood pressure by intravenous administration of phenylephrine. Average age and blood pressure were similar in both sexes, but females had a significantly lower heart period (HP; higher HR). A major difference existed between the two sexes when the blood pressure was elevated by the bolus method. Females had a significantly (50%) smaller baroreflex sensitivity (regression coefficient), which inferred a gender-related difference in baroreceptor reflex control of HR. However, because a positive correlation existed between basal HP and baroreflex sensitivity, it was important to investigate whether this difference was related to the significantly lower basal HP in females. This possibility was ruled out because a similar difference still existed when the data were collected from another group of females who had basal HP values similar to those of males. This gender-related difference in baroreceptor reflex control of HR seems to depend on the pattern by which the pressor stimulus is evoked. The baroreceptor HP response to a slowly developing pressor response that was maintained at a steady-state level was very similar in both sexes. Because the HP response to abrupt (bolus-evoked) pressor stimuli mainly reflects the activity of the vagal component, our findings suggest that the cardiac vagal component seems to play a substantially smaller role in the baroreflex-mediated bradycardia in females. Author (Herner)

A95-68899

EFFECTS OF STRENGTH TRAINING ON TOTAL AND REGIONAL BODY COMPOSITION IN OLDER MEN

M. S. TREUTH Maryland Univ., College Park, MD, US, A. S. RYAN Maryland Univ., College Park, MD, US, R. E. PRATLEY Maryland Univ., College Park, MD, US, M. A. RUBIN Maryland Univ., College Park, MD, US, J. P. MILLER Maryland Univ., College Park, MD, US, B. J. NICKLAS Maryland Univ., College Park, MD, US, J. SORKIN Maryland Univ., College Park, MD, US, S. M. HARMAN Maryland Univ., College Park, MD, US, A. P. GOLDBERG Maryland Univ., College Park, MD, US, and B. F. HURLEY Maryland Univ., College Park, MD, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 614-620 (Contract(s)/Grant(s): NIH-PO1-AG-04402; NIH-RO1-AG-07660; NIH-KO8-AG-00494; NIH-MO1-RR-02719; NIH-IT32-AG-00219) (HTN-95-A0116) Copyright

The effects of a 16-wk strength-training program on total and regional body composition were assessed by dual-energy X-ray absorptiometry (DEXA), magnetic resonance imaging (MRI), and hydrodensitometry in 13 untrained healthy men. Nine additional men

served as inactive control. The strength-training program resulted in substantial increases in both upper and lower body strength. Total fat-free mass (FFM) increased by 2 kg, and total fat mass decreased by the same amount when measured by DEXA. When measured by hydrodensitometry, similar increases in FFM and decreases in fat mass were observed. When measured by DEXA, FFM was increased in the arms, legs, and trunk, whereas fat mass was reduced in the arms, legs, and trunk as a result of training. MRI analysis revealed significant increases in mid thigh muscle cross-sectional area and significant reductions in mid thigh subcutaneous fat. These changes in body composition were not associated with changes in serum concentration of growth hormone, insulin-like growth factor I, or testosterone. None of the measured variables changed significantly in the control subjects. Thus, strength training increases regional and total lean mass and decreases regional and total fat mass in middle-aged and older men. Author (revised by Herner)

A95-68900

INTERACTION OF THYROID HORMONE AND FUNCTIONAL OVERLOAD ON SKELETAL MUSCLE ISOMYOSIN EXPRESSION

STEVEN J. SWOAP California Univ., Irvine, CA, US, FADIA HADDAD California Univ., Irvine, CA, US, VINCENT J. CAIOZZO California Univ., Irvine, CA, US, ROBERT E. HERRICK California Univ., Irvine, CA, US, SAMUEL A. MCCUE California Univ., Irvine, CA, US, and KENNETH M. BALDWIN California Univ., Irvine, CA, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 621-629 (Contract(s)/Grant(s): NIH-AR-30346) (HTN-95-A0117) Copyright

This study examined the interaction of exogenous thyroid hormone 3,5,3'-triiodothyronine (T sub 3) and functional overload on skeletal muscle myosin heavy chain (MHC) expression, studied at both the protein and mRNA level of analysis. Animals were allocated to the following groups: (1) normal control, (2) overload control, (3) hyperthyroid control, and (4) hyperthyroid + overload. Overload of the rat plantaris was accomplished by surgical removal of its synergists (soleus and gastrocnemius), and the animals were made hyperthyroid by injections of T(sub 3) (350 microgram/kg every other day). After overload of 8 wk, muscle enlargement occurred by 53% for both overload groups. This was accompanied by a 330 and 82% increase in the relative content of type I and IIa MHC, respectively, and a corresponding decrease by 16 and 44% in type IIx and IIb MHC, respectively, in the overload control group. Changes in the relative and absolute content of mRNA for these MHCs paralleled the protein response. Exogenous T(sub 3) completely reversed the upregulation of type I MHC and the downregulation of type IIx associated with overload at both the protein and mRNA. However, T(sub 3) was only partially effective in blunting the downregulation of IIb MHC and the upregulation of IIa MHC (protein and mRNA) accompanying the overload. These data suggest the following: (1) T(sub 3) can override the overload-induced signal in upregulating type I MHC expression in fast muscle, (2) the faster IIx and IIb MHC pools are differentially regulated by T(sub 3) and mechanical loading, and (3) both T(sub 3) and mechanical loading likely exert their impact on MHC expression at the pretranslational level of regulation. Author (Herner)

A95-68901

SIMULATION OF CARDIOVASCULAR RESPONSE TO LOWER BODY NEGATIVE PRESSURE FROM 0 TO -40 MMHG

FRANCOIS M. MELCHIOR, R. SRINI SRINIVASAN, PHILIPPE H. THULLIER, and JEAN-MICHEL CLERE *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 630-640 (Contract(s)/Grant(s): DRET-90-1053J) (HTN-95-A0118) Copyright

This paper presents a mathematical model for simulation of the human cardiovascular response to lower body negative pressure (LBNP) up to -40 mmHg both under normal conditions and when arterial baroreflex sensitivity or leg blood capacity (LBC) is altered.

Development of the model assumes that the LBNP response could be explained solely on the basis of (1) blood volume redistribution, (2) left ventricular end-diastolic filling, (3) interaction between left ventricle and peripheral circulation, and (4) modulations of peripheral resistances and heart rate by arterial and cardiopulmonary baroreflexes. The model reproduced well experimental data obtained both under normal conditions and during complete autonomic blockade; thus it is validated for simulation of the cardiovascular response from 0 to -40 mmHg LBNP. We tested the ability of the model to simulate the changes in LBNP response due to a reduction in LBC. To assess these changes experimentally, six healthy men were subjected to LBNP of -15, -30, and -38 mmHg with and without wearing elastic compression stockings. Stockings significantly reduced LBC and attenuated the change in heart rate. The model accurately reproduced this result. The model is useful for assessing the influence of LBC or other parameters such as arterial baroreflex sensitivity in diminishing the orthostatic tolerance of humans after spaceflight, bed rest, or endurance training. Author (Herner)

A95-68902**EFFECT OF GLUCOSE INFUSION ON ENDURANCE PERFORMANCE AFTER BETA-ADRENOCEPTOR BLOCKER ADMINISTRATION**

VAN BAAK, M. A. Limburg Univ., Maastricht, Netherlands and J. M. V. MOOIJ Limburg Univ., Maastricht, Netherlands *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 641-646 (HTN-95-A0119) Copyright

To investigate the effect of glucose (Glc) infusion on endurance performance after beta-adrenoceptor blockade, eight healthy male volunteers performed four endurance cycle ergometer tests at 67% of maximal work load after 80 mg of propranolol (Pr) of placebo (Pl) were administered orally in combination with a continuous infusion of Glc (0.5 g/min) or saline (Sal). The order of the tests was randomized. Endurance times were 53 +/- 6 (SE), 64 +/- 7, 26 +/- 5, and 31 +/- 6 min after Pl + Sal, Pl + Glc, Pr + Sal, and Pr + Glc, respectively (P less than 0.001). Glc infusion increased endurance after Pl (P less than 0.05) but not after Pr. Glc infusion resulted in significantly higher plasma Glc concentrations during exercise compared with Sal infusion (P less than 0.001). Glc infusion had no effect on cardiorespiratory or other metabolic variables. Plasma ammonia concentration was increased during the Pr tests (P less than 0.001) but reached similar values at exhaustion in all tests and was not affected by Glc infusion. The maintenance of plasma Glc concentration during endurance exercise at or above preexercise levels did not improve the reduction of endurance performance after Pr, indicating that the availability of blood Glc is not a limiting factor in this process. Author (Herner)

A95-68903**CONTRIBUTION OF BODY COMPOSITION AND PHYSICAL ACTIVITY TO AGE-RELATED DECLINE IN PEAK DOT-VO2 IN MEN AND WOMEN**

MICHAEL J. TOTH, ANDREW W. GARDNER, PHILIP A. ADES, and ERIC T. POEHLMAN *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 647-652 (Contract(s)/Grant(s): NIH-AG-00219; NIH-AG-07857; NIH-KO4-AG-00564; NIH-AG-0556; NIH-KO8-AG-00426) (HTN-95-A0120) Copyright

We examined the contribution of variations in body composition and leisure time physical activity to the age-related decline in peak oxygen consumption (dot-VO2) in men and women. Healthy males 17 - 80 yr old (n = 378) and females 18 - 81 yr old (n = 224) were characterized for dot-VO2 from a treadmill test to exhaustion, fat-free mass and fat mass by underwater weighing, and leisure time physical activity. Peak dot-VO2 showed a greater absolute decline (P less than 0.05) with age in males (r = 0.70, slope = -0.034 l/min/yr; P less than 0.01) than in females (r = 0.78, slope = -0.028 l/min/yr; P less than 0.01). After statistically controlling for differences in fat-free mass and fat mass, the decline in peak dot-VO2 was diminished in both sexes, although a greater rate of decline per-

sisted in males (r = -0.47, slope = -0.016 l/min/yr; P less than 0.01) than in females (r = -0.39, slope = -0.009 l/min/yr; P less than 0.01). We found that the addition of leisure time physical activity (independent of body composition) to the regression model further attenuated the rate of decline in males (r = 0.40, slope = -0.013 l/min/yr; P less than 0.01) but did not alter the age-related decline in peak dot-VO2 in females (r = -0.39, slope = -0.009 l/min/yr; P less than 0.01). We conclude that (1) the loss of fat-free mass and the increase in adiposity contribute to the decline in peak dot-VO2 with age in men and women and (2) the decline in leisure time physical activity, independent of differences in body composition is associated with the age-related decline in peak dot-VO2 in males but not in females. Author (Herner)

A95-68904**IMPACT OF PARALLEL HETEROGENEITY ON A CONTINUUM MODEL OF THE PULMONARY ARTERIAL TREE**

GARY S. KRENZ, JIANMING LIN, CHRISTOPHER A. DAWSON, and JOHN H. LINEHAN *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 660-670 Research sponsored by Dept. of Veterans Affairs (Contract(s)/Grant(s): NIH-HL-19298) (HTN-95-A0121) Copyright

Model arterial trees were constructed following rules consistent with morphometric data. Simulated heterogeneous trees approximating these rules were constructed by assigning vessel diameters $Hat-D(sub m) = D(sub a)(2/(m + 1))(exp 1/Beta(sub 1))$, such that $m - 1$ vessels were larger than $Hat-D(sub m)$ (vessel length proportional to diameter). Vessels were connected, forming random bifurcating trees. Longitudinal intravascular pressure ($Q(sub cum)$) with respect to cumulative vascular volume ($Q(sub cum)$) was computed for Poiseuille flow. Strahler-ordered tree morphometry yielded estimates of $L(sub a)$, $D(sub a)$, $Beta(sub 1)$, $Beta(sub 2)$, and mean number ratio(B); B is defined by $N(sub K + 1) = B(sub K)$, where k is total number of Strahler orders minus Strahler order number. The resulting $P(Q(sub cum))$ relationship was compared with that of the simulated tree, where Pa is total arterial pressure drop, Q is flow rate, $R(sub a) = (128/\mu L(sub a))/(\pi D(sub a)^4)$ (where μ is blood viscosity), and $Q(sub a)$ (volume of inlet artery) = $(1/4)D(sub a)^2/p/L(sub a)$. Results indicate that the equation, originally developed for homogeneous trees (*J. Appl. Physiol.* 72: 2225-2237, 1992), provides a good approximation to the heterogeneous tree $P(Q(sub cum))$. Author (Herner)

A95-68905**EXERCISE O2 TRANSPORT MODEL ASSUMING ZERO CYTOCHROME PO2 AT DOT-VO2(SUB MAX)**

JOHN W. SEVERINGHAUS *California Univ., San Francisco, CA, US Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 671-678 (HTN-95-A0122) Copyright

An analogy is drawn between cytochrome aa(sub 3) function and a polarographic cathode at which the potential of -0.6 V captures all O2 diffusing to the surface, achieving maximal O2 consumption (dot-VO2(sub max)) by eliminating O2 backpressure and outward diffusion from the surface, defined herein as zero surface PO2. The relationship of O2 consumption (as % dot-VO2(sub max)) to muscle venous, myoglobin, and cytochrome PO2 is modeled assuming that cytochrome aa(sub 3) PO2 reaches zero at dot-VO2(sub max), incorporating published data on the profile of leg venous PO2, pH, and blood lactate vs. work. Equations describe hemoglobin and myoglobin O2 dissociation and the Bohr effect of acid on O2 unloading. The O2 gradient from capillary blood to cytochrome aa(sub 3) is assumed to be proportional to O2 consumption. The model suggests that (1) to extract 75% of the O2 from myoglobin at dot-VO2(sub max), myoglobin must lie 90% down the O2 gradient from capillary to cytochrome; (2) the Bohr effect adds 15 - 30% to dot-VO2(sub max) and keeps venous PO2 almost constant as work rises from 60 to 100% of dot-VO2(sub max); and (3) in steady heavy work, the rising arterial lactate may impede lactate excretion from

muscle, reduce anaerobic ATP generation, and shift the energy balance toward aerobic metabolism. The zero PO₂ hypothesis facilitates modeling and may be the key to understanding the physiological limitation of work. Author (Herner)

A95-68906

HIGHER EXERCISE PERFORMANCE AND LOWER DOT-VO₂(SUB MAX) IN TIBETAN THAN HAN RESIDENTS AT 4,700 M ALTITUDE

RI-LI GE, QIU-HONG CHEN, LI-HAU WANG, DING GEN, PING YANG, KEISHI KUBO, KEISAKU FUJIMOTO, YUKINORI MATSUZAWA, KAZUHIKO YOSHIMURA, MICHIKO TAKEOKA et al. *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 684-691 (HTN-95-A0123) Copyright

To examine the hypothesis that the pathway of adaptation to high altitude in natives differs considerably from that in newcomers, we measured maximal O₂ uptake (dot-VO₂(sub max)), minute ventilation, anaerobic threshold (AT), blood lactate, and blood gases during maximal exercise in 17 lifelong Tibetan residents and 14 acclimatized Han Chinese newcomers living at the high altitude of 4,700 m. The two groups were similar in age, height, and weight, and the subjects were nonathletes. Although dot-VO₂(sub max) was significantly lower in the Tibetans than in the Hans, at maximal exercise effort the exercise workload was greater. The mean AT values (in % dot-VO₂(sub max)) in the Tibetan and Han subjects were 84.1 and 61.6%, respectively. Minute ventilation at maximal exercise was significantly lower in the Tibetans than in the Hans, whereas heart rate at maximal effort was equivalent in the two groups. The Tibetans showed lower blood lactate value than did the Hans both before and at the end of exercise. We conclude that the Tibetan natives have higher exercise performance and AT but lower dot-VO₂(sub max) and blood lactate concentration than do acclimatized Han newcomers. These results may reflect the effects of genetic or peripheral adaptation factors in the Tibetan natives.

Author (revised by Herner)

A95-68907

ROLES OF INTRA- AND EXTRACELLULAR CARBONIC ANHYDRASE IN ALVEOLAR-CAPILLARY CO₂ EQUILIBRATION

THOMAS A. HEMING, ERICH K. STABENAU, CARLOS G. VANOYE, HAMID MOGHADASI, and AKHIL BIDANI *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 697-705 Research sponsored by Moody Foundation (HTN-95-A0124) Copyright

Alveolar-capillary CO₂ equilibrium involves diffusive equilibration of CO₂ across the blood-gas barrier and chemical equilibration of perfusate CO₂-HCO₃⁻-H⁺ reactions. These processes are governed by different, but related, driving forces and conductances. The present study examined the importance of pulmonary carbonic anhydrase (CA) for diffusive and reactive CO₂ equilibration in isolated rat lungs. Lungs were perfused with salines containing membrane-impermeant or -permeant inhibitors of CA. Measurements of CO₂ excretion rate, equilibrated venous and arterial PCO₂ and pH, and postcapillary pH and PCO₂ disequilibria were used, together with our previous model of CO₂-HCO₃⁻-H⁺ reactions and transport in saline-perfused capillaries, to compute the relevant driving forces and conductances. Reactive CO₂ equilibration was markedly affected by extracellular (vascular) CA activity but not by the activity of intracellular (cytosolic) CA. The driving force for CO₂ diffusion was strongly influenced by vascular CA activity. The conductance for CO₂ diffusion was independent of CA activity. The minimum conductance for CO₂ diffusion was estimated to be 700 - 800 ml/min/Torr. The results indicate that extracellular vascular CA activity influences both diffusive and reactive CO₂ equilibration. However, cytosolic CA has no detectable role in alveolar-capillary CO₂ equilibration. Author (Herner)

A95-68908

VASCULAR IMPEDANCE ANALYSIS IN DOG LUNG WITH DETAILED MORPHOMETRIC AND ELASTICITY DATA

R. Z. GAN Memphis State Univ., Memphis, TN, US and R. T. YEN Memphis State Univ., Memphis, TN, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 706-717 (HTN-95-A0125) Copyright

On the basis of experimentally measured morphometric and elasticity data and model-derived mean pressure-flow conditions, we attempt a theoretical modeling of pulsatile flow in the whole lung. In the model we use the 'elastic tube' for arteries and veins, and the vascular impedance in arteries and veins follows Womersley's theory and electric analogue. We employ the 'sheet-flow' theory to describe the flow in the capillaries and to obtain the microvascular impedance matrix. The characteristic impedance of each order along the vascular tree, the input impedance at the capillary entrance and exit, and the pulmonary arterial input impedance at the main pulmonary artery are computed under certain physiological conditions. Using the pulsatile flow model, we investigate the effects of arterial vascular obstruction on pulmonary vascular impedance. The model-derived data are compared with the available experimental results in the literature. Author (Herner)

A95-68909

DIAMETERS OF JUXTACAPILLARY VENULES DETERMINED BY OIL-DROP METHOD IN RAT LUNG

RALPH SADURSKI St. Luke's-Roosevelt Hospital Center, Columbia Univ., New York, NY, US, HIROSHI TSUKADA St. Luke's-Roosevelt Hospital Center, Columbia Univ., New York, NY, US, XIAOYOU YING St. Luke's-Roosevelt Hospital Center, Columbia Univ., New York, NY, US, SUNITA BHATTACHARYA St. Luke's-Roosevelt Hospital Center, Columbia Univ., New York, NY, US, and JAHAR BHATTACHARYA St. Luke's-Roosevelt Hospital Center, Columbia Univ., New York, NY, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 718-725 Research sponsored by New York Lung Association (Contract(s)/Grant(s): NIH-HL-36024) (HTN-95-A0126) Copyright

We report a new method for precise quantification of lung microvascular diameter. Isolated blood-perfused rat lungs (500-g Sprague-Dawley rats) at constant inflation pressure (alveolar pressure (PA)) and stopped blood flow were viewed by microscopy and video. Subpleural venules of the second and third postcapillary generations were microinjected with oil colored with Sudan Black. Vascular pressure (Pvas) was varied in steps, and at each step the horizontal diameter (D(sub H)) and the length of the oil-filled segment were determined by microcaliper measurements of the replayed video image. At PA = 5 cmH₂O, a decrease in Pvas from 25 to 0 cmH₂O decreased D(sub H) in the second-generation venules from 55 +/- 2 (SE) to 41 +/- 1 micrometer (n = 13) and in the third-generation venules from 96 +/- 6 to 73 +/- 6 micrometer (n = 6). The constant-volume oil-filled segment conformed to the cylinder formula in that decreases in D(sub H) correlated linearly with 1/square root of length, thereby indicating that all Pvas values venular geometry was constant and probably circular and probably circular in cross section. The decrease in Pvas to -5 cmH₂O did not further decrease D(sub H). At Pvas = 10 - 25 cmH₂O, an increase in PA to 15 cmH₂O did not significantly increase D(sub H), although the increase in PA did diminish the slope (compliance) of the D(sub H)-Pvas relationship in second- but not third-generation venules. We conclude that (1) lung expansion decreases compliance of juxtacapillary venules, (2) venules retain circular cross sections at Pvas between -5 and 25 cmH₂O, and (3) venules are patent at subzero Pvas. Author (Herner)

A95-68910

HYPOXIA SIMILARLY IMPAIRS METABOLIC RESPONSES TO CUTANEOUS AND CORE COLD STIMULI IN CONSCIOUS RATS

GORDON G. GIESBRECHT Calgary Univ., Calgary, Canada, JAMES E. FEWELL Calgary Univ., Calgary, Canada, DAVID MEGIRIAN Calgary Univ., Calgary, Canada, ROLLIN BRANT Calgary Univ., Calgary, Canada, and JOHN E. REMMERS Calgary Univ., Calgary, Canada *Journal of Applied Physiology* (ISSN 8750-

7587) vol. 77, no. 2 August 1994 p. 726-730 Research sponsored by Medical Research Council of Canada, Manitoba Health Research Council, and Canadian Lung Association (Contract(s)/Grant(s): NIH-HL-42470) (HTN-95-A0127) Copyright

Cold exposure elicits several thermoregulatory responses, including an increased metabolic heat production from shivering and nonshivering thermogenesis. The increased metabolism can be in response to body core and/or body cutaneous cooling. Hypoxic hypoxia has been shown to attenuate the metabolic response to cutaneous cooling. We measured metabolic heat production in adult conscious rats during independent cutaneous and core cooling, during normoxia and hypoxia to (1) test the hypothesis that hypoxia suppresses the metabolic response to independent core cooling and (2) determine whether hypoxia acts preferentially on the response to cutaneous or core cooling. The animals were studied in a temperature-controlled metabolic chamber, and body core temperature was controlled by an abdominal heat exchange coil. Ambient temperature was varied (10, 19, and 28 C) while core temperature was clamped at 37 C or core temperature was varied (33, 35, and 37 C) at a stable ambient temperature of 28 C. Our data indicate that although the sensitivity of the metabolic response to core cooling is about five to six times that to cutaneous cooling, hypoxia similarly attenuates thermoregulatory responses to both stimuli.

Author (Herner)

A95-68911

INFLUENCE OF EXERCISE TRAINING ON MYOCARDIAL BETA-ADRENERGIC SIGNAL TRANSDUCTION: DIFFERENTIAL REGULATION WITH AGE

PHILIP J. SCARPACE, YE SHU, and NIHAL TUMER *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 737-741 Research sponsored by American Heart Association, Florida Affiliate and Dept. of Veterans Affairs (HTN-95-A0128) Copyright

Exercise training is known to increase cardiovascular performance and decrease heart rate. Because activation of adenylyl cyclase is an important factor in beta-adrenergic signal transduction and in the decline in signal transduction with age, we hypothesized that some of the effects of exercise training may be mediated by changes in postreceptor activation of adenylyl cyclase. To this end, we assessed isoproterenol-, G protein-, and forskolin-mediated activation of adenylyl cyclase as well as G protein immunoreactivity in the myocardium from young and senescent F-344 rats with and without prior exercise training by treadmill running. Isoproterenol, beta-gamma-imidoguanosine 5'-triphosphate, and forskolin stimulation of adenylyl cyclase activity declined by approximately 50% with age. Training increased the stimulation in the senescent rats and decreased the stimulation in the young rats such that the age-related decline in signal transduction was no longer significant. G(sub s) alpha protein immunoreactivity was unchanged by age or training. These data suggest that in young rats exercise training decreases beta-adrenergic signal transduction, whereas in older rats training increases signal transduction, partially offsetting the decline in signal transduction with age.

Author (Herner)

A95-68912

MAGNESIUM ATTENUATES PULMONARY HYPERTENSION DUE TO HYPOXIA AND GROUP B STREPTOCOCCI

MARK E. ANDERSON, TARA M. BURNETTE, DENNIS R. GEISER, and WARICHA JANJINDAMAI *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 751-756 Research sponsored by East Tennessee Foundation (HTN-95-A0129) Copyright

We investigated whether hypermagnesemia alleviates hypoxic or group B streptococcal (GBS) pulmonary hypertension (PH). Hypoxic PH was induced and maintained in 14 lambs by continuous ventilation with 12% oxygen. GBS PH was induced and maintained in 16 lambs by the continuous infusion of 5 - 10 x 10⁸ (exp 8) colony-forming units /kg/h of GBS. After the onset of PH, lambs were randomized to receive either magnesium sulfate (MgSO₄, intermit-

tent boluses of 0.38 mmol/kg, with a continuous infusion of 0.15 mmol/kg/h) or a similar volume of normal saline. Hypermagnesemia lowered pulmonary arterial pressure (PAP) and delayed the fall in systemic arterial pressure and stroke volume index seen in the control animals (each P less than 0.05). At a serum magnesium concentration (Mg) of 2.75 +/- 0.25 mmol/l, PAP was 27 +/- 3 compared with 40 +/- 4 Torr in the control animals (Mg = 0.87 +/- 0.06 mmol/l; P less than 0.05). In the GBS PH trial, hypermagnesemia prevented the continued increase in PAP seen in the control animals. At Mg = 2.15 +/- 0.07 mmol/l, PAP fell 2 +/- 1 Torr from prandomization values, whereas it rose 4 +/- 2 Torr in the control animals (Mg = 0.59 +/- 0.07 mmol/l; P less than 0.05). However, during the same time the systemic arterial pressure fell further in the magnesium-treated animals (-19 +/- 1 vs. -2 +/- 5 Torr). MgSO₄ attenuates PH in both models but may cause systemic hypotension in sepsis.

Author (Herner)

A95-68913

EFFECT OF HYPEROXIA AT 1 AND 2 ATA ON HYPOXIA AND HYPERCAPNIA IN HUMAN SKIN DURING EXPERIMENTAL INFLAMMATION

NEIL C. ABBOT, J. SWANSON BECK, FIONA M. T. CARNOCHAN, JAMES H. GIBBS, DAVID K. HARRISON, PHILIP B. JAMES, and J. GRAHAM LOWE *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 767-773 Research sponsored by Scottish Home and Health Dept. (HTN-95-A0130) Copyright

Transcutaneous PO₂ and PCO₂ measurements and estimates of skin respiration were monitored at different levels of inspired PO₂ in 20 healthy adults during the first 4 days of the tuberculin reaction, a convenient model of acute inflammation. Hyperoxia at 1 and 2 ATA significantly increased transcutaneous PO₂ levels in undisturbed and in inflamed skin but did not fully correct the relative hypoxia at the site of inflammation. Hypercapnia was reduced with O₂ breathing at 2 ATA. The apparent rate of O₂ consumption at the reaction site was raised during hyperoxia, most prominently at 2 ATA. The most intense reactions showed a central relative slowing of laser-Doppler blood flow indicative of microcirculatory impairment. The extent of the relative hypoxia and hypercapnia was greatest in these strongest reactions. The density of lymphocytes and monocytes in biopsies of 48-h reactions was loosely related to the corresponding transcutaneous PO₂ measurements. The present study provides evidence that diffusion barriers, in addition to increased local respiration, can contribute to the apparent hypoxia and hypercapnia of this inflammatory model.

Author (Herner)

A95-68914

LUNG VOLUME SPECIFICITY OF INSPIRATORY MUSCLE TRAINING

GEORGE E. TZELEPIS, DIEGO L. VEGA, MARK E. COHEN, and F. DENNIS MCCOOL *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 789-794 Research sponsored by Dept. of Veterans Affairs and Chicago Lung Association (HTN-95-A0131) Copyright

We examined the extent to which training-related increases of inspiratory muscle (IM) strength are limited to the lung volume (VL) at which the training occurs. IM strength training consisted of performing repeated static maximum inspiratory maneuvers. Three groups of normal volunteers performed these maneuvers at one of three lung volumes: residual volume (RV), relaxation volume (Vrel), or Vrel plus one-half of inspiratory capacity (Vrel + 1/2IC). A control group did not train. We constructed maximal inspiratory pressure-VL curves before and after a 6-wk training period. For each group, we found that the greatest improvements in strength occurred at the volume at which the subjects trained and were significantly greater for those who trained at low (36% for RV and 26% for Vrel) than at high volumes (13% for Vrel + 1/2IC). Smaller increments in strength were noted at volumes adjacent to the training volume. The range of vital capacity (VC) over which strength was increased was greater for those who trained at low (70% of VC) than at high VL (20% of VC).

We conclude that the greatest improvements in IM strength are specific to the VL at which training occurs. However, the increase in strength, as well as the range of volume over which strength increased, is greater for those who trained at the lower VL.

Author (Herner)

A95-68915

PRESSURE-FLOW SPECIFICITY OF INSPIRATORY MUSCLE TRAINING

GEORGE E. TZELEPIS, DIEGO L. VEGA, MARK E. COHEN, ASHOK M. FULAMBARKER, KISHOR K. PATEL, and F. DENNIS MCCOOL *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 795-801 Research sponsored by Dept. of Veterans Affairs and Chicago Lung Association (HTN-95-A0132) Copyright

The inspiratory muscles (IM) can be trained by having a subject breathe through inspiratory resistive loads or by use of unloaded hyperpnea. These disparate training protocols are characterized by high inspiratory pressure or high inspiratory flow, respectively. We tested the hypothesis that the posttraining improvements in IM pressure or flow performance are specific to training protocols in a way that is similar to force-velocity specificity of skeletal muscle training. IM training was accomplished in 15 normal subjects by use of three protocols: high inspiratory pressure-no flow (group A, n = 5), low-inspiratory pressure-high flow (group B, n = 5), and intermediate inspiratory pressure and flow (group C, n = 5). A control group (n = 4) did no training. Before and after training, we measured esophageal pressure (Pes) and inspiratory flow (dot-VI) during single maximal inspiratory efforts against a range of external resistances including an occluded airway. Efforts originated below relaxation volume (Vrel), and peak Pes and dot-VI were measured at Vrel. Isovolumic maximal Pes-dot-VI plots were constructed to assess maximal inspiratory pressure-flow performance. Group A (pressure training) performed 30 maximal static inspiratory maneuvers at Vrel daily, group B (flow training) performed 30 sets of three maximal inspiratory maneuvers with no added external resistance daily, and group C (intermediate training) performed 30 maximal inspiratory efforts on a midrange external resistance (7 mm ID) daily. Subjects trained 5 days/wk for 6 wk. Data analysis included comparison of posttraining Pes-dot-VI slopes among training groups. After training, group A increased maximal Pes (Pes(sub max); 37%) but not maximal dot-VI (dot-VI(sub max)) and group B increased dot-VI(sub max) (17%) but not Pes(sub max); group C increased Pes(sub max) (19%) and dot-VI(sub max) (14%). The posttraining slopes were dependent on the training protocol. We conclude that training protocols characterized by generating high inspiratory pressure or high inspiratory flow will specifically increase Pes(sub max) or dot-VI(sub max), respectively. In contrast, intermediate training protocols produce a more uniform increase in dot-VI(sub max) and flow.

Author (Herner)

A95-68916

SUPERNORMAL MUSCLE FIBER CONDUCTION VELOCITY DURING INTERMITTENT ISOMETRIC EXERCISE IN HUMAN MUSCLE

J. H. VAN DER HOEVEN *University Hospital Groningen, Groningen, Netherlands* and F. LANGE *University Hospital Groningen, Groningen, Netherlands* *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 802-806 (HTN-95-A0133) Copyright

Muscle fiber conduction velocity (MFCV) and surface electromyographic parameters were studied in the brachial biceps muscle of healthy males during voluntary intermittent isometric contractions at 50% of maximum force. Recovery in the following 15 min was then observed. The measurements were performed during duty cycles of 33, 25, and 10%. The main finding was a supernormal MFCV during the exercise phase when the duty cycle was 25 and 20%. The level continued to increase during the recovery phase. During the exercise phase when the duty cycle was 33%, the MFCV decreased slightly (suggesting that the local anaerobic threshold had been surpassed) but increased during recovery to supernormal

values. The ratio of median frequency to MFCV was constant during all experiments, indicating that the changes in median frequency reflect those in MFCV. We suggest that the supernormal MFCV was due to a combination of altered membrane properties, muscle fiber swelling, and temperature increase and hypothesize that the changes of electrical properties formed part of an adaptive mechanism of the muscle fiber membrane during exercise. In that respect, the increase of the MFCV could be a component of the well-known warm-up effect.

Author (Herner)

A95-68917

MAXIMUM RATE OF FORCE DEVELOPMENT IS INCREASED BY ANTAGONIST CONDITIONING CONTRACTION

MARK D. GRABINER *Cleveland Clinic Foundation, Cleveland, OH, US* *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 807-811 Research sponsored by Chattecx Corp.

(HTN-95-A0134) Copyright

Nine subjects performed maximum contractions with the knee extensor muscles on an isokinetic device set at 4.36 rad/s. The knee extensions were preceded by isometric conditioning contractions of the antagonist knee flexor muscles. The conditioning contraction forces were 0, 25, 50, 75, and 100% of the maximum isometric force. The purpose of the study was to determine (1) the effects of graded conditioning contractions with the knee flexor muscles on the maximum performance of the quadriceps femoris and (2) whether the effects of the conditioning contractions are similarly distributed to components of the quadriceps femoris. The increased initial quadriceps femoris activation levels associated with the conditioning contractions were similarly distributed to quadriceps femoris components. In contrast with previously published research on exercise performed at substantially lower knee extension velocities, the maximum knee extension force and the work performed were not affected by the conditioning contractions. However, the maximum rate of force development increased significantly (P less than 0.05). The intensity of the knee flexor conditioning contractions likely influences agonist motor unit activation variables, but this influence may be related to the speed of the agonist contraction.

Author (Herner)

A95-68918

DETECTION OF FREE RADICALS BY ELECTRON SPIN RESONANCE IN RAT DIAPHRAGM AFTER RESISTIVE LOADING

G. BORZONE *Ohio State Univ., Columbus, OH, US*, B. ZHAO *Ohio State Univ., Columbus, OH, US*, A. J. MEROLA *Ohio State Univ., Columbus, OH, US*, L. BERLINER *Ohio State Univ., Columbus, OH, US*, and T. L. CLANTON *Ohio State Univ., Columbus, OH, US* *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 812-818 Research sponsored by Sigma Xi Research Society

(Contract(s)/Grant(s): NIH-RR-03126)

(HTN-95-A0135) Copyright

Indirect evidence supports free radical production in the diaphragm under excessive mechanical loads in both in vitro and in situ preparations. We hypothesized that free radicals are produced in the diaphragm with loads in vivo at a sufficient concentration to be detected by electron spin resonance (ESR) spectroscopy. Anesthetized rats underwent severe inspiratory resistive loading for 2.5 - 3 h with maintenance of blood oxygenation and arterial blood pressure by breathing 70% oxygen. The ESR spectra of four samples (freeze-clamped at liquid nitrogen temperature) from each experimental animal were compared with the spectra from a control animal breathing air and a control animal breathing 70% oxygen. We observed (1) an approximately 30% increase in intensity of free radical signal in experimental animals (n = 10) compared with control animals breathing oxygen (n = 10; P less than 0.01) and control animals breathing air (n = 10; P less than 0.05), (2) that oxygen alone had no effect on the ESR spectrum, and (3) the intensity of the ESR signal decreased approximately 25% in the experimental group when samples were taken 10 min postmortem, whereas no differ-

51 LIFE SCIENCES (GENERAL)

ence in signal was observed for control animals. We conclude that the diaphragm shows an increased production of free radicals associated with respiratory failure induced by resistive breathing.

Author (Hemer)

A95-68919

EFFECTS OF HYPERINFLATION AND CPAP ON WORK OF BREATHING AND RESPIRATORY FAILURE IN DOGS

ELIZABETH SHADE Johns Hopkins Asthma and Allergy Center, Baltimore, MD, US, YASHUHIRO KAWAGOE Johns Hopkins Asthma and Allergy Center, Baltimore, MD, US, ROY G. BROWER Johns Hopkins Asthma and Allergy Center, Baltimore, MD, US, SOLBERT PERMUTT Johns Hopkins Asthma and Allergy Center, Baltimore, MD, US, and HENRY E. FESSLER Johns Hopkins Asthma and Allergy Center, Baltimore, MD, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 819-827

(HTN-95-A0136) Copyright

Increased end-expiratory lung volume (EELV) and airway resistance are both characteristic features of obstructive lung disease. Increased EELV alone loads the respiratory muscles and may cause respiratory failure, changes that could be reversed by continuous positive airway pressure (CPAP). To study the effects of elevated EELV on respiration without increased airway resistance, we used a mechanical analogue of airway closure to increase EELV in six spontaneously breathing anesthetized dogs. Hyperinflation of 0.84 +/- 0.11 liter for 30 min decreased minute ventilation from 4.8 +/- 0.37 to 3.5 +/- 0.21 l/min and increased arterial PCO₂ from 40.3 +/- 1.5 to 73.2 +/- 8.1 Torr (both P less than 0.01). Inspiratory work per breath increased 3-fold, work per liter increased 3.7-fold, and work per minute increased 2.8-fold (all P less than 0.01). CPAP at 15 cmH₂O restored minute ventilation to 4.3 +/- 0.3 l/min and reduced arterial PCO₂ to 54 +/- 6.6 Torr (NS vs. baseline). All measurements of inspiratory work were also restored to baseline, but cardiac output was reduced (baseline 3.09 +/- 0.36, hyperinflation 2.71 +/- 0.36, hyperinflation + CPAP 1.94 +/- 0.29 l/min; P less than 0.05, baseline vs. hyperinflation + CPAP). We conclude that increases in EELV mimic important features of airway obstruction, increase inspiratory work, and can cause respiratory failure independent of increased respiratory failure independent of increased airway resistance. This respiratory failure is reversed by CPAP at the potential expense of hemodynamic compromise.

Author (Hemer)

A95-68920

PERFORMANCE AND METABOLIC EFFECTS OF BENZODIAZEPINE DURING SUBMAXIMAL EXERCISE

K. COLLOMP, M. FORTIER, S. COOPER, A. LONG, S. AHMAIDI, C. PREFAUT, F. WRIGHT, M. PICOT, and M. G. COTE *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 828-833 Research sponsored by Canadian Center for Drug-free Sport

(HTN-95-A0137) Copyright

The present study examined whether benzodiazepine (BZ) intake alters performance and selected hormonal and metabolic variables during submaximal exercise. Seven triathletes completed two cycling trials at 85% maximum O₂ uptake starting 3 h after an ingestion of either a placebo (PLA) of gelatin or BZ (1.5 mg lorazepam) and continuing until exhaustion, according to a double-blind randomized protocol. Blood samples were collected at rest; 5, 10, and 15 min; and exhaustion for dopamine (DA), norepinephrine (NE), epinephrine (Epi), adrenocorticotropic hormone (ACTH), cortisol (CORT), insulin (INS), free fatty acid, blood glucose, and lactate (La) determinations. Time of cycling was not significantly changed after BZ or PLA administration. A decrease in CORT and an increase in INS (P less than 0.05) were observed with BZ before cycling. In comparison with rest, exercise resulted in a decrease in INS and an increase in all the other variables investigated (P less than 0.001), but DA, NE, Epi, ACTH, CORT, La, and free fatty acid were significantly less elevated under BZ (P less than 0.05). No change was found in glucose and INS levels between the two treatments at the end of the test. There was a strong correlation under both PLA and BZ conditions between DA, NE, Epi, and ACTH and also

between Epi and La levels. From these data, BZ intake did appear to alter metabolism but did not influence performance during intense submaximal exercise.

Author (Hemer)

A95-68921

INFLUENCE OF LUNG VOLUME DEPENDENCE OF UPPER AIRWAY RESISTANCE DURING CONTINUOUS NEGATIVE AIRWAY PRESSURE

F. SERIES Centre de Pneumologie de l'Hopital Laval, Univ. Laval, Sainte Foy, Canada and I. MARC Centre de Pneumologie de l'Hopital Laval, Univ. Laval, Sainte Foy, Canada *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 840-844 Research sponsored by Respiratory Health Network of Centres of Excellence of Canada

(HTN-95-A0138) Copyright

To quantify the contribution of lung volume dependence of upper airway (UA) on continuous negative airway pressure (CNAP)-induced increase in upper airway resistance, we compared the changes in supralaryngeal resistance during an isolated decrease in lung volume and during CNAP in eight normal awake subjects. Inspiratory supralaryngeal resistance was measured at isoflow during four trials, during two CNAP trials where the pressure in a nasal mask was progressively decreased in 3- to 5-cmH₂O steps and during two continuous positive extrathoracic pressure (CPEP) trials where the pressure around the chest (in an iron lung) was increased in similar steps. The CNAP and CPEP trials were done in random order. During the CPEP trial, the neck was covered by a rigid collar to prevent compression by the cervical seal of the iron lung. In each subject, resistance progressively increased during the experiments. The increase was linearly correlated with the pressure increase in the iron lung and with the square of the mask pressure during CNAP. There was a highly significant correlation between the rate of rise in resistance between CNAP and CPEP: the steeper the increase in resistance with decreasing lung volume, the steeper the increase in resistance with decreasing airway pressure. Lung volume dependence in UA resistance can account for 61% of the CNAP-induced increase in resistance. We conclude that in normal awake subjects the changes in supralaryngeal resistance induced by CNAP can partly be explained by the lung volume dependence of this resistance.

Author (Hemer)

A95-68922

PULMONARY VASCULAR RESISTANCE DISTRIBUTION AND RECRUITMENT OF MICROVASCULAR SURFACE AREA

K. A. OVERHOLSER Vanderbilt Univ., Nashville, TN, US, NIOMI A. LOMANGINO Vanderbilt Univ., Nashville, TN, US, R. E. PARKER Vanderbilt Univ., Nashville, TN, US, N. A. POU Vanderbilt Univ., Nashville, TN, US, and T. R. HARRIS Vanderbilt Univ., Nashville, TN, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 845-855

(Contract(s)/Grant(s): NIH-HL-19153; NIH-HL-07411; NIH-RR-06558)

(HTN-95-A0139) Copyright

To test the hypothesis that the distribution of hemodynamic resistance is involved in the control of pulmonary capillary surface area, we measured permeability-surface area product (PS) and longitudinal resistance distribution (LRD) as functions of perfusion rate in isolated rabbit lungs under zone II conditions (n = 10) and through the zone II - III transition (n = 4). PS, considered to be indicative of functioning capillary surface area, was measured with the aid of the diffusion-limited tracer (C-14)propanediol, whereas LRD was determined using a viscous bolus technique. LRD was seen to change character with increasing flow and increasing PS/surface area, becoming bimodal with low central resistance as full capillary recruitment was approached in zone III. Effects of hypoxic ventilation were studied in zone II in five lungs; it was found that hypoxia altered the LRD and eradicated the normoxic dependence of PS/surface area on perfusion rate. It was concluded that LRD is involved in the determination of functioning capillary surface area.

Author (Hemer)

A95-68923**ENERGY BALANCE AT HIGH ALTITUDE OF 6,542 M**

KLAAS R. WESTERTERP, BENGT KAYSER, LOEK WOUTERS, JEAN-LOUIS LE TRONG, and JEAN-PAUL RICHALET *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 862-866 Research sponsored by Sandoz Nutrition and Groupe de Reflexion sur le Probleme de l'Eau dans les Environnements d'Exception (HTN-95-A0140) Copyright

Weight loss due to malnutrition and possibly intestinal malabsorption is a well-known phenomenon in high-altitude climbers. Up to approximately 5,000 m, energy balance may be attained and intestinal energy digestibility remains normal. To see whether (1) energy balance may also be attained at 6,542 m and, if not (2) whether decreased energy digestibility would play a significant role in the energy deficit, energy intake (EI), energy expenditure, body composition, and energy digestibility of 10 subjects (4 women, 6 men; 27 - 44 yr) were assessed during a 21-day sojourn on the summit of Mt. Sajama, Bolivia (6,542 m). EI was measured during two 3-day intervals: EI1 (days 7 - 9) and EI2 (days 17 - 19). Total fecal energy loss during EI1 was calculated from fecal energy measured by bomb calorimetry. Average daily metabolic rate (ADMR) at altitude was measured in six subjects (2 women, 4 men) using doubly labeled water over a 10-day interval (days 9 - 19). Basal metabolic rate was measured before and after the expedition by respiratory gas analysis. Body composition was estimated from skinfolds and body mass before and during the altitude sojourn. Subjects were in negative energy balance throughout the observation period. The activity level, expressed as ADMR to basal metabolic rate, was 1.56 - 2.39. The loss of fat mass (3.7 +/- 1.5 kg) represented 74 +/- 15% of the loss of body mass. Energy content of the feces was 21 kJ/g dry wt, and gross energy digestibility amounted to 85%. The energy deficit increased to 3.5 MJ/day after correction for the decreased energy digestibility. In conclusion, energy balance was not attained at 6,542 m. The resulting energy deficit appeared to result mostly from malnutrition, and only a limited part could be attributed to malabsorption. Author (Herner)

A95-68924**RESPIRATORY RESPONSE TO INHALED CO₂ DURING POSITIVE INSPIRATORY PRESSURE IN HUMANS**

P. SCHEID Hopital Henri Mondor, Creteil, France, F. LOFASO Hopital Henri Mondor, Creteil, France, D. ISABEY Hopital Henri Mondor, Creteil, France, and A. HARF Hopital Henri Mondor, Creteil, France *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 876-882 Research sponsored by Fondation pour la Recherche Medicale (Contract(s)/Grant(s): DFG-II-B-7-SCHE-46/7-1) (HTN-95-A0141) Copyright

To investigate ventilatory CO₂ sensitivity during inspiratory pressure support (IPS), we administered inspiratory CO₂ (fractional concn (FI(sub CO₂)) 0.01, 0.03, or 0.05) in eight normal subjects without (CTRL) or with (Pinsp) positive inspiratory airway pressure (5 or 10 cmH₂O). At CTRL and low IPS, CO₂ inhalation led to a significant increase in tidal volume (VT) with nearly identical slopes in the plot of VT vs. end-tidal PCO₂. At the high IPS level, VT at FI(sub CO₂) of 0 was significantly above the value at lower Pinsp and did not increase with CO₂ unless FI(sub CO₂) was elevated to greater than 0.03. There was very little effect of either Pinsp or FI(sub CO₂) on respiratory frequency and respiratory timing. The data suggest that the CO₂ sensitivity of ventilation is similar at low levels of IPS as during CTRL. However, at high levels of IPS, VT is determined largely by the passive inflation and, thus, independent of CO₂. CO₂ has to be elevated to increase the respiratory drive before VT becomes CO₂ sensitive. Author (Herner)

A95-68925**CHEST WALL INTERRUPTER RESISTANCE IN ANESTHETIZED PARALYZED HUMANS**

E. D'ANGELO, E. PRANDI, M. TAVOLA, E. CALDERINI, and J. MILIC-EMILI *Journal of Applied Physiology* (ISSN 8750-7587)

vol. 77, no. 2 August 1994 p. 883-887 Research sponsored by National Atlantic Treaty Organization-CNR Senior Guest Fellowship (HTN-95-A0142) Copyright

Tracheal (Ptr) and esophageal (Pes) pressure and flow were measured in 12 supine anesthetized paralyzed normal subjects aged 16-22 yr. The subjects were ventilated with a fixed inflation volume (range 0.57 - 0.62 liter) and with different constant flows ranging between 0.24 and 1.12 l/s. A rapid airway shutter (closing time 10 - 15 ms) was used to briefly occlude (0.4 - 0.9 s) the airways at end inspiration for 33 - 44 consecutive breaths. At each flow level, Ptr and Pes records obtained during end-inspiratory occlusions were ensemble averaged to allow for the cardiac artifacts. The interrupter resistances of the chest wall and respiratory system were assessed as the rapid fall in Pes and Ptr with occlusion divided by the flow preceding the occlusion. Interrupter resistances of both the chest wall and lung were independent of flow and averaged 0.4 +/- 0.1 and 1.5 +/- 0.4 (SD) cmH₂O/s/l, respectively. The contribution of the chest wall to the total interrupter resistance was approximately 27% at flows less than or equal to 1 l/s. Author (Herner)

A95-68926**HYPOXIA INCREASES GLUCOSE TRANSPORT AT BLOOD-BRAIN BARRIER IN RATS**

SAMI I. HARIK, RAMIN A. BEHMAND, and JOSEPH C. LAMANNA *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 896-901 Research sponsored by Shriners-Burns and Fogarty International (Contract(s)/Grant(s): NIH-HL-36829; NIH-HL-07354) (HTN-95-A0143) Copyright

Prolonged hypoxia causes several adaptive changes in systemic physiology and tissue metabolism. We studied the effects of hypobaric hypoxia on glucose transport at the blood-brain barrier (BBB) in the rat. We found that hypoxia increased the density of brain microvessels seen on immunocytochemical stains using an antibody to the glucose transporting protein GLUT. In addition, we found that hypoxia increased the density of GLUT in isolated cerebral microvessels as determined by specific cytochalasin B binding. The higher GLUT density in isolated cerebral microvessels was evident after 1 wk of hypoxia and was associated with decreased activity of gamma-glutamyltranspeptidase. Consistent with these findings, we also demonstrated that 3 wk of hypobaric hypoxia caused increased unidirectional transport of glucose at the BBB in several brain regions in vivo, as determined by the doubly labeled single-pass indicator-fractionation atrial bolus injection method in anesthetized rats. We conclude that chronic hypobaric hypoxia is associated with increased glucose transport at the BBB. Author (Herner)

A95-68927**HYPERBARIC OXYGENATION INCREASES AROUSAL AND BREATHING MOVEMENTS IN FETAL LAMBS**

M. H. TIKTINSKY-RUPP, S. U. HASAN, B. BISHOP, and F. C. MORIN, III *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 902-911 Research sponsored by Alberta Lung Association and Alberta Children's Hospital (Contract(s)/Grant(s): NIH-RO1-HL-41387) (HTN-95-A0144) Copyright

Oxygenation produced by distending the lungs with 100% O₂ increases the occurrence of arousal and fetal breathing movements (FBM), particularly during non-rapid-eye-movement (NREM) sleep, in fetal sheep of equal to or greater than 35 days of gestation. We studied the breathing and behavioral responses in arterial PO₂ (Pa(sub O₂)) without lung distension in fetuses between 128 and 132 days gestation. Twelve fetuses were chronically instrumented to record FBM, behavioral state, blood pressure, arterial blood gas tensions, and pH. Fetal Pa(sub O₂) was raised by having the ewe breathe 100% O₂ at 3 atmosphere absolute pressure spontaneously (group 1) or with mechanical ventilation to control fetal arterial PCO₂ (group 2). Hyperbaric oxygenation raised fetal Pa(sub O₂) by 20 Torr in both groups. During hyperbaric oxygenation, the occurrence of arousal increased severalfold in both groups. The occurrence of FBM increased during arousal in both groups, during rapid-eye-

51 LIFE SCIENCES (GENERAL)

movement sleep in group 1, and during NREM sleep in group 2. The timing of diaphragmatic activity during arousal and the variability of diaphragmatic activity during NREM sleep were different than those in rapid-eye-movement sleep. We conclude that oxygenation without lung distension increases the occurrence of arousal and of FBM, principally during arousal and NREM sleep, in fetuses of less than or equal to 135 days of gestation. Author (Herner)

A95-68928

PULMONARY GAS EXCHANGE DURING EXERCISE IN ATHLETES. 1: VENTILATION-PERFUSION MISMATCH AND DIFFUSION LIMITATION

SUSAN R. HOPKINS, DONALD C. MCKENZIE, ROBERT B. SCHOENE, ROBB W. GLENNY, and H. THOMAS ROBERTSON *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 912-917 Research sponsored by Medical Research Council of Canada and British Columbia Lung Association (HTN-95-A0145) Copyright

To investigate pulmonary gas exchange during exercise in athletes, 10 high aerobic capacity athletes (maximal aerobic capacity = 5.15 ± 0.52 l/min) underwent testing on a cycle ergometer at rest, 150 W, 300 W, and maximal exercise (372 ± 22 W) while trace amounts of six inert gases were infused intravenously. Arterial blood samples, mixed expired gas samples, and metabolic data were obtained. Indexes of ventilation-perfusion (dot-VA/dot-Q) mismatch were calculated by the multiple inert gas elimination technique. The alveolar-arterial difference for O₂ (AaD(sub O₂)) was predicted from the inert gas model on the basis of the calculated dot-VA/dot-Q mismatch. dot-VA/dot-Q heterogeneity increased significantly with exercise and was predicted to increase the AaD(sub O₂) by greater than 17 Torr during heavy and maximal exercise. The observed AaD(sub O₂) increased significantly more than that predicted by the inert gas technique during maximal exercise (10 ± 10 Torr). These data suggest that this population develops diffusion limitation during maximal exercise, but dot-VA/dot-Q mismatch is the most important contributor (greater than 60%) to the wide AaD(sub O₂) observed. Author (Herner)

A95-68929

LOWER LIMIT OF BODY FAT IN HEALTHY ACTIVE MEN

KARL E. FRIEDL, ROBERT J. MOORE, LESTER E. MARTINEZ-LOPEZ, JAMES A. VOGEL, E. WAYNE ASKEW, LOUIS J. MARCHITELLI, REED W. HOYT, and CLAIRE C. GORDON *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 933-940 (HTN-95-A0146) Copyright

We examined body composition changes in 55 normal young men during an 8-wk Army combat leadership training course involving strenuous exercise and low energy intake, with an estimated deficit of 5.0 ± 2.0 MJ/day and a resultant $15.7 \pm 3.1\%$ weight loss. Percent body fat (BF) measured by dual-energy X-ray absorptiometry (DEXA) averaged 14.3% (range 6 - 26%) and $5.8 \pm 1.8\%$ (range 4 - 11%) at the beginning and end of the course, respectively. Men who achieved a minimum percent BF (4 - 6%) by 6 wk demonstrated only small additional total and subcutaneous fat losses in the final 2 wk and sacrificed increasingly larger proportions of fat-free mass. Percent BF estimated from skinfold thickness reflected changes in fat mass, although actual percent BF was overestimated. Instead of reaching a plateau after fat stores were substantially depleted, abdominal, hip, and thigh girths continued to decline with body weight loss. Final percent BF for the leanest men was similar to that observed after a 25% body weight reduction in the 1950 Minnesota study (5.2% by underwater weighing), and height-corrected final fat mass was the same, suggesting that these values represent a minimal body fat content in healthy men and that weight loss subsequent to achieving this level is contributed from the fat-free mass. Our results suggest that 4 - 6% BF approximately 2.5 kg fat represents the lower limit for healthy men, as assessed by DEXA or by underwater weighing. Author (Herner)

A95-68930

CATECHOLAMINE RESPONSES TO SHORT-TERM HIGH-INTENSITY RESISTANCE EXERCISE OVERTRAINING

A. C. FRY, W. J. KRAEMER, F. VAN BORSELEN, J. M. LYNCH, N. T. TRIPLETT, L. P. KOZIRIS, and S. J. FLECK *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 941-946 (HTN-95-A0147) Copyright

Seventeen weight-trained males were divided into an overtraining group that weight trained their legs daily for 2 wk with 100% 1 repetition maximum relative intensity on a squat machine and a control group that exercised 1 day/wk with low relative intensity (50% 1 repetition maximum). Test batteries including strength assessments and resting and exercise-induced concentrations of epinephrine and norepinephrine were conducted at the beginning, middle, and end (tests 1 - 3, respectively) of the study. Strength capabilities decreased by test 3 for the OT group. Resting catecholamine concentrations did not change either group during the study, whereas exercise-induced concentrations of both epinephrine and norepinephrine significantly increased by tests 2 and 3 for only the OT group. Correlation coefficients suggested decreased responsiveness of skeletal muscle to sympathetic nervous system activity. It appears that altered exercise-induced sympathetic nervous system accompanies high relative intensity resistance exercise overtraining and may be among the initial responses to the onset of the previously theoretical sympathetic overtraining syndrome. Author (Herner)

A95-68931

INTERACTIVE EFFECTS OF EMPHYSEMA AND MALNUTRITION ON DIAPHRAGM STRUCTURE AND FUNCTION

MICHAEL I. LEWIS, STEPHEN A. MONN, WEN-ZHI ZHAN, and GARY C. SIECK *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 947-955 Research sponsored by American Lung Association of California (Contract(s)/Grant(s): NIH-HL-01907; NIH-HL-34817; NIH-HL-37680; NIH-HL-01907) (HTN-95-A0148) Copyright

Interactive effects of emphysema (EMP) and prolonged nutritional deprivation (ND) on contractile, morphometric, and metabolic properties of hamster diaphragm muscle (DIA) were examined. Six months after induction of EMP (intratracheal elastase), saline-treated controls (CTL) and EMP hamsters of similar body weights were subjected to ND over 6 wk. Isometric contractile and fatigue properties of costal DIA were determined in vitro. DIA fibers were histochemically classified as type I or II, and fiber succinate dehydrogenase activity and cross-sectional area were determined using quantitative microscopic procedures. From histochemical sections, the number of capillaries per fiber (C/F) and per fiber cross-sectional area (C/A) were determined. ND resulted in progressive loss of body weight. ND did not affect reduction in optimal length (L(sub O)) of DIA fibers in EMP compared with CTL and ND-CTL hamsters. Maximum specific force (i.e., force/unit area) was reduced by approximately 25% in EMP animals compared with CTL. ND did not improve or exacerbate the reduction in specific force with EMP. ND attenuated improved fatigue resistance of DIA in EMP animals. No differences in fiber type proportions were noted among experimental groups. Significant atrophy of type I and II DIA fibers was noted after ND. Atrophy was proportionately greater in type II fibers of ND-EMP when referenced to EMP animals. Thus adaptive hypertrophy of type II DIA fibers in EMP animals was abolished. Fiber succinate dehydrogenase activity was significantly increased in type I and II fibers in EMP DIA. ND did not affect this metabolic adaptation of DIA fibers to persistent loads imposed by EMP. In EMP hamsters, C/F was increased for type I and II fibers, whereas C/A was unaffected. Significant increments in both indexes of capillary were evident in ND-EMP DIA. We conclude that interaction of ND with EMP has serious negative clinical implications, in that total force production, endurance, and functional force reserve of the DIA will be significantly curtailed. Author (Herner)

A95-68932

HIGH PHYSIOLOGICAL LEVELS OF EPINEPHRINE DO NOT ENHANCE MUSCLE GLYCOGENOLYSIS DURING TETANIC STIMULATION

ALAN CHESLEY Guelph Univ., Guelph, Canada, DAVID J. DYCK Guelph Univ., Guelph, Canada, and LAWRENCE L. SPRIET Guelph Univ., Guelph, Canada Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 956-962 Research sponsored by Natural Sciences and Engineering Research Council of Canada (HTN-95-A0149) Copyright

This study examined whether high physiological concentrations of epinephrine (EPI) would enhance muscle glycogenolysis during intense muscular contractions. Muscles of the rat hindlimb were perfused for 12 min at rest and 45 s of tetanic stimulation (1.0-Hz train rate, 100-ms train duration at 80 Hz) without EPI (control) or with 15 or 35 nM EPI. In the EPI groups the muscles were perfused with EPI for the last 2 min of rest perfusion and throughout stimulation. Glycogenolysis in the white gastrocnemius, red gastrocnemius, plantaris, and soleus muscles during stimulation was unaffected by the presence of EPI in the perfusion medium. In addition, muscle lactate and hindlimb lactate efflux were similar in EPI and control groups. It is concluded that EPI is not important for enhancing glycogenolysis in rat muscles composed predominantly of fast-twitch fibers during intense short-term tetanic stimulation.

Author (Herner)

A95-68933

LUNG VOLUMES AND EXPIRATORY FLOW LIMITATION DURING EXERCISE IN INTERSTITIAL LUNG DISEASE

D. D. MARCINIUK Royal Univ. Hospital, Saskatchewan, Canada, G. SRIDHAR Royal Univ. Hospital, Saskatchewan, Canada, R. E. CLEMENS Royal Univ. Hospital, Saskatchewan, Canada, T. A. ZINTEL Royal Univ. Hospital, Saskatchewan, Canada, and C. G. GALLAGHER Royal Univ. Hospital, Saskatchewan, Canada Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 963-973 Research sponsored by Saskatchewan Lung Association and Saskatchewan Health Research Board (HTN-95-A0150) Copyright

Lung volumes were measured at rest and during exercise by an open-circuit N₂-washout technique in patients with interstitial lung disease (ILD). Exercise tidal flow-volume (F-V) curves were also compared with maximal F-V curves to investigate whether these patients demonstrated flow limitation. Seven patients underwent 4 min of constant work rate bicycle ergometer exercise at 40, 70, and 90% of their previously determined maximal work rates. End-expiratory lung volume and total lung capacity were measured at rest and near the end of each period of exercise. There was no significant change in end-expiratory lung volume or total lung capacity when resting measurements were compared with measurements at 40, 70, and 90% work rates. During exercise, expiratory flow limitation was evident in four patients who reported stopping exercise because of dyspnea. In the remaining patients who discontinued exercise because of leg fatigue, no flow limitation was evident. In all patients, the mean ratio of maximal minute ventilation to maximal ventilatory capacity (calculated from maximal F-V curves) was 67%. We conclude that lung volumes during exercise do not significantly differ from those at rest in this population and that patients with ILD may demonstrate expiratory flow limitation during exercise. Furthermore, because most patients with ILD are not breathing near their maximal ventilatory capacity at the end of exercise, we suggest that respiratory mechanics are not the primary cause of their exercise limitation.

Author (Herner)

A95-68934

SURFACE ACTIVITY OF LIPID EXTRACT SURFACTANT IN RELATION TO FILM AREA COMPRESSION AND COLLAPSE
SAMUEL SCHUERCH, DAVID SCHUERCH, TORE CURSTEDT,

and BENGT ROBERTSON Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 974-986 Research sponsored by Medical Research Council of Canada, Alberta Heritage Foundation for Medical Research, Swedish Medical Research Council, and Oscar II:s Jubileumsfond (HTN-95-A0151) Copyright

The physical properties of modified porcine surfactant (Curosurf), isolated from minced lungs by extraction with chloroform-methanol and further purified by liquid-gel chromatography, were investigated with the captive bubble technique. Bubble size, and thus the surface tension of an insoluble film at the bubble surface, is altered by changing the pressure within the closed bubble chamber. The film surface tension and area are determined from the shape (height and diameter) of the bubble. Adsorption of fresh Curosurf is characterized by stepwise decreases in surface tension, which can easily be observed by sudden quick movements of the bubble apex. The 'adsorption clicks' imply a cooperative movement of large collective units of molecules, approximately 10(exp 14) (corresponding to approximately 120 ng of phospholipid) or approximately 10(exp 18) molecules/sq m, into the interface during adsorption. Films formed in this manner are already highly enriched in dipalmitoyl phosphatidylcholine, as seen by the extremely low compressibility, close to that of dipalmitoyl phosphatidylcholine. Near-zero minimum tensions are obtained, even at phospholipid concentrations as low as 50 microgram/ml. During dynamic cycling (20 - 50 cycles/min), low minimum surface tensions, good film stability, low compressibility, and maximum surface tensions between 30 and 40 mN/m are possible only if the films are not overcompressed near zero surface tension; i.e., the overall film area compression should not substantially exceed 30%.

Author (Herner)

A95-68935

INDEX FINGER POSITION AND FORCE OF THE HUMAN FIRST DORSAL INTEROSSEUS AND ITS ULNAR NERVE ANTAGONIST

I. ZIJDEWIND Amsterdam Univ., Academisch Medisch Centrum, Amsterdam, Netherlands and D. KERNELL Groningen Univ., Groningen, Netherlands Journal of Applied Physiology (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 987-997 Research sponsored by Nederlands Organization for Scientific Research (HTN-95-A0152) Copyright

In normal subjects, maximum voluntary contraction (MVC) and electrical ulnar nerve stimulation (UNS; 30-Hz bursts of 0.33 s) were systematically compared with regard to the forces generated in different directions (abduction/adduction and flexion) and at different degrees of index finger abduction. With a 'resting' hand position in which there was no index finger abduction, UNS produced about one-half of the abduction force elicited by an MVC (mean ratio 51%). Qualitatively, such a discrepancy would be expected, because UNS activates two index finger muscles with opposing actions in the abduction/adduction plane of torques: the first dorsal interosseus (FDI) and the first palmar interosseus (FPI). The abduction forces produced by MVC and UNS were very sensitive to index finger abduction angle: at a maximum degree of abduction, the UNS-generated force even reversed its direction of action to adduction (with FPI dominating) and the abduction MVC declined to 37% of that in the resting hand position. Inasmuch as these declines in MVC and UNS-generated abduction force could not be explained by a change in moment arm, the main alternative seemed to be abduction-associated alterations in FDI fiber length (analysis by previously published biomechanical data). The FDI and FPI were further compared by application of a UNS-generated fatigue test (5-min burst stimulation), with the index finger kept at a 'neutral' angle, i.e., the abduction angle at which, in the unfatigued state, the forces of the FDI and FPI were in balance (zero net UNS-generated abduction/adduction force). There was no major difference in the fatigability of the two muscles, yet the fatigue reactions of the two muscles were not fully identical.

Author (Herner)

A95-68936

CARDIOPULMONARY ADAPTATIONS TO PNEUMONECTOMY IN DOGS. 4: MEMBRANE DIFFUSING CAPACITY AND CAPILLARY BLOOD VOLUME

C. C. W. HSIA Texas Univ. Southwestern Medical Center, Dallas, TX, US, L. F. HERAZO Texas Univ. Southwestern Medical Center, Dallas, TX, US, M. RAMANATHAN Texas Univ. Southwestern Medical Center, Dallas, TX, US, and R. L. JOHNSON, JR. Texas Univ. Southwestern Medical Center, Dallas, TX, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 998-1005 Research sponsored by American Heart Association (Contract(s)/Grant(s): NIH-R01-HL-40070; NIH-TL-07362) (HTN-95-A0153) Copyright

Lung diffusing capacity for carbon monoxide ($DL(\text{sub CO})$) and its components, membrane diffusing capacity ($Dm(\text{sub CO})$) and capillary blood volume (V_c), as well as pulmonary blood flow (dot-QC), were measured at rest at several lung volumes and during treadmill exercise by a rebreathing technique in four adult dogs after right pneumonectomy (R-PNX) and in six matched control dogs (Sham) 6 - 12 mo after surgery. In both groups, lung inflation at rest was associated with a small increase in $DL(\text{sub CO})$ and $Dm(\text{sub CO})$ but not in V_c . After R-PNX, total $DL(\text{sub CO})$ was lower by 30% at peak exercise compared with control values. When compared with $DL(\text{sub CO})$ in a normal left lung, $DL(\text{sub CO})$ in the remaining lung continued to increase along the normal relationship with respect to dot-QC up to a cardiac output equivalent to 34 l/min through both lungs of the Sham dog. There was no evidence of an upper limit of $DL(\text{sub CO})$ being reached. The augmentation of $DL(\text{sub CO})$ from rest to exercise was associated with corresponding increases in $Dm(\text{sub CO})$ and V_c ; after R-PNX, both $Dm(\text{sub CO})$ and V_c continued to increase with respect to dot-QC along similar relationships as in control dogs without reaching an upper limit, suggesting a much larger alveolar-capillary reserve for gas exchange by diffusion than previously recognized. At higher levels of blood flow through the remaining lung, $DL(\text{sub CO})$ was greater in adult dogs after R-PNX than after left pneumonectomy (Carlin et al., (1991)), suggesting that additional sources of compression, e.g., lung growth, exist after removal of greater than 50% of lung. Author (Herner)

A95-68937

(31)P-MRS AND SIMULTANEOUS QUANTIFICATION OF DYNAMIC HUMAN QUADRICEPS EXERCISE IN A WHOLE BODY MR SCANNER

J. B. RODENBURG, R. W. DE BOER, J. A. L. JENESON, C. J. A. VAN ECHTEL, and P. R. BAER *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 1021-1029 (HTN-95-A0154) Copyright

An ergometer for dynamic quadriceps exercise in a magnetic resonance (MR) scanner is physiologically validated, and its technical aspects are presented. The reproducibility of heart rate (HR), O_2 consumption (dot- VO_2), and power (P) during two graded exercise on the MR ergometer was good ($n = 8$). Graded exercises on the MR ergometer and on a cycle ergometer ($n = 17$) were similar with respect to the regression lines between (1) HR and dot- VO_2 and (2) HR and P; also peak P did not differ significantly. Peak HR, peak dot- VO_2 , and the slope of the regression line between P and dot- VO_2 were lower for MR exercise. During quadriceps exercise in an MR scanner ($n = 12$), peak P was 64 - 143 W for the right leg, with corresponding inorganic phosphate-to-phosphocreatine ratios of 0.85 - 7.2. It is concluded that continuous noninvasive assessment of energy metabolism with (31)P-MR spectroscopy and quantification of power output can be performed simultaneously during dynamic quadriceps exercise, without major reduction of the spectral resolution or the signal-to-noise ratio, and that exercise on this MR ergometer currently is the best possible approximation of cycling exercise for MR purposes. Author (Herner)

A95-68938

EVANS BLUE DYE IN THE ASSESSMENT OF PERMEABILITY-SURFACE AREA PRODUCT IN PERFUSED RAT LUNGS

MOHAMMED M. DALLAL Northwestern Univ. Medical School, Chicago, IL, US and SHIH-WEN CHANG Dept. of Veterans Affairs Lakeside Medical Center, Chicago, IL, US *Journal of Applied Physiology* (ISSN 8750-7587) vol. 77, no. 2 August 1994 p. 1030-1035 Research sponsored by Dept. of Veterans Affairs (HTN-95-A0155) Copyright

Evans blue dye (EBD) has been used extensively as a marker of extravascular protein leakage. We assessed the utility of EBD as an albumin marker in the measurement of permeability-surface area product (PS) in perfused rat lungs and compared the results with PS values obtained using I-125-labeled albumin. In isolated perfused rat lungs, PS was measured by exposure to a perfusate containing EBD (600 microgram/ml) and I-125-albumin (1 micro Ci) for exactly 3 min, followed by washout of the intravascular space with fresh perfusate for 6 min. In lungs from normal rats, we found that PS obtained by EBD (PS-EBD) was fivefold higher than PS obtained by I-125-albumin. Similarly, in rats pretreated with Salmonella enteritidis lipopolysaccharide (2 mg/kg iv), PS-EBD was much higher than PS-I-125. This discrepancy between PS-EBD and PS-I-125 was not explained by difference in the amount of free marker in perfusate, because the albumin-binding rate for both markers was very high. In addition, prolonged perfusion (40 min) with EBD did not significantly affect pulmonary vasoreactivity or water content in rat lungs. A detailed comparison of the kinetics of lung tissue uptake of the two markers showed an initial phase of rapid lung uptake of EBD, followed by parallel uptake of both markers up to 60 min of perfusion. We conclude that although EBD does not cause obvious lung injury, it is not a reliable marker for measurement of vascular permeability in perfused rat lungs. This is most likely due to rapid binding of EBD to lung tissue proteins. Author (revised by Herner)

N95-19536# Naval Research Lab., Washington, DC.

PROCEEDINGS OF NATURAL SCIENCES SYMPOSIUM

Jul. 1994 84 p Symposium held in Washington, DC, 14 Jun. 1993 (AD-A285452; NRL/PU/6000-94-252) Avail: CASI HC A05/MF A01

This symposium included seven Nobel Laureates' presentations: Prospects for the Physical Sciences; Creativity and Managed Research; Science — The Preeminent Random Process; Dealing with Genes; Understanding Life as Chemistry; The Impossible Takes a Little Longer; and Reflections of an Era. This last presentation is made by Dr. Jerome Karle, Nobel Laureate, Chemistry, 1985, for whom the conference marked a 75th birthday and 45th year in government service. The focus of these talks was an outlook on the 'Future Implications of Current Advances in the Natural Sciences'. CASI

N95-19613# Department of Energy, Washington, DC. Office of Energy Research.

ACTIVITIES OF THE DIVISION OF ENERGY BIOSCIENCES

Annual Summary Report, FY 1994

Sep. 1994 172 p

(DE95-001046; DOE/ER-0621P) Avail: CASI HC A08/MF A02

The Energy Biosciences program was initiated in 1979 for the purpose of fulfilling the need of basic information about plants and microorganisms relating to varied energy matters. The program generates basic information that contributes significantly to future technologies involving alternate fuel generation, petroleum replacements, sustained industrial activities along with means of improving environmental conditions. The Energy Biosciences program during Fiscal Year 1994 received 139 new research applications following the screening of numerous preapplications. Of the applications received, 21 projects were funded. DOE

AEROSPACE MEDICINE

Includes physiological factors; biological effects of radiation; and effects of weightlessness on man and animals.

A95-65873

CHARACTERIZATION OF FRACTURE TOUGHNESS OF RENAL CALCULI USING A MICROINDENTATION TECHNIQUE

P. ZHONG Univ. of Texas Southwestern Medical Center at Dallas, Dallas, TX, C. J. CHUONG, and G. M. PREMINGER *Journal of Materials Science Letters* (ISSN 0261-8028) vol. 12, no. 18 September 15, 1993 p. 1460-1462 refs (BTN-94-EIX94361122293) Copyright

We report our results on the characterization of fracture toughness of renal calculi of various compositions. A microindentation technique was used to determine stone hardness. The results showed distinct differences in fracture toughness of renal calculi which confirms the observations of clinical experience: viz. COM and brushite stones are much more resistant to shock-wave fragmentation than MAPH and CA stones. EI

A95-65904

FLUID FLOW AND HEAT TRANSFER IN THE CRESCENT-SHAPED LUMEN CATHETER

M. A. EBADIAN Florida Inst. Univ., Miami, FL and H. Y. ZHANG *Journal of Applied Mechanics, Transactions ASME* (ISSN 0021-8936) vol. 60, no. 3 September 1993 p. 721-727 (BTN-94-EIX94361122408) Copyright

This paper presents a numerical investigation of fluid flow, frequency response in the fully developed region, and convective heat transfer in the entrance region of the crescent-shaped lumen catheter. The catheter is commonly used in the biomedical field to clinically diagnose heart disease and also to treat vessel blockage in surgery. The catheter is subjected to a constant wall temperature. The solution to discretization of the momentum and energy equations is obtained by using the numerically generated boundary fitted coordinate system. According to this method, the complex domain in the physical plane is transformed into a regular square domain in the computational plane. The control volume-based finite difference method is then used to discretize the transformed governing equations. Results for the thermal entry region flow, frequency response, and heat transfer are presented in graphical form. The representative curves illustrating variations of the flow rate, frequency response, damping coefficient, bulk temperature, and the Nusselt numbers with pertinent parameters in the entire thermal entry region are plotted. The optimized catheter design for diagnostic use in the medical industry is also presented graphically. Author (EI)

A95-65949

MAXIMALLY SMOOTH IMAGE RECOVERY IN TRANSFORM CODING

YAO WANG Polytechnic Univ, Brooklyn, NY, QIN-FAN ZHU, and LEONARD SHAW *IEEE Transactions on Communications* (ISSN 0090-6778) vol. 41, no. 10 October 1993 p. 1544-1551 refs (BTN-94-EIX94361135047) Copyright

This paper considers the reconstruction of images from partial coefficients in block transform coders and its application for packet loss recovery in image transmission over ATM networks. The proposed algorithm makes use of the smoothness property of common image signals and produces a maximally smooth image among all those with the same coefficients and boundary conditions. It recovers each damaged block by minimizing the intersample variation within the block and across the block boundary. The optimal solution is achievable through two linear transformations, where the transform matrices depend on the loss pattern and can be calculated in advance. The reconstruction of contiguously damaged blocks is accomplished iteratively, using the previous solution as the boundary conditions in each new step. This technique is applicable

to any unitary block-transform and is very effective for recovering the DC and low-frequency coefficients. When applied to still image coders using the discrete cosine transform (DCT), high quality images have been reconstructed in the absence of many DC and low-frequency coefficients over spatially adjacent blocks. When the damaged blocks are made isolated by the use of block interleaving, satisfactory results have been obtained even when all the coefficients are missing. Author (EI)

A95-65976

QUANTITATIVE SPECT BRAIN IMAGING: EFFECTS OF ATTENUATION AND DETECTOR RESPONSE

D. R. GILLAND Duke Univ Medical Cent, Durham, NC, R. J. JASZCZAK, J. E. BOWSER, T. G. TURKINGTON, Z. LIANG, and K. L. GREER *IEEE Transactions on Nuclear Science* (ISSN 0018-9499) vol. 40, no. 3 June 1993 p. 295-299 refs (BTN-94-EIX94361133479) Copyright

The purpose of this work was to implement two reconstruction methods that compensate for attenuation and detector response, a 3D maximum likelihood-EM method (ML) and a filtered backprojection method (FB) with Metz filter and Chang attenuation compensation, and compare the methods in terms of quantitative accuracy and image noise. The methods were tested on simulated data of the 3D Hoffman brain phantom. The simulation incorporated attenuation and distance-dependent detector response. Bias and standard deviation of reconstructed voxel intensities were measured in the grey and white matter regions. The results with ML showed that in both the grey and white matter regions as the number of iterations increased. In both regions, ML had smaller standard deviation than FB for a given bias. Reconstruction times for the ML method have been greatly reduced through efficient coding, limited source support, and any computing attenuation factors only along rays perpendicular to the detector. Author (EI)

A95-68268* National Aeronautics and Space Administration. Langley Research Center, Hampton, VA.

RADIATION EXPOSURE PREDICTIONS FOR LONG-DURATION-STAY MARS MISSIONS

SCOTT A. STRIEPE, LISA C. SIMONSEN, and JOHN E. NEALY *Journal of the Astronautical Sciences* (ISSN 0021-9142) vol. 42, no. 2 April-June 1994 p. 131-142 refs (BTN-95-EIX95032453043) Copyright

In this study, the ionizing radiation environment is estimated, using the Mission Radiation Calculation (MIRACAL) program, for several long-duration-stay Mars missions proposed for early in the 21(st) century. Both minimum energy and fast transfer missions are evaluated and their 30-day maximum, annual maximum, and total slab skin and blood-forming organ (BFO) doses are compared. When large flares were included while the astronauts were on the surface, the delivered dose did not significantly contribute to the total dose (less than 4 cSv BFO dose, or 8 percent of the guideline annual limit, for the most energetic event simulated) due to the substantial protection provided by the Martian atmosphere. However, dose delivered by large flares during transit is dependent on vehicle shielding and distance from the Sun. All of the fast transfer missions studied had lower total and annual maximum doses than the corresponding minimum energy transfer missions (on average, 30% less for missions having no large flares and the shielding thicknesses evaluated in this study). For all the missions studied, having the astronauts spend one-third of their day during transit in a 10 g/sq cm storm shelter resulted in an approximate 10% reduction in the total mission dose. Author (EI)

A95-69746

A MULTIMODAL METHOD FOR ASSESSING AND TREATING AIRSICKNESS

R. JEFFREY JACKSON U.S. Air Force Academy, Colorado Springs, CO, US *International Journal of Aviation Psychology* (ISSN 1050-8414) vol. 4, no. 1 1994 p. 85-96 (HTN-95-90719) Copyright

Airsickness is a disruptive phenomenon in military flight train-

ing and operational settings. Despite its occurrence, several studies have demonstrated efficacy in treating this syndrome. Although generally successful, these treatment strategies appear to be guided by their presumed effectiveness rather than by theoretical rationale. In support of these varied treatments, a multimodal conceptual model is proposed. This model attempts to integrate those factors bearing on the etiology of airsickness and, based on a careful assessment, directs and explains the treatment efforts.

Author (Herner)

N95-19537# Physics and Electronics Lab. TNO, The Hague (Netherlands).

HIGH PERFORMANCE COMPUTING FOR MEDICAL IMAGE INTERPRETATION

A. C. DUMAY Oct. 1993 72 p

(AD-A285457; FEL-93-B160; TDCK-93-2783) Avail: CASI HC A04/MF A01

Medical image processing aims at improvement of the image quality in order to support the medical diagnosis and therapy. Globally, there can be four subjects distinguished for image processing, including: (1) data fusion, (2) object/feature extraction, (3) analysis, and (4) visualization. Data fusion, or registration, concerns the combination of images acquired with various imaging modalities into a single image. Object/feature extraction, or segmentation, aims at the recognition and extraction of objects and object features from image data. And the analysis involves the interpretation of the object features from which conclusions can be drawn. Visualization is of utmost importance for each of the forementioned subjects. With the introduction of computer tomography (CT) and magnetic resonance imaging (MRI) three-dimensional images can be acquired from patients. The processing of 3D images such that the time required for that purpose is in acceptable limits makes the use of high performance computing (HPC) techniques desirable. Currently (commercially) available implementations do not satisfy the performance requirements for routine use in a clinical setting. As a result, such systems will not be used at all or at high costs of personnel. High performance computing medical image interpretation includes data fusion, object/feature extraction, analysis, and visualization of 3D medical images. Modules for these purposes may be integrated in existing medical systems to support interactive visualization, quantitative analysis, radiotherapy planning, and neurosurgery planning. Integrated into a high performance computing medical image interpretation system (HPMI(sup 2)S), one has the tools for developing anatomical models for simulation and training for example minimal access surgery (MAS) and triage. DTIC

N95-19695# Keele Univ. (England). Dept. of Chemistry.

PHOTOBIOLOGY IN MEDICINE Final Proceedings Report, 30 Sep. - 13 Oct. 1993

T. GEORGE TRUSCOTT, ed. 13 Oct. 1993 200 p Sponsored by European Office of Aerospace Research and Development (AD-A284843; CSP-93-1041) Avail: CASI HC A09/MF A03

Man has recognized an association of light with life and medicine for over 3000 years. Today the major challenges to this topic include the elucidation of photochemical reactions involved in photobiology at the molecular level. This includes the use of a variety of modern probing techniques that directly measure the reactivity of excited states and free radicals involved in biological reactions. This textbook is based on such an approach and has arisen from some of the lectures delivered at the NATO ASI held at Hotel Capo Caccia near the Centre for Advanced Research in Photobiology (CARP) in Cardegia, Italy. The ASI took place from 30 Sep. - 13 Oct. 93 and involved a total membership of 90. Photobiology in medicine is a wide-ranging subject that obviously cannot be totally covered comprehensively in one book alone. Rather, chosen topics of current interest in photomedicine were focused upon and these are linked to the major underlying themes and techniques in photophysics and photochemistry. DTIC

N95-19698# Alabama Univ., Birmingham, AL.

CUMULATIVE EFFECTS OF REPEATED BRIEF CEREBRAL

ISCHEMIA Final Report

H. P. HETHERINGTON and K. A. CONGER 1994 48 p (Contract(s)/Grant(s): F49620-92-J-0362)

(AD-A284234; AFOSR-94-0481TR) Avail: CASI HC A03/MF A01

The purpose of this grant was to investigate the metabolic and physiological factors determining the extent of the ischemic damage in a model of brief repetitive cerebral ischemia. The model is intended to simulate the effects of Gz induced blackout experienced by fighter pilots undergoing high gravitational stress maneuvers. To achieve this goal a rat model was developed whereby the ischemia was remotely induced by inflation (under computer control) of an occlusive cuff placed about the common carotid artery. Metabolic parameters were determined using in vivo NMR spectroscopy measurements throughout the ischemic and reflow periods. Lactate production was found to be highly reproducible and not limited by blood glucose levels. High energy phosphate decreases were correlated with loss of high frequency EEG. To evaluate the correlation between histologic damage and measures of regional metabolism we have carried out H-1 spectroscopic imaging experiments with histologic evaluations of the brain three hours after reperfusion. These studies were acquired with 5 minute (two dimensional mapping across the cortex) and 16 sec time resolution (one dimensional mapping) during the ischemia and reflow. Preliminary results of regional clearance kinetics of lactate shows a strong correlation with the presence of histologic damage. DTIC

N95-19714# Rothe Development, Inc., San Antonio, TX.

ECHO IMAGING TECHNIQUES DETERMINE THE SIZE OF INTRAVASCULAR BUBBLES IN DECOMPRESSION

SICKNESS Final Report

ROBERT M. OLSON Jul. 1994 30 p

(Contract(s)/Grant(s): F33615-89-D-0604)

(AD-A284849; AL/CF-TR-1994-0033) Avail: CASI HC A03/MF A01

The size of altitude induced intravenous bubbles was determined. These bubbles, referred to as venous gas emboli (VGE), are thought to be a major factor in the onset and severity of decompression sickness (DCS). Ten volunteer subjects were monitored for altitude induced VGE in the inferior vena cava (IVC) with a HP Sonos 1000 echo imaging system. Bubble size was determined indirectly and by in-vitro sizing methods because ultrasonic images do not necessarily represent true VGE size. Stratification of bubbles of known size in a water filled mechanical analog of the IVC determined the upper size limit, which was 300 microns in these experiments; bubbles larger than 300 microns roll along the top of the vessel. The lower size limit was the size of the smallest bubbles which the Sonos 1000 could image, determined indirectly by in-vitro microbubble flotation rates and survival times measured ultrasonically and calibrated microscopically. The diameter of the smallest VGE which echo imaging systems can detect in the IVC was found to be 30-40 microns. Used in this determination was the fact that these microbubbles are too small to float; they survive less than a minute. In conclusion, the size of the ultrasonically detected VGE in the interior of the IVC of decompressed subjects can be measured; for subjects at 29,500 ft. bubble size varied from 30-300 microns. DTIC

N95-19746* National Aeronautics and Space Administration, Washington, DC.

AEROSPACE MEDICINE AND BIOLOGY: A CONTINUING BIBLIOGRAPHY WITH INDEXES (SUPPLEMENT 397)

Jan. 1995 57 p

(NASA-SP-7011(397); NAS 1.21:7011(397)) Avail: CASI HC A04

This bibliography lists 122 reports, articles and other documents introduced into the NASA Scientific and Technical Information System during Jan. 1995. Subject coverage includes: aerospace medicine and physiology, life support systems and man/system technology, protective clothing, exobiology and extraterrestrial life, planetary biology, and flight crew behavior and performance.

Author

N95-19811 Army Research Inst. of Environmental Medicine, Natick, MA.

DISTRIBUTED MEDICAL DATABASE SYSTEM FOR REAL-TIME MONITORING OF THE HEALTH AND RISK EXPOSURE OF MILITARY RESEARCH STUDY VOLUNTEERS AT USARIEM

MATTHEW J. REARDON and DONNA CARDINAL May 1994 67 p Limited Reproducibility: More than 20% of this document may be affected by microfiche quality (AD-A279193; USARIEM-T94-13) Avail: Issuing Activity (Defense Technical Information Center (DTIC))

This report describes a computerized medical database system designed to track the health and study related risk exposures and adverse incidents of soldiers participating as volunteer test subjects in USARIEM human-use research studies. This database system was implemented as an Oracle application on the USARIEM DEC VAX. The data collected at remote computer terminals are transmitted via modem and LAN to secure digital storage media at USARIEM. Database file access is controlled using methods to ensure test subject confidentiality and anonymity with regard to medical information. Only physician users are allowed access to medical details in the database. This database system was designed to improve the tracking of research study related risks and medical incidents as well as improve the coordination of medical care provided to test subjects by USARIEM physicians. DTIC

N95-19827 Army Research Inst. of Environmental Medicine, Natick, MA.

ANNUAL HISTORICAL REPORT, CALENDAR YEAR 1993

Apr. 1994 174 p Limited Reproducibility: More than 20% of this document may be affected by microfiche quality (AD-A280178) Avail: CASI HC A08

This report contains information concerning the mission, organization, key staff, overall funding and significant research accomplishments of the US Army Research Institute of Environmental Medicine, a subordinate element of the US Army Medical Research and Development Command, for calendar year 1992. Also included are listings of published reports, abstracts, presentations and key briefings for each Research Division of the Institute and significant accomplishments and appointments of the professional staff. DTIC

53

BEHAVIORAL SCIENCES

Includes psychological factors; individual and group behavior; crew training and evaluation; and psychiatric research.

A95-69742* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

PILOT INTERACTION WITH COCKPIT AUTOMATION 2: AN EXPERIMENTAL STUDY OF PILOTS' MODEL AND AWARENESS OF THE FLIGHT MANAGEMENT SYSTEM

NADINE B. SARTER Ohio State University, Columbus, OH, US and DAVID D. WOODS Ohio State University, Columbus, OH, US International Journal of Aviation Psychology (ISSN 1050-8414) vol. 4, no. 1 1994 p. 1-28

(Contract(s)/Grant(s): NCC2-592) (HTN-95-90715) Copyright

Technological developments have made it possible to automate more and more functions on the commercial aviation flight deck and in other dynamic high-consequence domains. This increase in the degrees of freedom in design has shifted questions away from narrow technological feasibility. Many concerned groups, from designers and operators to regulators and researchers, have begun to ask questions about how we should use the possibilities afforded by technology skillfully to support and expand human

performance. In this article, we report on an experimental study that addressed these questions by examining pilot interaction with the current generation of flight deck automation. Previous results on pilot-automation interaction derived from pilot surveys, incident reports, and training observations have produced a corpus of features and contexts in which human-machine coordination is likely to break down (e.g., automation surprises). We used these data to design a simulated flight scenario that contained a variety of probes designed to reveal pilots' mental model of one major component of flight deck automation: the Flight Management System (FMS). The events within the scenario were also designed to probe pilots' ability to apply their knowledge and understanding in specific flight contexts and to examine their ability to track the status and behavior of the automated system (mode awareness). Although pilots were able to 'make the system work' in standard situations, the results reveal a variety of latent problems in pilot-FMS interaction that can affect pilot performance in nonnormal time critical situations.

Author (Herner)

A95-69743

USING KNOWLEDGE EXPLORATION TOOLS TO STUDY AIRSPACE COMPLEXITY IN AIR TRAFFIC CONTROL

RICHARD H. MOGFORD CTA Incorporated, Pleasantville, NJ, US, ELIZABETH D. MURPHY CTA Incorporated, Pleasantville, NJ, US, and JEREMY A. GUTTMAN CTA Incorporated, Pleasantville, NJ, US International Journal of Aviation Psychology (ISSN 1050-8414) vol. 4, no. 1 1994 p. 29-45 Research sponsored by the Federal Aviation Administration (HTN-95-90716) Copyright

A study was conducted at the Federal Aviation Administration's (FAA's) Jacksonville en route air traffic control center to identify factors that contribute to airspace complexity. Direct (verbal reports) and indirect (multidimensional scaling) procedures were used to identify potential factors. A list of candidate factors was created by combining the data from the direct and indirect knowledge exploration methods. Important complexity factors were identified by determining their simple and multiple correlations with overall sector complexity as judged by a group of Traffic Management Unit personnel (flow controllers). A final list of 16 complexity factors was developed and is suggested as a reference for future research in the area. An evaluation of the knowledge extraction techniques indicated that, although little unique information was generated by the indirect procedure, it was useful for the identification of complexity factors when combined with data from direct sources. Further research to validate the identified sector complexity factors is recommended. Author (Herner)

A95-69744

USE OF WORKLOAD REDLINES: A KC-135 CREW-REDUCTION APPLICATION

JUSTIN D. RUEB U.S. Air Force Academy, Colorado Springs, CO, US, MICHAEL A. VIDULICH Armstrong Laboratory, Wright-Patterson AFB, OH, US, and JOHN A. HASSOUN Crew Station Evaluation Facility, Wright-Patterson AFB, OH, US International Journal of Aviation Psychology (ISSN 1050-8414) vol. 4, no. 1 1994 p. 47-64

(HTN-95-90717) Copyright

In recent years, workload assessment has played an increased role in system design and evaluation. However, one concern in workload assessment has been the identification of how much workload is too much. This study established and used workload redlines in determining the feasibility of a two-person (no navigator) KC-135 conceptual cockpit design. Ten KC-135 crews and 2 KC-10 crews were required to fly several simulator missions over a period of 1 week. Objective and subjective measures of performance and workload were taken during and after each flight. The various performance measures and the subjective-workload measures provided convergent results supporting the feasibility of the conceptual design. Recommendations on the use of redline procedures and their future application are discussed. Author (Herner)

A95-69745* National Aeronautics and Space Administration. Langley Research Center, Hampton, VA.

COGNITIVE REPRESENTATIONS OF FLIGHT-DECK INFORMATION ATTRIBUTES

WENDELL R. RICKS NASA. Langley Research Center, Hampton, VA, US, JON E. JONSSON McDonnell Douglas Aerospace-West, Long Beach, CA, US, and WILLIAM H. ROGERS Bolt Beranek and Newman, Inc., Cambridge, MA, US International Journal of Aviation Psychology (ISSN 1050-8414) vol. 4, no. 1 1994 p. 65-83 (HTN-95-90718) Copyright

A large number of aviation issues are generically being called flight-deck information management issues, underscoring the need for an organization or classification structure. One objective of this study was to empirically determine how pilots organize flight-deck information attributes and — based upon that data — develop a useful taxonomy (in terms of better understanding the problems and directing solutions) for classifying flight-deck information management issues. This study also empirically determined how pilots model the importance of flight-deck information attributes for managing information. The results of this analysis suggest areas in which flight-deck researchers and designers may wish to consider focusing their efforts. Author (Herner)

N95-19538# Institute for Human Factors TNO, Soesterberg (Netherlands).

LEARNING EFFECTS ON STRATEGY SELECTION IN A DYNAMIC TASK ENVIRONMENT AS A FUNCTION OF TIME PRESSURE

J. H. KERSTHOLT Jun. 1994 17 p (AD-A285458; TNO-TM-1994-B-12; TDCK-94-0055) Avail: CASI HC A03/MF A01

Previous research on strategy selection in dynamic task environments indicated that subjects preferred to request information first, before an action was applied, even when the straightforward application of actions would have resulted in more optimal performance. Furthermore, this strategy was also used when subjects only had limited time for diagnosis. In the present experiment it was investigated whether the amount of training could account for the limited use that subjects made of the task dynamics. Subjects were required to monitor the changing fitness level of an athlete, by means of a graph on a computer screen, and to apply treatments whenever necessary. They could request various symptoms that would provide an indication for the cause underlying a possible fitness decline. The subjects either received limited training or elaborate training and they had either sufficient time for diagnosis, or worked under time pressure. The results showed that the amount of training did not affect the strategy that subjects used: half of the subjects used a judgment-oriented strategy and the other half used an action-oriented strategy in both training conditions. However, the well trained subjects were superior in selecting information and they processed the information at a faster rate than the subjects with only minimal training. The time pressure effects replicated previous findings: subjects used the same strategy and speeded up information processing. A high level of time pressure only deteriorated the information integration processes of the subjects who had received only minimal training, but not that of the well-trained subjects. DTIC

N95-19564# Army Aeromedical Research Lab., Fort Rucker, AL. Aircrew Health and Performance Div.

COLLECTION OF REAL-TIME, MULTICHANNEL EEG DATA FROM HELICOPTER PILOTS IN FLIGHT: A FEASIBILITY STUDY Final Report

JOHN A. CALDWELL, JR., JAMES A. LEWIS, STEPHEN R. DARLING, ROBERT M. DILLARD, and PARLEY P. JOHNSON May 1994 61 p (AD-A285050; USAARL-94-26) Avail: CASI HC A04/MF A01

An initial assessment of whether valid electroencephalographic (EEG) data could be collected on helicopter pilots in flight was conducted using 20 volunteers. Each subject provided a brief eyes-

open and eyes-closed EEG in the laboratory and then in a helicopter. During helicopter flights, data were monitored and recorded on the ground in real-time via a radio telemetry system. Initial hardware problems resulted in the exclusion of several records, but these problems were resolved toward the end of the study. Analysis of the remaining acceptable EEG records indicated that the typical increase in alpha activity (7.5-13.0 Hz) due to eye closure was clearly observable at every analyzed electrode site regardless of whether the testing occurred in the laboratory or in the helicopter. Further examinations of delta, theta, and beta bands showed that the EEG data were not overly contaminated by artifacts, although it was much more difficult to find 'clean' segments on records collected in the helicopter than in the laboratory. It was concluded that it is possible to record and telemeter useable EEG from the in-flight helicopter environment. DTIC

N95-19583# Prins Maurits Lab. TNO, Rijswijk (Netherlands). **MULTIPLE-TASK PERFORMANCE: A CRITICAL REVIEW OF THE LITERATURE AND A COGNITIVE NEUROSCIENCE FRAMEWORK**

J. E. KORTELING 10 Mar. 1994 91 p (AD-A285133; IZF-1994-B-5; TDCK-94-0043) Avail: CASI HC A05/MF A01

In our modern society, technological developments have altered the nature of jobs and tasks. In many work situations, operators are required to monitor, control, and manipulate information via complex technological systems. Such systems typically involve performance of several tasks in a limited period of time. In order to be able to optimize such technological systems, knowledge with regard to complex-task performance is needed, based on which technical products, processes, and systems involved in daily life can be matched to the capabilities and limitations of people. DTIC

N95-19602# New York Univ., New York, NY. Dept. of Psychology. **FACILITATION AND INTERFERENCE IN IDENTIFICATION OF PICTURES AND WORDS Final Report, 1 Dec. 1991 - 31 May 1994**

JOAN G. SNODGRASS 5 Oct. 1994 54 p (Contract(s)/Grant(s): F49620-92-J-0119) (AD-A285882; AFOSR-94-0670TR) Avail: CASI HC A04/MF A01

This research is concerned with long-term facilitation and short-term interference and facilitation in identification of pictures and words. The long-term facilitation occurs when subjects are exposed to some representation of the item during a study episode, and then show improved identification of that item during a retention test. This type of facilitation is known as priming (or long-term priming) and the retention test is known as an implicit or indirect test because subjects are not instructed to think back to the prior study episode during the test. Much of our recent research has concerned the relationship between performance on the implicit test of picture fragment completion and the explicit test of recognition memory. Our major interest has been on the importance of maintaining the same surface features between study and test on performance in both implicit and explicit tests: Contrary to previous findings that explicit tests are impervious to surface changes and only sensitive to changes in meaning, we have found performance decrements from changes in surface features in explicit as well as implicit tests. These surface changes have been as subtle as differences in the level of fragmentation between study and test and as extreme as differences in the form of item (picture vs. word) between study and test. The research carried out under the grant has exploited this similarity between explicit and implicit tests within a components-of-information model of memory which accommodates both associations and dissociations between the two classes of tests. DTIC

N95-19740 Rice Univ., Houston, TX.

COMPARING PERFORMANCE ON IMPLICIT MEMORY TESTS Annual Technical Report, 1 Aug. 1993 - 31 Jul. 1994 HENRY ROEDIGER, III 31 Aug. 1994 9 p Limited Reproducibility: More than 20% of this document may be affected by

microfiche quality

(Contract(s)/Grant(s): F49620-92-J-0437)
(AD-A284861; AFOSR-94-0577TR) Avail: CASI HC A02

The second year of this grant saw progress on nine projects. In particular, in the past year five papers or chapters have been published, six are in press, three are in preparation, and data are being collected on several new projects. We have completed projects on the following topics: (1) effects of imagery on nonverbal implicit tests; (2) effects of high priority events on implicit tests; (3) specificity of priming on verbal and nonverbal perceptual tests; (4) direct comparison of two methods of testing for contamination of implicit tests by conscious recollection; (5) the experimental basis of serial position effects; and (6) a new paradigm for the study of false memories. Four or five other projects should be completed during the final year of the grant. DTIC

54

MAN/SYSTEM TECHNOLOGY AND LIFE SUPPORT

Includes human engineering; biotechnology; and space suits and protective clothing.

A95-68263

ATMOSPHERIC PRESSURE WITHIN LUNAR STRUCTURE
PAUL S. NOWAK Gonzaga Univ., Spokane, WA, JEFFREY JANAKUS, and CHARLES E. MITCHELL Journal of Aerospace Engineering (ISSN 0893-1321) vol. 7, no. 4 October 1994 p. 398-410 refs
(BTN-95-EIX95032452326) Copyright

Design and construction of a structure on the Moon requires addressing a host of issues not encountered on Earth. Since there is no atmosphere on the Moon, a lunar structure must contain an artificial atmosphere. One critical design issue is the magnitude of the pressure of this atmosphere. Much of the current literature on the design of lunar structures assumes a pressure of 101.3 kPa (14.7 psi), corresponding to that at sea level on Earth, which is an order of magnitude larger than any other loading on the structure. An assessment of the outcome of lowering the internal pressure for a lunar structure is presented that accounts for human physiology, plant growth, mechanical equipment for gas circulation, structural aspects, leak rate, decompression, flammability, combustion, and economic issues. Options for the magnitude and content of an internal atmosphere for a lunar structure are given. Results clearly show that there is a great savings if the pressure is lowered by an amount that does not greatly affect the inhabitants' physiology or safety. Author (EI)

A95-68265

DESIGN AND CONSTRUCTION OF SHIELDED LUNAR OUTPOST

S. D. JOLLY Univ. of Colorado, Boulder, CO, J. HAPPEL, and S. STURE Journal of Aerospace Engineering (ISSN 0893-1321) vol. 7, no. 4 October 1994 p. 417-434 refs
(BTN-95-EIX95032452328) Copyright

The construction of an outpost on the Moon in which humans can live and work for periods exceeding six months will require special countermeasures to adapt to the hostile environment present at the lunar surface. Various inherent dangers such as meteoroids, galactic cosmic radiation, solar proton events, and large thermal extremes will drive the design configuration of the outpost. Other considerations such as lunar soil mechanics, equipment performance, mass delivery, risk, reliability, and tele-operability act strongly as constraints that shape and control the design alternatives. Analysis of these fundamental relationships have resulted in lunar civil engineering guidelines, which are unique to this domain, and these in turn have pointed to research areas needing further attention. A preliminary design is presented for a lunar outpost shelter. Additionally, the design methodology is explored, and early enabling

technologies are identified to facilitate an understanding of lunar shelter designs from an integrated system standpoint.

Author (EI)

A95-68277

MINIMIZING HUMAN-MACHINE INTERFACE FAILURES IN HIGH RISK SYSTEMS

JOHN J. SUDANO Government Electronic Systems, Moorestown, NJ and MARTIN MARIETTA IEEE Aerospace and Electronic Systems Magazine (ISSN 0885-8985) vol. 9, no. 10 October 1994 p. 17-20 refs
(BTN-95-EIX95042474623) Copyright

Technology now permits the building of very complex man-machine systems with centralized controls, with the result that many processes can be run by relatively few individual workers. Studies of failures within these complex systems indicate that they are usually the consequence of a series of highly complex coincidences. There is an institutional neglect or misunderstanding of the implications of low-probability, high-consequence events for the design of complex man-machine systems. Bhopal, Challenger, Chernobyl, the Swiss Chemical Spill, the Exxon Valdez, Seveso, Tenerife Airplane Crash, Three Mile Island - incidents like these emphasize the need to better understand the mechanisms of disasters in complex, high-risk systems. Hardware failures are the best understood component of failures in such systems. Software has historically been less well understood, but great strides are presently being made in understanding interactions between software design errors in these systems. Human error is the most complex and least understood factor in the failures of complex systems, accounting for as much as 60% to 80% of complex system failures and as much as 96% of simple system failures. We must stop designing systems in which we virtually guarantee that operator errors will occur with catastrophic consequences. The greatest payback in reducing high risk system accidents is to reduce catastrophes induced or exacerbated by human error. This paper discusses some task breakdowns between the human element and software/hardware. These task allocations allow the complex man-machine system to be designed more robustly and prevent human error so as to reduce possible catastrophic consequences. Author (EI)

A95-68352

OXYGEN SENSOR DEVELOPMENT FOR LIFE SUPPORT
JEFFERY T. CHEUNG Rockwell Science Center and SCOTT R. JOHNSON Aerospace Engineering (Warrendale, Pennsylvania) (ISSN 0736-2536) vol. 14, no. 9 September 1994 p. 7-9
(BTN-95-EIX95042477107) Copyright

Metal-oxide semiconductor thin films offer advantages as oxygen detection devices. Unlike potentiometric cells with a solid electrolyte, such as those in the space shuttle orbiter, these thin films have low to medium operating-temperature range to minimize power use and maximize longevity, satisfying requirements for space mission applications. Moreover, the entire sensor could be patterned into a miniaturized, integrated, solid-state package with a linear response in the oxygen concentration range of interest (10-30%). ZnO thin film is particularly touted as the active sensing material able to meet all abovementioned attributes, aside from exhibiting selectivity and long-term selectivity. EI

A95-69223

FEASIBILITY STUDY OF INFLATABLE STRUCTURES FOR A LUNAR BASE

PAUL S. NOWAK Colorado State Univ, Fort Collins, CO, United States, WILLY Z. SADEH, and JEFFREY JANAKUS Journal of Spacecraft and Rockets (ISSN 0022-4650) vol. 31, no. 3 May-June 1994 p. 453-457
(BTN-95-EIX95041503792) Copyright

The design of a structure on the moon requires addressing a host of issues not encountered on earth. A modular inflatable structure consisting of thin membranes of composite material integrated with supporting columns and arches is proposed. An initial linear analysis of the structure is briefly reviewed. The actual

response of an inflatable membrane is nonlinear and, hence, a nonlinear numerical analysis of the stresses and displacements was undertaken. Results indicate that an inflatable structure is a feasible concept for a lunar structure. Author (EI)

N95-19627* Kent State Univ., OH.

ON THE DEVELOPMENT OF AN EXPERT SYSTEM FOR WHEELCHAIR SELECTION

GREGORY R. MADEY, CHARLOTTE A. BHANSIN (Cleveland Clinic Foundation, OH.), SULAIMAN A. ALARAINI, and MOHAMED A. NOUR *In* NASA. Johnson Space Center, Third CLIPS Conference Proceedings, Volume 1 p 2-12 Nov. 1994
Avail: CASI HC A03/MF A03

The presentation of wheelchairs for the Multiple Sclerosis (MS) patients involves the examination of a number of complicated factors including ambulation status, length of diagnosis, and funding sources, to name a few. Consequently, only a few experts exist in this area. To aid medical therapists with the wheelchair selection decision, a prototype medical expert system (ES) was developed. This paper describes and discusses the steps of designing and developing the system, the experiences of the authors, and the lessons learned from working on this project. Wheelchair Advisor, programmed in CLIPS, serves as diagnosis, classification, prescription, and training tool in the MS field. Interviews, insurance letters, forms, and prototyping were used to gain knowledge regarding the wheelchair selection problem. Among the lessons learned are that evolutionary prototyping is superior to the conventional system development life-cycle (SDLC), the wheelchair selection is a good candidate for ES applications, and that ES can be applied to other similar medical subdomains. Author

55

SPACE BIOLOGY

Includes exobiology; planetary biology; and extraterrestrial life.

A95-66751* National Aeronautics and Space Administration, Washington, DC.

LIFE SCIENCES AND SPACE RESEARCH 24 (4): PLANETARY BIOLOGY AND ORIGINS OF LIFE; TOPICAL MEETING OF THE COSPAR INTERDISCIPLINARY SCIENTIFIC COMMISSION F (MEETING F3) OF THE COSPAR PLENARY MEETING, 29TH, WASHINGTON, DC, AUG. 28-SEP. 5, 1992

J. M. GREENBERG, editor Univ. of Leiden, Leiden, Netherlands, J. ORO, editor Univ. of Houston, Houston, TX, US, A. BRACK, editor CNRS, Orleans, France, D. L. DEVINCENZI, editor NASA. Ames Research Center, Moffett Field, CA, US, A. BANIN, editor Hebrew Univ., Rehovot, Israel, E. I. FRIEDMANN, editor Florida State Univ., Tallahassee, FL, US, J. D. RUMMEL, editor Marine Biological Lab., Woods Hole, MA, US, F. RAULIN, editor Univ. de Paris, Creteil, France, C. P. MCKAY, editor NASA. Ames Research Center, Moffett Field, CA, US, H. BALTSCHIEFFSKY, editor Stockholm Univ., Stockholm, Sweden et al. *Advances in Space Research* (ISSN 0273-1177) vol. 15, no. 3 March 1995 449 p. Research sponsored by NASA (ISBN 0-08-042540-2; HTN-95-10500) Copyright

The proceedings include sessions on extraterrestrial organic chemistry and the origins of life; life on Mars: past, present and future; planetary protection of Mars missions; chemical evolution on Titan; origins and early evolution of biological (a) energy transduction and membranes (b) information and catalysis; and carbon chemistry and isotopic fractionations in astrophysical environments. For individual titles, see A95-66752 through A95-66807. Hermer

A95-66752

CIRCUMSTELLAR CHEMISTRY FROM MICROWAVE AND MM-WAVE SPECTROSCOPY

J. H. BIEGING Univ. of Arizona, Tucson, AZ, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research* (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 3-14
Copyright

Stars in their late stages of evolution often shed matter in the form of a cool wind which is molecular in composition. These winds are a major source of replenishment of the interstellar gas and dust, so they furnish a large part of the raw materials for new generations of stars and planets. The chemistry of the circumstellar envelope depends strongly on the photospheric abundances of the elements, especially C and O. If C/O greater than 1, a rich organic chemistry is observable in the microwave and mm-wavelength emission lines of the reaction products. This paper reviews the observational evidence for the presence of organic molecules and their formation pathways in circumstellar envelopes, with special emphasis on rotational spectra at microwave and millimeter wavelengths. Author (Hermer)

A95-66764

ORIGIN OF AMINO ACIDS IN THE EARLY SOLAR SYSTEM

J. F. KERRIDGE UCLA, Los Angeles, CA, US *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research* (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 107-111
Copyright

Synthesis of meteoritic amino acids probably took place in the aqueous sub-surface regions of none or more asteroid-sized parent bodies. Starting material for those reactions apparently consisted of a population of more simple compounds including a suite of aliphatic precursors characterised by (1) complete structural diversity, (2) prevalence of branched- over straight-chain species, (3) exponential decrease in abundance with increasing C number, (4) large enrichment in D, and, probably, (5) systematic decrease in C-13/C-12 with increasing C number. Those properties were apparently acquired during synthesis of the precursors by ion-molecule reactions in a presolar molecular cloud. Author (Hermer)

A95-66766

FORMATION OF BIOORGANIC COMPOUNDS IN PLANETARY ATMOSPHERES BY COSMIC RADIATION

K. KOBAYASHI Yokohama National Univ., Hodogaya, Yokohama, Japan, T. KANEKO Yokohama National Univ., Hodogaya, Yokohama, Japan, M. TSUCHIYA Yokohama National Univ., Hodogaya, Yokohama, Japan, T. SAITO Univ. of Tokyo, Tanashi, Tokyo, Japan, T. YAMAMOTO Inst. of Space and Astronautical Science, Sagami, Japan, J. KOIKE Tokyo Inst. of Tech., Midori, Yokohama, Japan, and T. OSHIMA Tokyo Inst. of Tech., Midori, Yokohama, Japan *Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research* (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 127-130 Research sponsored by the Institute of Space and Astronautical Science
Copyright

Simulated planetary atmospheres (mixtures of simple gases) were irradiated with high energy particles to simulate an action of cosmic rays. When a mixture of carbon monoxide, nitrogen and water was irradiated with 2.8-40 MeV protons, a wide variety of bioorganic compounds including amino acids, imidazole, and uracil were identified in the products. The amount of amino acids was proportional to the energy deposit to the system. Various kinds of simulated planetary atmospheres, such as 'Titan type' and 'Jovian

type', were also irradiated with high energy protons, and gave amino acids in the hydrolyzed products. Since cosmic rays are a universal energy source in space, it was suggested that formation of bioorganic compounds in planetary atmospheres is inevitable in the course of cosmic evolution. Author (Herner)

A95-66768* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

EXO BIOLOGY RESEARCH ON SPACE STATION FREEDOM
J. L. HUNTINGTON NASA. Ames Research Center, Moffett Field, CA, US, D. M. STRATTON NASA. Ames Research Center, Moffett Field, CA, US, and T. W. SCATTERGOOD NASA. Ames Research Center, Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 135-138
Copyright

The Gas-Grain Simulation Facility (GGSF) is a multidisciplinary experiment laboratory being developed by NASA at Ames Research Center for delivery to Space Station Freedom in 1998. This facility will employ the low-gravity environment of the Space Station to enable aerosol experiments of much longer duration than is possible in any ground-based laboratory. Studies of fractal aggregates that are impossible to sustain on Earth will also be enabled. Three research areas within exobiology that will benefit from the GGSF are described here. An analysis of the needs of this research and of other suggested experiments has produced a list of science requirements which the facility design must accommodate. A GGSF design concept developed in the first stage of flight hardware development to meet these requirements is also described. Author (Herner)

A95-66770* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

EXO BIOLOGICAL EXPLORATION OF MARS
H. P. KLEIN Santa Clara Univ., Santa Clara, CA, US and D. L. DEVINCENZI NASA. Ames Research Center, Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 151-156
Copyright

Of all the other planets in the solar system, Mars remains the most promising for further elucidating concepts about chemical evolution and the origin of life. Strategies were developed to pursue three exobiological objectives for Mars exploration: determining the abundance and distribution of the biogenic elements and organic compounds, detecting evidence of an ancient biota on Mars, and determining whether indigenous organisms exist anywhere on the planet. The three strategies are quite similar and, in fact, share the same sequence of phases. In the first phase, each requires global reconnaissance and remote sensing by orbiters to select sites of interest for detailed in situ analyses. In the second phase, lander missions are conducted to characterize the chemical and physical properties of the selected sites. The third phase involves conducting 'critical' experiments at sites whose properties make them particularly attractive for exobiology. These critical experiments would include, for example, identification of organics, detection of fossils, and detection of extant life. The fourth phase is the detailed analysis of samples returned from these sites in Earth-based laboratories to confirm and extend previous discoveries. Finally, in the fifth phase, human exploration is needed to establish the geological settings for the earlier findings or to discover and explore sites that are not accessible to robotic spacecraft. Author (Herner)

A95-66771* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

SITE SELECTION FOR MARS EXO BIOLOGY

J. FARMER NASA. Ames Research Center, Moffett Field, CA, US, D. DES MARAIS NASA. Ames Research Center, Moffett Field, CA, US, R. GREELEY Arizona State Univ., Tempe, AZ, US, R. LANDHEIM Arizona State Univ., Tempe, AZ, US, and H. KLEIN Santa Clara Univ., Santa Clara, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 157-162 Research sponsored by NASA
Copyright

The selection of sites on Mars that have a high priority for exobiological research is fundamental for planning future exploration. The most immediate need is to identify targets for high resolution orbital imaging during the Mars Observer and Mars '94/'96 missions that can be used to refined site priorities for surface exploration. We present an objective approach to site selection whereby individual sites are selected and scored, based on the presence of key geological features which indicate high priority environments. Prime sites are those that show evidence for the prolonged activity of liquid water and which have sedimentary deposits that are likely to have accumulated in environments favorable for life. High priority areas include fluvio-lacustrine (stream-fed lake systems), springs, and periglacial environments. Sites where mineralization may have occurred in the presence of organisms (e.g. springs) are given high priority in the search for a fossil record on Mars. A systematic review of Viking data for 83 sites in the Mars Landing Site Catalog (MLSC) resulted in the selection of 13 as being of exobiological interest. The descriptions of these sites were expanded to address exobiological concerns. An additional five sites were identified for inclusion in the second edition of the MLSC. We plan to broaden our site selection activities to include a systematic global reconnaissance of Mars using Viking data, and will continue to refine site priorities for exobiological research based on data from future missions in order to define strategies for surface exploration. Author (Herner)

A95-66772* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

LIFE ON MARS? 1: THE CHEMICAL ENVIRONMENT
A. BANIN Hebrew Univ., Rehovot, Israel and R. L. MANCINELLI NASA. Ames Research Center, Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 163-170
Copyright

The origin of life at its abiotic evolutionary stage, requires a combination of constituents and environmental conditions that enable the synthesis of complex replicating macromolecules from simpler monomeric molecules. It is very likely that the early stages of this evolutionary process have been spontaneous, rapid and widespread on the surface of the primitive Earth, resulting in the formation of quite sophisticated living organisms within less than a billion years. To what extent did such conditions prevail on Mars? Two companion-papers will review and discuss the available information related to the chemical, physical and environmental conditions on Mars and assess it from the perspective of potential exobiological evolution. Author (Herner)

A95-66773* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

LIFE ON MARS? 2: PHYSICAL RESTRICTIONS
R. L. MANCINELLI NASA. Ames Research Center, Moffett Field, CA, US and A. BANIN Hebrew Univ., Rehovot, Israel Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th,

Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 171-176

Copyright

The primary physical factors important to life's evolution on a planet include its temperature, pressure and radiation regimes. Temperature and pressure regulate the presence and duration of liquid water on the surface of Mars. The prolonged presence of liquid water is essential for the evolution and sustained presence of life on a planet. It has been postulated that Mars has always been a cold dry planet; it has also been postulated that early Mars possessed a dense atmosphere of CO₂ (greater than or equal to 1 bar) and sufficient water to cut large channels across its surface. The degree to which either of these postulates is true correlates with the suitability of Mars for life's evolution. Although radiation can destroy living systems, the high fluxes of UV radiation on the martian surface do not necessarily stop the origin and early evolution of life. The probability for life to have arisen and evolved to a significant degree on Mars, based on the postulated ranges of early martian physical factors, is almost solely related to the probability of liquid water existing on the planet for at least hundreds of millions to billions of years.

Author (Hemer)

A95-66774

CHEMICAL STUDIES ON THE POSSIBLE EXISTENCE OF LIFE ON MARS

C. PONNAMPERUMA Univ. of Maryland, College Park, MD, US, R. NAVARRO-GONZALEZ Universidad Nacional Autonoma de Mexico, Mexico, Mexico, and Y. HONDA Naruto Univ. of Education, Naruto City, Japan Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 177-184

Copyright

Although there is no direct evidence yet for the existence of life on Mars, it is reasonable to conclude that the emergence of life on Earth, which appears to have been controlled by universal laws of physics and chemistry, may have been repeated elsewhere in the universe. The dual approach of synthesis and analysis in our experimental studies has provided ample evidence in support of this hypothesis.

Author (Hemer)

A95-66775

PHOSPHORUS AS A POTENTIAL GUIDE IN THE SEARCH FOR EXTINCT LIFE ON MARS

G. WECKWERTH Deutsche Forschungsanstalt fuer Luft-und Raumfahrt, Linder Hoehe, Koeln, Germany and M. SCHIDLOWSKI Max-Planck-Institut fuer Chemie, Mainz, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 185-191

Copyright

In contrast to the search for extant organisms, the quest for fossil remains of life on Mars need not be guided by the presence of water and organic compounds on the present surface. An appropriate tracer might be the element phosphorus which is a common constituent of living systems. Utilizing terrestrial analogues, it should preferentially exist in the form of sedimentary calcium phosphate (phosphorites), which would have readily resisted changing conditions on Mars. Moreover, higher ratios of Phosphorus/Thorium (P/Th) in phosphorites in comparison to calcium phosphates from magmatic rocks give us the possibility to distinguish them from inorganically formed phosphorus deposits at or close to the Martian surface. Identification of anomalous phosphorus enrichments by remote sensing or in situ analysis could be promising approaches for selecting areas preferentially composed of rocks with remains of extinct life.

Author (Hemer)

A95-66776* National Aeronautics and Space Administration. Ames Research Center, Moffett Field, CA.

SEARCH FOR LIFE ON MARS: EVALUATION OF TECHNIQUES

D. E. SCHWARTZ NASA. Ames Research Center, Moffett Field, CA, US, R. L. MANCINELLI NASA. Ames Research Center, Moffett Field, CA, US, and M. R. WHITE NASA. Ames Research Center, Moffett Field, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 193-197

Copyright

An important question for exobiology is, did life evolve on Mars? To answer this question, experiments must be conducted on the martian surface. Given current mission constraints on mass, power, and volume, these experiments can only be performed using proposed analytical techniques such as: electron microscopy, X-ray fluorescence, X-ray diffraction, alpha-proton backscatter, gamma-ray spectrometry, differential thermal analysis, differential scanning calorimetry, pyrolysis gas chromatography, mass spectrometry, and specific element detectors. Using prepared test samples consisting of 1% organic matter (bovine serum albumin) in palagonite and a mixture of palagonite, clays, iron oxides, and evaporites, it was determined that a combination of X-ray diffraction and differential thermal analysis coupled with gas chromatography provides the best insight into the chemistry, mineralogy, and geological history of the samples.

Author (Hemer)

A95-66782

A PHYSICAL AND CHEMICAL CHARACTERIZATION OF MARTIAN PERMAFROST AS A POSSIBLE HABITAT FOR VIABLE MICROORGANISMS

V. OSTROUMOV Russian Academy of Sciences, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 229-236

Copyright

Data from experiments with model samples show that ion transfer coefficients in the water-rich permafrost on Mars must be three orders of magnitude less than those of terrestrial permafrost. The effects of low temperatures and of carbon dioxide have been accounted for. Exchange between cells and the environment is impeded in Martian permafrost. The microscopic distributional heterogeneity of concentration, pH, Eh, and other physicochemical parameters may be more pronounced in the permafrost of Mars. We present a classification of unfrozen water types in the permafrost that is based on the structures of unfrozen water films and on their functions with respect to cells. Any viable microorganisms on Mars probably exists with minimum metabolism in compact zones with energy carriers and high transfer coefficients. These zones may be microvolumes of unfrozen water in which cells accumulate.

Author (Hemer)

A95-66785

MODERN ASPECTS OF PLANETARY PROTECTION AND REQUIREMENTS TO STERILIZATION OF SPACE HARDWARE

V. V. DEMIDOV Russian Academy of Sciences, Moscow, Russia, A. A. GONCHAROV Russian Academy of Sciences, Moscow, Russia, V. B. OSIPOV Scientific Industrial Association Biotechnology, Moscow, Russia, and V. I. TROFIMOV Scientific Industrial Association Biotechnology, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 251-255

Copyright

The viewpoint of a working group of Russian experts on the problem of planetary protection for future manned and unmanned

Mars mission is presented. Recent data on Martian environment and on survival of terrestrial microorganisms in extreme conditions were used for detailed analysis and overview of planetary protection measures in regard to all possible flight situations including accidental landing. The special emphasis on 'Mars-94' mission was done. This analysis resulted in revised formulation of spacecraft sterilization requirements and possible measures for their best implementation. New general combined approach to spacecraft sterilization is proposed. It includes penetrating radiation and heat treatment of spacecraft parts and components which is to be carried out before the final assembly of spacecraft and gaseous radiation sterilization of the whole spacecraft during the flight to Mars (or from Mars for return mission).

Author (Hemer)

A95-66786* Jet Propulsion Lab., California Inst. of Tech., Pasadena, CA.

DESIGNING PLANETARY PROTECTION INTO THE MARS OBSERVER MISSION

T. H. SWEETSER Jet Propulsion Lab., California Inst. of Tech., Pasadena, CA, US, C. A. HALSELL Jet Propulsion Lab., California Inst. of Tech., Pasadena, CA, US, and R. J. CESARONE Jet Propulsion Lab., California Inst. of Tech., Pasadena, CA, US Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 257-260 Research sponsored by NASA

Copyright

Planetary protection has been an important consideration during the process of designing the Mars Observer mission. It affected trajectory design of both the interplanetary transfer and the orbits at Mars; these in turn affected the observation strategies developed for the mission. The Project relied mainly on the strategy of collision avoidance to prevent contamination of Mars. Conservative estimates of spacecraft reliability and Martian atmosphere density were used to evaluate decisions concerning the interplanetary trajectory, the orbit insertion phase at Mars, and operations in orbit at Mars and afterwards. Changes in the trajectory design, especially in the orbit insertion phase, required a refinement of those estimates.

Author (Hemer)

A95-66787

PLANETARY ENVIRONMENT PROTECTION ID NO: F3.3 - M.1.05 IMPLICATIONS FOR THE DEVELOPMENT OF A NETWORK OF SURFACE STATIONS ON MARS

G. E. N. SCOON Directorate of Scientific Programmes, Noordwijk, Netherlands, A. CHICARRO Directorate of Scientific Programmes, Noordwijk, Netherlands, G. H. SCHWEHM Directorate of Scientific Programmes, Noordwijk, Netherlands, and F. SPIERO ESA ESTEC, Noordwijk, Netherlands Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 261-272

Copyright

The European Space Agency's (ESA) studies of a Comet Nucleus Sample Return mission (ROSETTA) as its Planetary Cornerstone in its long-term program 'Horizon 2000' and the Marsnet mission, a potential contribution of the Agency to an international network of surface stations on Mars, has revived the interest in the present state of Planetary Protection requirements. MARSNET was one of the four candidate missions selected in April 1991 for further Design Feasibility (Phase A) studies. Furthermore, of all space agencies participating in planetary exploration activities only the United States National Aeronautics and Space Administration (NASA) had a well established Planetary Protection Policy on Viking and other relevant planetary missions, whereas ESA is considering the feasibility and potential impact of a planetary protection policy on its Marsnet mission, within the framework of a tight budgetary envelope applicable to ESA's Medium (M) class missions. This paper will discuss in general terms the impact

of Planetary Protection measures, its implications for Marsnet and the issues arising from this for the implementation of the mission in ESA's scientific program.

Author (Hemer)

A95-66788

THE EXPERIMENTAL STUDY OF MICROBIAL CONTAMINATION OF THE SPACE HARDWARE

V. B. VASIN Scientific Industrial Association Biotechnology, Moscow, Russia and V. I. TROFIMOV Scientific Industrial Association Biotechnology, Moscow, Russia Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 273-276

Copyright

The role of potential contaminants of design materials and products of space technology - aerobic and anaerobic prokaryotes (myxobacteria, eubacteria, corinebacteria, actinomycetes), and eukaryotes (micromycetes), psychrophilic, mesophilic and thermophilic forms, and chemolithotrophic microorganisms is discussed in this paper. The methods of analysis of microbial contamination in the solution of problem of planetary protection are considered. The necessity of the use of ultrasound at the evaluation of surface and subsurface contamination of specimens is demonstrated; methods of determination of buried contamination (with the use of organic solvents and mechanical pulverization) are discussed. The data on buried and subsurface contamination for some materials and electronic parts together with microflora resistivity to sterilizing treatment are given.

Author (Hemer)

A95-66789

PLANETARY PROTECTION CONSIDERATIONS FOR MARSNET AND MARS SAMPLE RETURN MISSIONS

C. LINDBERG DLR, Linder Hoehe, Koeln, Germany and G. HORNECK DLR, Linder Hoehe, Koeln, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 277-280

Copyright

The European Space Agency (ESA) MarsNet mission proposal consists most probably of a trio of Mars landers. These landers each contain a variety of scientific equipment. The network of stations demands for a definition of its planetary protection requirements. With respect to the MarsNet mission only forward contamination problems will be considered. Future involvement of European efforts in planetary exploration including sample returns will also raise the problem of back contamination. A tradeoff study for the overall scientific benefit with respect to the approximative cost is necessary. Planetary protection guide-lines will be proposed by an interdisciplinary and international board of experts working in the fields of both biology and planetary science. These guide-lines will have to be flexible in order to be modified with respect to new research results, e. g. on adaptation of microorganisms to extreme (space) conditions. Experiments on the survival of microorganisms at conditions of simulated Mars surface and subsurface will have to be conducted in order to obtain a baseline data collection as a reference standard for future guide-lines.

Author (Hemer)

A95-66792

TITAN'S ATMOSPHERE COMPOSITION: CERTAINTIES AND SPECULATIONS

D. GAUTIER Observatoire de Paris-Meudon, Meudon, France Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 295-301

Copyright

Firm results concerning the thermal structure, the composition, the seasonal effects of the atmosphere of Titan, as well as the superrotation of its stratosphere are reviewed. The nature of the surface of the satellite, the possible presence of argon in the atmosphere and the structure and composition of clouds and aerosols are, among other topics, still speculative. The implications of the observed deuterium enrichment on the origin of ices in the outer part of the nebula are controversial. Author (Herner)

A95-66795

RAINDROPS ON TITAN

R. D. LORENZ Univ. of Kent, Canterbury, UK Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 317-320 Research sponsored by the UK Science and Engineering Research Council (SERC) and the Univ. of Kent Copyright

Some of the aspects of methane precipitation on Titan are considered. In particular, descent velocities are computed. It is found that raindrops fall much slower than on Earth. Additionally, the maximum size of raindrops on Titan is over 9mm, compared with under 6mm on Earth. The composition of drops will vary with altitude. Implications of these properties for Titan and the Huygens mission are considered. Author (Herner)

A95-66796

THE LOW TEMPERATURE ORGANIC CHEMISTRY OF TITAN'S GEOFLUID

F. RAULIN Univ. Paris - Val de Marne, Creteil, France, P. BRUSTON Univ. Paris - Val de Marne, Creteil, France, P. PAILLOUS Univ. Paris - Val de Marne, Creteil, France, and R. STERNBERG Univ. Paris - Val de Marne, Creteil, France Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 321-333 Research sponsored by the Centre National d'Etudes Spatiales and the Programme National de Planetologie Copyright

Organic chemistry on Titan and prebiotic chemistry on Earth involve the same N-containing organics: nitriles and their oligomers. Couplings of their chemistry in the three parts of Titan's geofluid (atmosphere, aerosols and surface) seem to play a key role in the organic chemical evolution of the planet. If liquid water was present on Titan, then a prebiotic chemistry, involving eutectics, similar to that of the early Earth, may have occurred. However, liquid water is currently absent and a prebiotic chemistry based only on N-organics may be evolving now on Titan. The other consequence of the low temperatures of Titan is the possible formation of organics unstable at room temperature and very reactive. So far, these compounds have not been systematically searched for in experimental studies of Titan's organic chemistry. C₄N₂ has already been detected on Titan. Powerful reactants in organic chemistry, CH₂N₂, and CH₃N₃, may be also present. They exhibit spectral signatures in the mid-IR strong enough to allow their detection at the 10-100 ppb level. They may be detectable on future IR spectra (ISO and Cassini) of Titan. Author (Herner)

A95-66801

PRODUCTION AND EVOLUTION OF CARBONACEOUS MATERIAL BY ION IRRADIATION IN SPACE

G. STRAZZULLA Univ. di Catania, Catania, Italy, G. A. BARATTA Citta Univ., Catania, Italy, and F. SPINELLA Citta Univ., Catania, Italy Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3

March 1995 p. 385-399 Research sponsored by the IDP Raman Spectra and the Italian Space Agency (ASI) Copyright

We review recent experimental studies concerning the evolution, driven by ion irradiation, of carbonaceous material from frozen gas to a refractory molecular solid. Under further irradiation the latter changes to a polymer-like material and ultimately to amorphous carbon. Most of the results have been obtained by 'in situ' and remote Infrared (IR) and Raman spectroscopy. The results have been applied to demonstrate that molecular solids may be easily formed by irradiation of frozen mantles in dense interstellar clouds. Polymer-like material and amorphous carbons may result by further irradiation of organic mantles on grains in the diffuse interstellar medium. Those grains, during the aggregation to form extended bodies like comets (T-Tau phase of the Sun), are further modified. These latter are also irradiated, after the comet formation, during their long stay in the Oort cloud. In particular it has been suggested that comet may develop an ion-produced cometary organic crust that laboratory evidences show to be stable against temperature increases experienced during passages near the Sun. The comparison between the Raman spectra of some IDP (Interplanetary Dust Particles) and the Raman spectra of some ion-produced amorphous carbons, is also discussed. Author (Herner)

A95-66803

CARBONACEOUS MATTER IN COMETARY DUST AND COMA

F. R. KRUEGER Engineering Office, Darmstadt, Germany Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 407-411 Copyright

The analysis of carbonaceous matter in p/Halley's dust and coma via mass spectrometry of positive ions is reviewed. Dust impact generated ions were analyzed by the PUMA instrument aboard VEGA 1, and coma plasma ions by the PICCA instrument aboard GIOTTO. For the organic molecules results an overall C:H:O:N ratio of 1.:1.4:0.6:0.1. Most of this polymer material can formally be understood as an aggregation of monomers C₂H₂, CH₂O, and HCN. Special emphasis is given to possible aromatic, especially heterocyclic, and other unsaturated ions, and their importance for abiotic chemical and prebiotic evolution. Aspects of the potential heterogeneous catalysis in liquid water at the inorganic grain backbone structure found by this analysis, too, are also treated. Author (Herner)


A95-66806

ISOTOPIC RATIOS IN PLANETARY ATMOSPHERES

C. DE BERGH Observatoire de Paris, Meudon, France Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992. A95-66751 Advances in Space Research (ISSN 0273-1177) vol. 15, no. 3 March 1995 p. 427-440 Copyright

Recent progress on measurements of isotopic ratios in planetary or satellite atmospheres include measurements of the Deuterium/Hydrogen (D/H) ratio in the methane of Uranus, Neptune and Titan and in the water of Mars and Venus. Implications of these measurements on our understanding of the formation and evolution of the planets and satellite are discussed. Our current knowledge of the carbon, nitrogen and oxygen isotopic ratios in the atmospheres of these planets, as well as on Jupiter and Saturn, is also reviewed. We finally show what progress can be expected in the very near future due to some new ground-based instrumentation particularly well suited to such studies, and to forthcoming space missions. Author (Herner)

Typical Subject Index Listing



The subject heading is a key to the subject content of the document. The title is used to provide a description of the subject matter. When the title is insufficiently descriptive of document content, a title extension is added, separated from the title by three hyphens. The accession number and the page number are included in each entry to assist the user in locating the abstract in the abstract section. If applicable, a report number is also included as an aid in identifying the document. Under any one subject heading, the accession numbers are arranged in sequence.

A

ABIOTENESIS
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Carbonaceous matter in cometary dust and coma p 218 A95-66803

ACETAZOLAMIDE
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration [HTN-95-A0124] p 201 A95-68907

ACETIC ACID
Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799

ACOUSTIC IMAGING
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714

ADENINES
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767

ADENOSINES
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767

ADIPOSE TISSUES
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women [HTN-95-A0120] p 200 A95-68903

ADRENERGENICS
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911

ADRENOCORTICOTROPIN (ACTH)
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

AEROEMBOLISM
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714

AEROSOLS
Exobiology research on Space Station Freedom p 215 A95-66768
The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796

AEROSPACE ENVIRONMENTS
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751

AEROSPACE MEDICINE
A multimodal method for assessing and treating airsickness [HTN-95-90719] p 209 A95-69746
Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746

AGE FACTOR
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911

AGGREGATES
The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794

AGING (BIOLOGY)
Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women [HTN-95-A0120] p 200 A95-68903

AIR FLOW
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure [HTN-95-A0138] p 204 A95-68921

AIR LAND INTERACTIONS
Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793

AIR POLLUTION
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895

AIR TRAFFIC CONTROL
Using knowledge exploration tools to study airspace complexity in air traffic control [HTN-95-90716] p 211 A95-69743

AIR TRAFFIC CONTROLLERS (PERSONNEL)
Using knowledge exploration tools to study airspace complexity in air traffic control [HTN-95-90716] p 211 A95-69743

AIR WATER INTERACTIONS
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897

AIRCRAFT DESIGN
Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744

AIRCRAFT PILOTS
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study [AD-A285050] p 212 N95-19564

AIRSPACE
Using knowledge exploration tools to study airspace complexity in air traffic control [HTN-95-90716] p 211 A95-69743

ALBUMINS
Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs [HTN-95-A0155] p 208 A95-68938

ALTITUDE ACCLIMATIZATION
Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923

ALTITUDE SICKNESS
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714

ALVEOLI
Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration [HTN-95-A0124] p 201 A95-68907
Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

AMINO ACIDS
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
Origin of amino acids in the early solar system p 214 A95-66764
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Extinction of dinosaurs: A possible novel cause p 191 A95-66769
Chemical studies on the possible existence of life on Mars p 216 A95-66774
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800

AMMONIA
Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902

ANALOGIES
Facilitation and interference in identification of pictures and words [AD-A285882] p 212 N95-19602

ANOXIA
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888

ANTARCTIC REGIONS
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777

ANTHROPOMETRY
Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929

ANTIADRENERGENICS
Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902

ARGON
Titan's atmosphere composition: Certainties and speculations p 217 A95-66792

ARID LANDS
Biochemical constraints for survival under martian conditions p 192 A95-66778

ARTERIES
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle [HTN-95-A0106] p 196 A95-68889
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree [HTN-95-A0121] p 200 A95-68904

ASPARTIC ACID

Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763

ASPERGILLUS

Biochemical constraints for survival under martian conditions p 192 A95-66778
Studies in the search for life on Mars p 192 A95-66779

ASTRONOMICAL SPECTROSCOPY

Laboratory astrophysics and molecular astronomy of pure carbon molecules p 189 A95-66753
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804
Isotopic ratios in planetary atmospheres p 218 A95-66806

ATMOSPHERIC CIRCULATION

Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263

ATMOSPHERIC COMPOSITION

Titan's atmosphere composition: Certainties and speculations p 217 A95-66792
The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794
Raindrops on Titan p 218 A95-66795
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797
Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263

ATMOSPHERIC PRESSURE

Life on Mars? 1: The chemical environment p 215 A95-66772
Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263

AUTOMATIC CONTROL

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742

AVIATION PSYCHOLOGY

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742
Using knowledge exploration tools to study airspace complexity in air traffic control [HTN-95-90716] p 211 A95-69743
Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745
A multimodal method for assessing and treating airsickness [HTN-95-90719] p 209 A95-69746

B

BACILLUS

Studies in the search for life on Mars p 192 A95-66779

BACTERIA

Biogeochemical evidence of microbial activity on Mars p 192 A95-66780
A primitive cyanobacterium as pioneer microorganism for terraforming Mars p 193 A95-66784

BARORECEPTORS

Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise [HTN-95-A0103] p 196 A95-68886
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901

BIBLIOGRAPHIES

Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746

BIOCHEMISTRY

Activities of the Division of Energy Biosciences [DE95-001046] p 208 N95-19613
Photobiology in medicine [AD-A284843] p 210 N95-19695

BIOGEOCHEMISTRY

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
Exobiological exploration of Mars p 215 A95-66770
Site selection for Mars exobiology p 215 A95-66771

Life on Mars? 2: Physical restrictions p 215 A95-66773

Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

BIOINSTRUMENTATION

Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems [BTN-94-EIX94501433007] p 195 A95-67426

BIOLOGICAL EVOLUTION

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800

BIOLOGICAL MODELS (MATHEMATICS)

Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree [HTN-95-A0121] p 200 A95-68904
Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

BIOMASS

Activities of the Division of Energy Biosciences [DE95-001046] p 208 N95-19613

BIOSPHERE

Cometary origin of the biosphere: A progress report p 189 A95-66757
Comets and life in the Universe p 190 A95-66761

BIOTECHNOLOGY

Activities of the Division of Energy Biosciences [DE95-001046] p 208 N95-19613

BLOOD

Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902

BLOOD CIRCULATION

Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698

BLOOD PRESSURE

Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898

BLOOD VESSELS

Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

BLOOD-BRAIN BARRIER

Hypoxia increases glucose transport at blood-brain barrier in rats [HTN-95-A0143] p 205 A95-68926

BLUE GREEN ALGAE

A 'crylic' microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781

BODY COMPOSITION (BIOLOGY)

Effects of strength training on total and regional body composition in older men p 199 A95-68899
Contribution of body composition and physical activity to age-related decline in peak dot-V_O2 in men and women [HTN-95-A0120] p 200 A95-68903
Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923
Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929

BODY FLUIDS

Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897

BOHR THEORY

Exercise O₂ transport model assuming zero cytochrome PO₂ at dot-V_O2(sub max) [HTN-95-A0122] p 200 A95-68905

BONE MARROW

Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading [HTN-95-A0113] p 198 A95-68896

BRAIN

Quantitative SPECT brain imaging: Effects of attenuation and detector response [BTN-94-EIX94361133479] p 209 A95-65976
Hypoxia increases glucose transport at blood-brain barrier in rats [HTN-95-A0143] p 205 A95-68926

BRAIN CIRCULATION

Hypoxia increases glucose transport at blood-brain barrier in rats [HTN-95-A0143] p 205 A95-68926

BREATHING

Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading [HTN-95-A0135] p 203 A95-68918
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs [HTN-95-A0144] p 205 A95-68927

BRONCHI

Serial distribution of airway mechanical properties in dogs: Effects of histamine [HTN-95-A0110] p 197 A95-68893

BUBBLES

Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897
Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714

C

C-135 AIRCRAFT

Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744

CALCIUM

Life on Mars? 2: Physical restrictions p 215 A95-66773
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934

CALCIUM METABOLISM

Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci [HTN-95-A0129] p 202 A95-68912

CALCIUM PHOSPHATES

Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775

CALCULI

Characterization of fracture toughness of renal calculi using a microindentation technique [BTN-94-EIX94361122293] p 209 A95-65873

CAPILLARIES

Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909
Interactive effects of emphysema and malnutrition on diaphragm structure and function [HTN-95-A0148] p 206 A95-68931
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume [HTN-95-A0153] p 208 A95-68936

CAPILLARIES (ANATOMY)

Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration [HTN-95-A0124] p 201 A95-68907
Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

CARBOHYDRATE METABOLISM

Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926

CARBON
Laboratory astrophysics and molecular astronomy of pure carbon molecules p 189 A95-66753
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765
Life on Mars? 2: Physical restrictions p 215 A95-66773
Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

CARBON COMPOUNDS
Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801

CARBON CYCLE
Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-66807

CARBON DIOXIDE
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle
[HTN-95-A0106] p 196 A95-68889
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration
[HTN-95-A0124] p 201 A95-68907
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

CARBON ISOTOPES
Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-66807

CARBON STARS
Laboratory astrophysics and molecular astronomy of pure carbon molecules p 189 A95-66753

CARBON 13
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763

CARBONACEOUS CHONDRITES
Cometary origin of the biosphere: A progress report p 189 A95-66757
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

CARBONIC ANHYDRASE
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration
[HTN-95-A0124] p 201 A95-68907

CARDIOGRAPHY
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891

CARDIOLOGY
Fluid flow and heat transfer in the crescent-shaped lumen catheter
[BTN-94-EIX94361122408] p 209 A95-65904

CARDIOVASCULAR SYSTEM
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg
[HTN-95-A0118] p 199 A95-68901
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936

CAROTID SINUS REFLEX
Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886

CASTOR OIL
Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909

CATALYSIS
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798

CATECHOLAMINE
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930

CATHETERIZATION
Fluid flow and heat transfer in the crescent-shaped lumen catheter
[BTN-94-EIX94361122408] p 209 A95-65904

CELL MEMBRANES (BIOLOGY)
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

CELLS (BIOLOGY)
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426

CEREBRUM
Cumulative effects of repeated brief cerebral ischemia
[AD-A284234] p 210 N95-19698

CHARACTER RECOGNITION
Facilitation and interference in identification of pictures and words
[AD-A285882] p 212 N95-19602

CHEMICAL COMPOSITION
Isotopic ratios in planetary atmospheres p 218 A95-66806

CHEMICAL EVOLUTION
Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762
Origin of amino acids in the early solar system p 214 A95-66764
Exobiological exploration of Mars p 215 A95-66770
Chemical studies on the possible existence of life on Mars p 216 A95-66774
Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775
Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799

CHEST
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925

CHIRAL DYNAMICS
Origin of amino acids in the early solar system p 214 A95-66764
Chemical studies on the possible existence of life on Mars p 216 A95-66774

CLIMATE CHANGE
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization
[HTN-95-60524] p 195 A95-68775

CLINICAL MEDICINE
A multimodal method for assessing and treating airsickness
[HTN-95-90719] p 209 A95-69746

CLOUDS (METEOROLOGY)
Titan's atmosphere composition: Certainties and speculations p 217 A95-66792
Raindrops on Titan p 218 A95-66795

COCKPITS
Use of workload redlines: A KC-135 crew-reduction application
[HTN-95-90717] p 211 A95-69744

CODING
Maximally smooth image recovery in transform coding
[BTN-94-EIX94361135047] p 209 A95-65949
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976

COGNITION
Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework
[AD-A285133] p 212 N95-19583

COGNITIVE PSYCHOLOGY
Cognitive representations of flight-deck information attributes
[HTN-95-90718] p 212 A95-69745
A multimodal method for assessing and treating airsickness
[HTN-95-90719] p 209 A95-69746
Comparing performance on implicit memory tests
[AD-A284861] p 212 N95-19740

COLD ACCLIMATIZATION
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910

COMET HEADS
Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758
Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759
Carbonaceous matter in cometary dust and coma p 218 A95-66803

COMET RENDEZVOUS ASTEROID FLYBY MISSION
Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758

COMETS
Cometary origin of the biosphere: A progress report p 189 A95-66757
Comets and life in the Universe p 190 A95-66761
Exobiology research on Space Station Freedom p 215 A95-66768
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802

COMPLEX SYSTEMS
Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX9504274623] p 213 A95-68277

COMPLEXITY
Using knowledge exploration tools to study airspace complexity in air traffic control
[HTN-95-90716] p 211 A95-69743

COMPUTER TECHNIQUES
High performance computing for medical image interpretation
[AD-A285457] p 210 N95-19537

COMPUTERIZED SIMULATION
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268

CONDUCTING POLYMERS
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426

CONFERENCES
Proceedings of Natural Sciences Symposium
[AD-A285452] p 208 N95-19536

CONTAMINATION
Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789

CONTAMINATION
The experimental study of microbial contamination of the space hardware p 217 A95-66788

CONTRACTION
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
Maximum rate of force development is increased by antagonist conditioning contraction p 203 A95-68917
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931

CONTROL BOARDS
Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System
[HTN-95-90715] p 211 A95-69742

CONVECTIVE HEAT TRANSFER
Fluid flow and heat transfer in the crescent-shaped lumen catheter
[BTN-94-EIX94361122408] p 209 A95-65904

CORE-MANTLE BOUNDARY
Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-66807

CORTICOSTEROIDS
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920

COSMIC DUST
Circumstellar chemistry from microwave and mm-wave spectroscopy p 214 A95-66752
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
Organic molecules in the gas phase of dense interstellar clouds p 189 A95-66755
Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762
Extinction of dinosaurs: A possible novel cause p 191 A95-66769
Carbonaceous matter in cometary dust and coma p 218 A95-66803
Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

COSMIC RAYS

Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization

[HTN-95-60524] p 195 A95-68775

CRETACEOUS-TERTIARY BOUNDARY

Extinction of dinosaurs: A possible novel cause p 191 A95-66769

CRUDE OIL

Activities of the Division of Energy Biosciences [DE95-001046] p 208 N95-19613

CUMULATIVE DAMAGE

Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698

CYCLIC HYDROCARBONS

Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759

CYTOCHROMES

Exercise O₂ transport model assuming zero cytochrome P₀₂ at dot-VO₂(sub max) [HTN-95-A0122] p 200 A95-68905

D

DATA BASE MANAGEMENT SYSTEMS

Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM [AD-A279193] p 211 N95-19811

DATA BASES

Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM [AD-A279193] p 211 N95-19811

DECISION MAKING

Using knowledge exploration tools to study airspace complexity in air traffic control [HTN-95-90716] p 211 A95-69743
Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538

DECOMPRESSION SICKNESS

Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714

DECONGESTANTS

Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898

DECONTAMINATION

The experimental study of microbial contamination of the space hardware p 217 A95-66788

DENSITOMETERS

Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929

DEOXYRIBONUCLEIC ACID

Biochemical constraints for survival under martian conditions p 192 A95-66778
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems [BTN-94-EIX94501433007] p 195 A95-67426

DESIGN ANALYSIS

Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744

DEUTERIUM

Titan's atmosphere composition: Certainties and speculations p 217 A95-66792

DIABETES MELLITUS

Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats [HTN-95-A0107] p 197 A95-68890

DIAPHRAGM (ANATOMY)

Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading [HTN-95-A0135] p 203 A95-68918
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans [HTN-95-A0141] p 205 A95-68924
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs [HTN-95-A0144] p 205 A95-68927
Interactive effects of emphysema and malnutrition on diaphragm structure and function [HTN-95-A0148] p 206 A95-68931

DIFFUSION

Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888

DIVING (UNDERWATER)

Tissue oxygen and carbon dioxide stores and breath-hold diving in humans [HTN-95-A0108] p 197 A95-68891

DOSAGE

Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898

DROP SIZE

Raindrops on Titan p 218 A95-66795

DROPS (LIQUIDS)

Raindrops on Titan p 218 A95-66795

E

EARTH ENVIRONMENT

Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-68807
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775

ELECTRIC SPARKS

Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763

ELECTRIC STIMULI

High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist [HTN-95-A0152] p 207 A95-68935

ELECTROCARDIOGRAPHY

Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise [HTN-95-A0103] p 196 A95-68886
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933

ELECTROENCEPHALOGRAPHY

Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study [AD-A285050] p 212 N95-19564

ENDOTHELIUM

Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle [HTN-95-A0106] p 196 A95-68889

ENDOTOXINS

Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894
Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs [HTN-95-A0155] p 208 A95-68938

ENVIRONMENT MODELS

Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767

ENVIRONMENT SIMULATION

Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

ENVIRONMENT SIMULATORS

Studies in the search for life on Mars p 192 A95-66779

ENVIRONMENTAL ENGINEERING

Annual historical report, calendar year 1993 [AD-A280178] p 211 N95-19827

ENZYME ACTIVITY

On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
Myosin isoforms in mammalian skeletal muscle [HTN-95-A0102] p 195 A95-68885
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle [HTN-95-A0106] p 196 A95-68889
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression [HTN-95-A0117] p 199 A95-68900
Interactive effects of emphysema and malnutrition on diaphragm structure and function [HTN-95-A0148] p 206 A95-68931

ENZYMES

Myosin isoforms in mammalian skeletal muscle [HTN-95-A0102] p 195 A95-68885

EPINEPHRINE

Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

ESOPHAGUS

Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs [HTN-95-A0136] p 204 A95-68919

ETIOLOGY

A multimodal method for assessing and treating airsickness [HTN-95-90719] p 209 A95-69746

EXERCISE PHYSIOLOGY

Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women [HTN-95-A0120] p 200 A95-68903
Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937

EXO BIOLOGY

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
Comets and life in the Universe p 190 A95-66761
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
Exobiology research on Space Station Freedom p 215 A95-66768
Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746

EXPERT SYSTEMS

On the development of an expert system for wheelchair selection p 214 N95-19627

EXPIRED AIR

Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895

EXPOSURE

Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM [AD-A279193] p 211 N95-19811

EXTINCTION

Extinction of dinosaurs: A possible novel cause p 191 A95-66769

EXTRATERRESTRIAL ENVIRONMENTS

Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263

F

FATTY ACIDS

Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

FEASIBILITY ANALYSIS

Feasibility study of inflatable structures for a lunar base [BTN-95-EIX95041503792] p 213 A95-69223

FIBERS

Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle [HTN-95-A0133] p 203 A95-68916
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

FINGERS

Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist [HTN-95-A0152] p 207 A95-68935

FINITE DIFFERENCE THEORY

Fluid flow and heat transfer in the crescent-shaped lumen catheter [BTN-94-EIX94361122408] p 209 A95-65904

FLIGHT CONTROL

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742

FLIGHT CREWS

Use of workload redlines: A KC-135 crew-reduction application
[HTN-95-90717] p 211 A95-69744

FLIGHT MANAGEMENT SYSTEMS

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System
[HTN-95-90715] p 211 A95-69742

FLIGHT SIMULATION

Use of workload redlines: A KC-135 crew-reduction application
[HTN-95-90717] p 211 A95-69744

FLOW GEOMETRY

Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease
[HTN-95-A0150] p 207 A95-68933

FLUID FILMS

Surface activity of lipid extract surfactant in relation to film area compression and collapse
[HTN-95-A0151] p 207 A95-68934

FLUID FLOW

Fluid flow and heat transfer in the crescent-shaped lumen catheter
[BTN-94-EIX94361122408] p 209 A95-65904

FOOD INTAKE

Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929

FORMALDEHYDE

Possible evidence of hydrocarbons released by the nucleus of Halley's comet
p 190 A95-66759
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices
p 194 A95-66802

FRACTIONATION

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751

FRACTURE STRENGTH

Characterization of fracture toughness of renal calculi using a microindentation technique
[BTN-94-EIX94361122293] p 209 A95-65873

FRAGMENTS

Facilitation and interference in identification of pictures and words
[AD-A285882] p 212 N95-19602

FREE RADICALS

Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918

G**GAS CHROMATOGRAPHY**

Search for life on Mars: Evaluation of techniques
p 216 A95-66776

GAS DETECTORS

Oxygen sensor development for life support
[BTN-95-EIX95042477107] p 213 A95-68352

GAS EXCHANGE

Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928

GENE EXPRESSION

Myosin isoforms in mammalian skeletal muscle
[HTN-95-A0102] p 195 A95-68885

GENETIC CODE

Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code
p 194 A95-66800

GENETIC ENGINEERING

Proceedings of Natural Sciences Symposium
[AD-A285452] p 208 N95-19536

GEOCHEMISTRY

A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms
p 216 A95-66782
Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars
p 217 A95-66787

GLUCOSE

Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats
[HTN-95-A0107] p 197 A95-68890
Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration
[HTN-95-A0119] p 200 A95-68902
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926

GLYCOGENS

High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation
[HTN-95-A0149] p 207 A95-68932

GRAVITATIONAL PHYSIOLOGY

Cumulative effects of repeated brief cerebral ischemia
[AD-A284234] p 210 N95-19698

H**HALLEY'S COMET**

Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future
p 190 A95-66758
Possible evidence of hydrocarbons released by the nucleus of Halley's comet
p 190 A95-66759

HARDNESS TESTS

Characterization of fracture toughness of renal calculi using a microindentation technique
[BTN-94-EIX94361122293] p 209 A95-65873

HEALTH

Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811

HEART RATE

Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans
[HTN-95-A0115] p 199 A95-68898
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner
[HTN-95-A0154] p 208 A95-68937

HELICOPTERS

Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564

HEMODYNAMICS

Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg
[HTN-95-A0118] p 199 A95-68901
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree
[HTN-95-A0121] p 200 A95-68904
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease
[HTN-95-A0150] p 207 A95-68933

HEMOGLOBIN

Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892
Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906

HIGH ALTITUDE BREATHING

Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906

HIGH ALTITUDE ENVIRONMENTS

Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923

HIGH GRAVITY ENVIRONMENTS

Cumulative effects of repeated brief cerebral ischemia
[AD-A284234] p 210 N95-19698

HIGH PRESSURE OXYGEN

Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

HISTOCHEMICAL ANALYSIS

Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931

HUMAN FACTORS ENGINEERING

On the development of an expert system for wheelchair selection
p 214 N95-19627

HUMAN PERFORMANCE

Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX95042474623] p 213 A95-68277
Learning effects on strategy selection in a dynamic task environment as a function of time pressure
[AD-A285458] p 212 N95-19538
Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework
[AD-A285133] p 212 N95-19583

HUMAN-COMPUTER INTERFACE

Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX95042474623] p 213 A95-68277

HYDROCARBONS

Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future
p 190 A95-66758

Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth
p 193 A95-66793

The physical nature of Titan's aerosols: Laboratory simulations
p 193 A95-66794

Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission
p 193 A95-66797

HYDROCYANIC ACID

Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry
p 190 A95-66760

Chemical studies on the possible existence of life on Mars
p 216 A95-66774

Role of trace metal ions in chemical evolution. The case of free-radical reactions
p 194 A95-66799

HYDROXYL COMPOUNDS

Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite
p 190 A95-66762

HYPERBARIC CHAMBERS

Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927

HYPERCAPNIA

Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919

HYPEROXIA

Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

HYPERTENSION

Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912

HYPERTHERMIA

Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892

HYPOXIA

Developmental differences in vascular responses to hypoxia in lungs of rabbits
[HTN-95-A0104] p 196 A95-68887

Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910

Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912

Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926

- ICE**
Titan's atmosphere composition: Certainties and speculations p 217 A95-66792
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802
- ICE ENVIRONMENTS**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
- IMAGE PROCESSING**
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949
High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537
- IMAGE RECONSTRUCTION**
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949
- IMAGE RESOLUTION**
High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537
- IMAGING TECHNIQUES**
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949
Quantitative SPECT brain imaging: Effects of attenuation and detector response [BTN-94-EIX94361133479] p 209 A95-65976
High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537
- IMMUNE SYSTEMS**
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading [HTN-95-A0113] p 198 A95-68896
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression [HTN-95-A0117] p 199 A95-68900
- IMPEDANCE**
Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908
- INFLATABLE STRUCTURES**
Feasibility study of inflatable structures for a lunar base [BTN-95-EIX95041503792] p 213 A95-69223
- INFORMATION MANAGEMENT**
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745
- INFORMATION PROCESSING (BIOLOGY)**
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745
Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538
Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework [AD-A285133] p 212 N95-19583
- INFRARED SPECTROSCOPY**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
- INJECTION**
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898
- INJURIES**
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894
- INSULIN**
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats [HTN-95-A0107] p 197 A95-68890
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920
- INTERNATIONAL COOPERATION**
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775
- INTERPLANETARY FLIGHT**
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268
- INTERSTELLAR GAS**
Organic molecules in the gas phase of dense interstellar clouds p 189 A95-66755
- INTERSTELLAR MATTER**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
- Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804
Isotopic ratios in planetary atmospheres p 218 A95-66806
- INTERSTELLAR RADIATION**
Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801
- INTRAVASCULAR SYSTEM**
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714
- IONIZING RADIATION**
Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268
- ISCHEMIA**
Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698
- ISOMORPHISM**
Myosin isoforms in mammalian skeletal muscle [HTN-95-A0102] p 195 A95-68885
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression [HTN-95-A0117] p 199 A95-68900
- J**
- JUPITER (PLANET)**
Isotopic ratios in planetary atmospheres p 218 A95-66806
- K**
- KINETIC ENERGY**
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
- KNEE (ANATOMY)**
Maximum rate of force development is increased by antagonist conditioning contraction [HTN-95-A0134] p 203 A95-68917
- L**
- LACTATES**
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920
- LACTIC ACID**
Exercise O₂ transport model assuming zero cytochrome P₀₂ at dot-VO₂(sub max) [HTN-95-A0122] p 200 A95-68905
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906
- LAKES**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
- LANDING MODULES**
Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785
- LANGMUIR-BLODGETT FILMS**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems [BTN-94-EIX94501433007] p 195 A95-67426
- LARYNX**
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure [HTN-95-A0138] p 204 A95-68921
- LEARNING**
Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538
- LEARNING THEORY**
Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538
- LEG (ANATOMY)**
(31P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937
- LIFE SCIENCES**
Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746
- LIFE SUPPORT SYSTEMS**
Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263
Oxygen sensor development for life support [BTN-95-EIX95042477107] p 213 A95-68352
Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746
- LIGHT (VISIBLE RADIATION)**
Photobiology in medicine [AD-A284843] p 210 N95-19695
- LIPIDS**
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897
Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934
- LOADS (FORCES)**
Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930
- LOCAL AREA NETWORKS**
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM [AD-A279193] p 211 N95-19811
- LONG DURATION SPACE FLIGHT**
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268
- LOW TEMPERATURE ENVIRONMENTS**
The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802
- LOWER BODY NEGATIVE PRESSURE**
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901
- LUNAR BASES**
Atmospheric pressure within lunar structure [BTN-95-EIX95032452326] p 213 A95-68263
Design and construction of shielded lunar outpost [BTN-95-EIX95032452328] p 213 A95-68265
Feasibility study of inflatable structures for a lunar base [BTN-95-EIX95041503792] p 213 A95-69223
- LUNAR ENVIRONMENT**
Design and construction of shielded lunar outpost [BTN-95-EIX95032452328] p 213 A95-68265
- LUNAR SHELTERS**
Design and construction of shielded lunar outpost [BTN-95-EIX95032452328] p 213 A95-68265
- LUNAR SURFACE**
Design and construction of shielded lunar outpost [BTN-95-EIX95032452328] p 213 A95-68265
- LUNG MORPHOLOGY**
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree [HTN-95-A0121] p 200 A95-68904
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure [HTN-95-A0138] p 204 A95-68921
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans [HTN-95-A0141] p 205 A95-68924
- LUNGS**
Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897
Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908
Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909
Pulmonary vascular resistance distribution and recruitment of microvascular surface area [HTN-95-A0139] p 204 A95-68922

Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925

Lung volumes and expiratory flow limitation during exercise in interstitial lung disease
[HTN-95-A0150] p 207 A95-68933

Surface activity of lipid extract surfactant in relation to film area compression and collapse
[HTN-95-A0151] p 207 A95-68934

Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936

Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs
[HTN-95-A0155] p 208 A95-68938

LYMPHOCYTES
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading
[HTN-95-A0113] p 198 A95-68896

M

MACROPHAGES
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading
[HTN-95-A0113] p 198 A95-68896

MAGNESIUM
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805

Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912

MAGNETIC STORMS
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization
[HTN-95-60524] p 195 A95-68775

MAGNETITE
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

MAN MACHINE SYSTEMS
Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX95042474623] p 213 A95-68277

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System
[HTN-95-90715] p 211 A95-69742

Use of workload redlines: A KC-135 crew-reduction application
[HTN-95-90717] p 211 A95-69744

Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397)
[NASA-SP-7011(397)] p 210 N95-19746

MARINE BIOLOGY
A 'crytic' microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781

MARS (PLANET)
Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751

Exobiological exploration of Mars p 215 A95-66770

Site selection for Mars exobiology p 215 A95-66771

Life on Mars? 1: The chemical environment p 215 A95-66772

Life on Mars? 2: Physical restrictions p 215 A95-66773

Chemical studies on the possible existence of life on Mars p 216 A95-66774

Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775

Search for life on Mars: Evaluation of techniques p 216 A95-66776

Biochemical constraints for survival under martian conditions p 192 A95-66778

Studies in the search for life on Mars p 192 A95-66779

Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

A 'crytic' microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781

A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782

Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

A primitive cyanobacterium as pioneer microorganism for terraforming Mars p 193 A95-66784

Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785

Designing planetary protection into the Mars observer mission p 217 A95-66786

Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

Isotopic ratios in planetary atmospheres p 218 A95-66806

Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268

MARS ATMOSPHERE
Exobiological exploration of Mars p 215 A95-66770

Life on Mars? 1: The chemical environment p 215 A95-66772

Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

MARS ENVIRONMENT
Life on Mars? 1: The chemical environment p 215 A95-66772

A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777

Biochemical constraints for survival under martian conditions p 192 A95-66778

Studies in the search for life on Mars p 192 A95-66779

A 'crytic' microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781

A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782

Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

A primitive cyanobacterium as pioneer microorganism for terraforming Mars p 193 A95-66784

MARS EXPLORATION
Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751

Exobiological exploration of Mars p 215 A95-66770

Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785

Designing planetary protection into the Mars observer mission p 217 A95-66786

Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789

Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268

MARS LANDING
Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

MARS PHOTOGRAPHS
Site selection for Mars exobiology p 215 A95-66771

MARS PROBES
Exobiological exploration of Mars p 215 A95-66770

MARS SAMPLE RETURN MISSIONS
Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789

MARS SURFACE
Exobiological exploration of Mars p 215 A95-66770

Site selection for Mars exobiology p 215 A95-66771

Life on Mars? 2: Physical restrictions p 215 A95-66773

Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

MARS SURFACE SAMPLES
Search for life on Mars: Evaluation of techniques p 216 A95-66776

MASS SPECTROSCOPY
Search for life on Mars: Evaluation of techniques p 216 A95-66776

MECHANICAL DEVICES
Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263

MEDICAL EQUIPMENT
Fluid flow and heat transfer in the crescent-shaped lumen catheter
[BTN-94-EIX94361122408] p 209 A95-65904

MEDICAL SCIENCE
Photobiology in medicine
[AD-A284843] p 210 N95-19695

Annual historical report, calendar year 1993
[AD-A280178] p 211 N95-19827

MEDICAL SERVICES
High performance computing for medical image interpretation
[AD-A285457] p 210 N95-19537

On the development of an expert system for wheelchair selection p 214 N95-19627

Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811

Annual historical report, calendar year 1993
[AD-A280178] p 211 N95-19827

MEMORY
Facilitation and interference in identification of pictures and words
[AD-A285882] p 212 N95-19602

Comparing performance on implicit memory tests
[AD-A284861] p 212 N95-19740

MENTAL PERFORMANCE
Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System
[HTN-95-90715] p 211 A95-69742

Cognitive representations of flight-deck information attributes
[HTN-95-90718] p 212 A95-69745

Comparing performance on implicit memory tests
[AD-A284861] p 212 N95-19740

METABOLISM
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798

Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923

Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929

Cumulative effects of repeated brief cerebral ischemia
[AD-A284234] p 210 N95-19698

METAL OXIDE SEMICONDUCTORS
Oxygen sensor development for life support
[BTN-95-EIX95042477107] p 213 A95-68352

METEORITES
Comets and life in the Universe p 190 A95-66761

Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

METHANE
Raindrops on Titan p 218 A95-66795

The low temperature organic chemistry of Titan's geofluid p 218 A95-66796

Isotopic ratios in planetary atmospheres p 218 A95-66806

MICROHARDNESS
Characterization of fracture toughness of renal calculi using a microindentation technique
[BTN-94-EIX94361122293] p 209 A95-65873

MICROORGANISMS
Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751

A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782

Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

The experimental study of microbial contamination of the space hardware p 217 A95-66788

Activities of the Division of Energy Biosciences
[DE95-001046] p 208 N95-19613

MILITARY OPERATIONS
Annual historical report, calendar year 1993
[AD-A280178] p 211 N95-19827

MILITARY TECHNOLOGY
Proceedings of Natural Sciences Symposium
[AD-A285452] p 208 N95-19536

MINERALS
Search for life on Mars: Evaluation of techniques p 216 A95-66776

MOISTURE CONTENT
Regional lung hematocrit variation and assessment of acute lung injury
[HTN-95-A0111] p 198 A95-68894

Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923

MOLECULAR BIOLOGY
Photobiology in medicine
[AD-A284843] p 210 N95-19695

MOLECULAR CLOUDS
Organic molecules in the gas phase of dense interstellar clouds p 189 A95-66755

MOON

- Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- MOTION SICKNESS**
A multimodal method for assessing and treating airsickness
[HTN-95-90719] p 209 A95-69746
- MURCHISON METEORITE**
Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry p 190 A95-66760
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
Origin of amino acids in the early solar system p 214 A95-66764

MUSCLE RELAXANTS

- Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920

MUSCLES

- Myosin isoforms in mammalian skeletal muscle
[HTN-95-A0102] p 195 A95-68885
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle
[HTN-95-A0106] p 196 A95-68889
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
Maximum rate of force development is increased by antagonist conditioning contraction
[HTN-95-A0134] p 203 A95-68917
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation
[HTN-95-A0149] p 207 A95-68932
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner
[HTN-95-A0154] p 208 A95-68937

MUSCULAR FATIGUE

- Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist
[HTN-95-A0152] p 207 A95-68935

MUSCULAR FUNCTION

- Myosin isoforms in mammalian skeletal muscle
[HTN-95-A0102] p 195 A95-68885
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
Maximum rate of force development is increased by antagonist conditioning contraction
[HTN-95-A0134] p 203 A95-68917
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation
[HTN-95-A0149] p 207 A95-68932

MUSCULAR STRENGTH

- Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
Lung volume specificity of inspiratory muscle training
[HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training
[HTN-95-A0132] p 203 A95-68915

- Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist
[HTN-95-A0152] p 207 A95-68935

MUSCULOSKELETAL SYSTEM

- High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation
[HTN-95-A0149] p 207 A95-68932
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist
[HTN-95-A0152] p 207 A95-68935

MYOCARDIUM

- Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age
[HTN-95-A0128] p 202 A95-68911

N

NAPHTHALENE

- Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804

NECK (ANATOMY)

- Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886

NEPTUNE (PLANET)

- Isotopic ratios in planetary atmospheres p 218 A95-66806

NEUROMUSCULAR TRANSMISSION

- Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916

NITRIC OXIDE

- Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle
[HTN-95-A0106] p 196 A95-68889
Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892

NITRILES

- Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

NITROGEN

- Life on Mars? 2: Physical restrictions p 215 A95-66773

NITROGEN METABOLISM

- Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923

NOREPINEPHRINE

- Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930

NUCLEOTIDES

- Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800

O

OBESITY

- Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats
[HTN-95-A0107] p 197 A95-68890

OORT CLOUD

- Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801

OPERATIONAL HAZARDS

- Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265

ORGANIC CHEMISTRY

- Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
- ORGANIC COMPOUNDS**
Comets and life in the Universe p 190 A95-66761
Exobiology research on Space Station Freedom p 215 A95-66768
Carbonaceous matter in cometary dust and coma p 218 A95-66803

OXIDATION-REDUCTION REACTIONS

- Exercise O2 transport model assuming zero cytochrome PO2 at dot-VO2(sub max)
[HTN-95-A0122] p 200 A95-68905

OXYGEN

- Oxygen sensor development for life support
[BTN-95-EIX95042477107] p 213 A95-68352

OXYGEN CONSUMPTION

- Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891
Contribution of body composition and physical activity to age-related decline in peak dot-VO2 in men and women
[HTN-95-A0120] p 200 A95-68903
Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age
[HTN-95-A0128] p 202 A95-68911
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner
[HTN-95-A0154] p 208 A95-68937
- OXYGEN METABOLISM**
Exercise O2 transport model assuming zero cytochrome PO2 at dot-VO2(sub max)
[HTN-95-A0122] p 200 A95-68905
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- OZONE**
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow
[HTN-95-A0112] p 198 A95-68895

P

PACKET SWITCHING

- Maximally smooth image recovery in transform coding
[BTN-94-EIX94361135047] p 209 A95-65949

PARALYSIS

- Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925

PATTERN RECOGNITION

- Facilitation and interference in identification of pictures and words
[AD-A265882] p 212 N95-19602

PERFORMANCE TESTS

- Facilitation and interference in identification of pictures and words
[AD-A265882] p 212 N95-19602

PERMAFROST

- A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

PERMEABILITY

- Regional lung hematocrit variation and assessment of acute lung injury
[HTN-95-A0111] p 198 A95-68894
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs
[HTN-95-A0155] p 208 A95-68938

PHENANTHRENE

- Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804

PHOSPHORUS

Life on Mars? 2: Physical restrictions p 215 A95-66773
 Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775

PHOSPHORUS ISOTOPES

(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937

PHOTOCHEMICAL REACTIONS

Photobiology in medicine [AD-A284843] p 210 N95-19695

PHYSICAL EXERCISE

Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906
 Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911
 Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920
 Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923
 Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929
 Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930
 Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933
 (31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937

PHYSICAL FITNESS

Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899
 Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902
 Contribution of body composition and physical activity to age-related decline in peak dot-VO2 in men and women [HTN-95-A0120] p 200 A95-68903
 Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911
 Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914
 Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915
 Maximum rate of force development is increased by antagonist conditioning contraction [HTN-95-A0134] p 203 A95-68917
 Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928

PHYSICAL WORK

Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895
 Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902
 Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle [HTN-95-A0133] p 203 A95-68916
 Maximum rate of force development is increased by antagonist conditioning contraction [HTN-95-A0134] p 203 A95-68917
 Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume [HTN-95-A0153] p 208 A95-68936

PHYSIOLOGICAL FACTORS

Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698
 Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746

PHYSIOLOGICAL RESPONSES

Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268

PILOT PERFORMANCE

Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742
 Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744

Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745

PLANETARY ATMOSPHERES

Formation of biogenic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
 Isotopic ratios in planetary atmospheres p 218 A95-66806

PLANETARY EVOLUTION

Comets and life in the Universe p 190 A95-66761
 Isotopic ratios in planetary atmospheres p 218 A95-66806

PLANETARY MAPPING

Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

PLANETARY ORBITS

Designing planetary protection into the Mars observer mission p 217 A95-66786

PLANETARY QUARANTINE

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28, Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
 Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785

Designing planetary protection into the Mars observer mission p 217 A95-66786
 Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789

POLAR CAPS

Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

POLYMERIC FILMS

Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems [BTN-94-EIX94501433007] p 195 A95-67426

POLYMERS

Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry p 190 A95-66760

POTASSIUM

Life on Mars? 2: Physical restrictions p 215 A95-66773

PREDICTIONS

The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775

PRESSURE BREATHING

Respiratory response to inhaled CO2 during positive inspiratory pressure in humans [HTN-95-A0141] p 205 A95-68924

PRESSURE SENSORS

Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909

PRIMITIVE EARTH ATMOSPHERE

Cometary origin of the biosphere: A progress report p 189 A95-66757
 Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793

PROKARYOTES

Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

PROSTAGLANDINS

Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887

PSYCHOLOGICAL TESTS

Comparing performance on implicit memory tests [AD-A284861] p 212 N95-19740

PSYCHOLOGY

Facilitation and interference in identification of pictures and words [AD-A285882] p 212 N95-19602

PULMONARY CIRCULATION

Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887

Tissue oxygen and carbon dioxide stores and breath-hold diving in humans [HTN-95-A0108] p 197 A95-68891

Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood [HTN-95-A0109] p 197 A95-68892

Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree [HTN-95-A0121] p 200 A95-68904

Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909

Pulmonary vascular resistance distribution and recruitment of microvascular surface area [HTN-95-A0139] p 204 A95-68922

Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928

Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume [HTN-95-A0153] p 208 A95-68936

Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs [HTN-95-A0155] p 208 A95-68938

PULMONARY FUNCTIONS

Tissue oxygen and carbon dioxide stores and breath-hold diving in humans [HTN-95-A0108] p 197 A95-68891

Serial distribution of airway mechanical properties in dogs: Effects of histamine [HTN-95-A0110] p 197 A95-68893

Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895

Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914

Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915

Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs [HTN-95-A0136] p 204 A95-68919

Chest wall interrupter resistance in anesthetized paralyzed humans [HTN-95-A0142] p 205 A95-68925

Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs [HTN-95-A0144] p 205 A95-68927

Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928

Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933

PYRENES

Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804

PYROLYSIS

Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

PYRUVATES

Exercise O2 transport model assuming zero cytochrome PO2 at dot-VO2(sub max) [HTN-95-A0122] p 200 A95-68905

Q

QUINONES

Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading [HTN-95-A0135] p 203 A95-68918

R

RADIATION COUNTERS

Quantitative SPECT brain imaging: Effects of attenuation and detector response [BTN-94-EIX94361133479] p 209 A95-65976

RADIATION DOSAGE

Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268

RADIATION EFFECTS

Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
 Photobiology in medicine [AD-A284843] p 210 N95-19695

RADIATION HAZARDS

The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775

RADIATION PROTECTION

- Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268
- RADIATION SHIELDING**
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- RADIATION TOLERANCE**
Life on Mars? 1: The chemical environment
p 215 A95-66772
- RADIO TELEMETRY**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564
- RANDOM PROCESSES**
Proceedings of Natural Sciences Symposium
[AD-A285452] p 208 N95-19536
- REAL TIME OPERATION**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811
- REBREATHING**
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936
- RED GIANT STARS**
Circumstellar chemistry from microwave and mm-wave spectroscopy
p 214 A95-66752
- REFLEXES**
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans
[HTN-95-A0115] p 199 A95-68898
- REFRACTORY PERIOD**
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
- REGRESSION ANALYSIS**
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree
[HTN-95-A0121] p 200 A95-68904
- RELIABILITY ANALYSIS**
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- REPTILES**
Extinction of dinosaurs: A possible novel cause
p 191 A95-66769
- RESEARCH AND DEVELOPMENT**
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811
Annual historical report, calendar year 1993
[AD-A280178] p 211 N95-19827
- RESEARCH MANAGEMENT**
Proceedings of Natural Sciences Symposium
[AD-A285452] p 208 N95-19536
- RESISTANCE**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
- RESPIRATION**
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891
Serial distribution of airway mechanical properties in dogs: Effects of histamine
[HTN-95-A0110] p 197 A95-68893
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
- RESPIRATORY IMPEDANCE**
Serial distribution of airway mechanical properties in dogs: Effects of histamine
[HTN-95-A0110] p 197 A95-68893
Pressure-flow specificity of inspiratory muscle training
[HTN-95-A0132] p 203 A95-68915
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
- RESPIRATORY PHYSIOLOGY**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

- Lung volume specificity of inspiratory muscle training
[HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training
[HTN-95-A0132] p 203 A95-68915
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928
- RESUSCITATION**
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans
[HTN-95-A0141] p 205 A95-68924
- RISK**
Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX95042474623] p 213 A95-68277
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811
- RUNNING**
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-1a(cp) rats
[HTN-95-A0107] p 197 A95-68890

S

- SATELLITE ATMOSPHERES**
Isotopic ratios in planetary atmospheres
p 218 A95-66806
- SATELLITE SURFACES**
The low temperature organic chemistry of Titan's geofluid
p 218 A95-66796
- SATURN (PLANET)**
Isotopic ratios in planetary atmospheres
p 218 A95-66806
- SEDIMENTS**
Phosphorus as a potential guide in the search for extinct life on Mars
p 216 A95-66775
Biogeochemical evidence of microbial activity on Mars
p 192 A95-66780
A 'crytic' microbial mat: A new model ecosystem for extant life on Mars
p 192 A95-66781
A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms
p 216 A95-66782
Preservation of cell structures in permafrost: A model for exobiology
p 192 A95-66783
- SELECTION**
On the development of an expert system for wheelchair selection
p 214 N95-19627
- SEMICONDUCTING FILMS**
Oxygen sensor development for life support
[BTN-95-EIX95042477107] p 213 A95-68352
- SHIVERING**
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910
- SIZE DETERMINATION**
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness
[AD-A284849] p 210 N95-19714
- SKIN (ANATOMY)**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- SLEEP**
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure
[HTN-95-A0138] p 204 A95-68921
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927
- SOCKS**
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg
[HTN-95-A0118] p 199 A95-68901
- SOILS**
Life on Mars? 2: Physical restrictions
p 215 A95-66773
Search for life on Mars: Evaluation of techniques
p 216 A95-66776
- SOLAR SYSTEM EVOLUTION**
Circumstellar chemistry from microwave and mm-wave spectroscopy
p 214 A95-66752
- SOLAR TERRESTRIAL INTERACTIONS**
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization
[HTN-95-60524] p 195 A95-68775
- SOLAR WIND**
Circumstellar chemistry from microwave and mm-wave spectroscopy
p 214 A95-66752
- SPACE ENVIRONMENT SIMULATION**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805
- SPACE STATION FREEDOM**
Exobiology research on Space Station Freedom
p 215 A95-66768
- SPACECRAFT**
The experimental study of microbial contamination of the space hardware
p 217 A95-66788
- SPACECRAFT STERILIZATION**
Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
The experimental study of microbial contamination of the space hardware
p 217 A95-66788
- STANDARD DEVIATION**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- STAPHYLOCOCCUS**
Studies in the search for life on Mars
p 192 A95-66779
- STELLAR COMPOSITION**
Laboratory astrophysics and molecular astronomy of pure carbon molecules
p 189 A95-66753
- STELLAR ENVELOPES**
Organic molecules in the gas phase of dense interstellar clouds
p 189 A95-66755
- STELLAR EVOLUTION**
Circumstellar chemistry from microwave and mm-wave spectroscopy
p 214 A95-66752
- STEREOCHEMISTRY**
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite
p 190 A95-66763
- STREPTOCOCCUS**
Studies in the search for life on Mars
p 192 A95-66779
Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912
- STRESS (PHYSIOLOGY)**
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg
[HTN-95-A0118] p 199 A95-68901
Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926
- STRUCTURAL DESIGN**
Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
Feasibility study of inflatable structures for a lunar base
[BTN-95-EIX95041503792] p 213 A95-69223
- STRUCTURAL ENGINEERING**
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- SUN**
Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268
- SURFACE PROPERTIES**
Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs
[HTN-95-A0155] p 208 A95-68938
- SURFACTANTS**
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble
[HTN-95-A0114] p 198 A95-68897
Surface activity of lipid extract surfactant in relation to film area compression and collapse
[HTN-95-A0151] p 207 A95-68934

SYNTHESIS (CHEMISTRY)

Organic analysis of hydrogen cyanide polymers:
Prebiotic and extraterrestrial chemistry

p 190 A95-66760

Were micrometeorites a source of prebiotic molecules
on the early Earth?

p 191 A95-66765

SYNTHETIC FUELS

Activities of the Division of Energy Biosciences

[DE95-001046] p 208 N95-19613

SYSTEM FAILURES

Minimizing human-machine interface failures in high risk
systems

[BTN-95-EIX95042474623] p 213 A95-68277

T**TASK COMPLEXITY**

Multiple-task performance: A critical review of the
literature and a cognitive neuroscience framework

[AD-A285133] p 212 N95-19583

TECHNETIUM

Regional lung hematocrit variation and assessment of
acute lung injury

[HTN-95-A0111] p 198 A95-68894

TEMPERATURE DEPENDENCE

Life on Mars? 1: The chemical environment

p 215 A95-66772

TEMPERATURE EFFECTS

Radiation and thermal stabilities of adenine
nucleotides

p 191 A95-66767

TERRAFORMING

A primitive cyanobacterium as pioneer microorganism
for terraforming Mars

p 193 A95-66784

THERMAL ANALYSIS

Search for life on Mars: Evaluation of techniques

p 216 A95-66776

THERMOREGULATION

Hypoxia similarly impairs metabolic responses to
cutaneous and core cold stimuli in conscious rats

[HTN-95-A0127] p 201 A95-68910

THIN FILMS

Oxygen sensor development for life support

[BTN-95-EIX95042477107] p 213 A95-68352

THREE DIMENSIONAL MODELS

High performance computing for medical image
interpretation

[AD-A285457] p 210 N95-19537

THYROID GLAND

Interaction of thyroid hormone and functional overload
on skeletal muscle isomyosin expression

[HTN-95-A0117] p 199 A95-68900

TIME DEPENDENCE

Learning effects on strategy selection in a dynamic task
environment as a function of time pressure

[AD-A285458] p 212 N95-19538

TITAN

Life sciences and space research 24 (4): Planetary
biology and origins of life: Topical Meeting of the COSPAR
Interdisciplinary Scientific Commission F (Meeting F3) of
the COSPAR Plenary Meeting, 29th, Washington, DC, Aug.
28-Sep. 5, 1992

[ISBN 0-08-042540-2] p 214 A95-66751

Titan's atmosphere composition: Certainties and
speculations

p 217 A95-66792

Surface-atmosphere interactions on Titan compared
with those on the pre-biotic Earth

p 193 A95-66793

The physical nature of Titan's aerosols: Laboratory
simulations

p 193 A95-66794

Raindrops on Titan

p 218 A95-66795

The low temperature organic chemistry of Titan's
geofluid

p 218 A95-66796

Analytical pyrolysis experiments of Titan aerosol
analogues in preparation for the Cassini Huygens
mission

p 193 A95-66797

TOXIC DISEASES

Extinction of dinosaurs: A possible novel cause

p 191 A95-66769

TRACE ELEMENTS

Chemical studies on the possible existence of life on
Mars

p 216 A95-66774

TRACHEA

Serial distribution of airway mechanical properties in
dogs: Effects of histamine

[HTN-95-A0110] p 197 A95-68893

Detection of free radicals by electron spin resonance
in rat diaphragm after resistive loading

[HTN-95-A0135] p 203 A95-68918

TRANSMISSION LOSS

Maximally smooth image recovery in transform coding

[BTN-94-EIX94361135047] p 209 A95-65949

U**ULTRASONIC TESTS**

Echo imaging techniques determine the size of
intravascular bubbles in decompression sickness

[AD-A284849] p 210 N95-19714

URANUS (PLANET)

Isotopic ratios in planetary atmospheres

p 218 A95-66806

V**VASOCONSTRICTION**

Developmental differences in vascular responses to
hypoxia in lungs of rabbits

[HTN-95-A0104] p 196 A95-68887

VASODILATION

Nitric oxide-independent response to acetylcholine by
terminal arterioles in rat cremaster muscle

[HTN-95-A0106] p 196 A95-68889

VENTILATION

Influence of lung volume dependence of upper airway
resistance during continuous negative airway pressure

[HTN-95-A0138] p 204 A95-68921

VENUS (PLANET)

Isotopic ratios in planetary atmospheres

p 218 A95-66806

VIABILITY

Biochemical constraints for survival under marian
conditions

p 192 A95-66778

Studies in the search for life on Mars

p 192 A95-66779

VIRUSES

Studies in the search for life on Mars

p 192 A95-66779

VISUAL PERCEPTION

Facilitation and interference in identification of pictures
and words

[AD-A285882] p 212 N95-19602

W**WATER**

Life on Mars? 2: Physical restrictions

p 215 A95-66773

WEIGHT REDUCTION

Lower limit of body fat in healthy active men

[HTN-95-A0146] p 206 A95-68929

WEIGHTLESSNESS SIMULATION

Cosmic dust analog simulation in a microgravity
environment: The STARDUST program

p 195 A95-66805

Interleukin-2 therapy reverses some
immunosuppressive effects of skeletal unloading

[HTN-95-A0113] p 198 A95-68896

WHEELCHAIRS

On the development of an expert system for wheelchair
selection

p 214 N95-19627

WORDS (LANGUAGE)

Facilitation and interference in identification of pictures
and words

[AD-A285882] p 212 N95-19602

WORKLOADS (PSYCHOPHYSIOLOGY)

Use of workload redlines: A KC-135 crew-reduction
application

[HTN-95-90717] p 211 A95-69744

Multiple-task performance: A critical review of the
literature and a cognitive neuroscience framework

[AD-A285133] p 212 N95-19583

Z**ZINC**

Cosmic dust analog simulation in a microgravity
environment: The STARDUST program


p 195 A95-66805

PERSONAL AUTHOR INDEX

AEROSPACE MEDICINE AND BIOLOGY / A Continuing Bibliography (Supplement 401)

May 1995

Typical Personal Author Index Listing



Listings in this index are arranged alphabetically by personal author. The title of the document is used to provide a brief description of the subject matter. The report number helps to indicate the type of document (e.g., NASA report, translation, NASA contractor report). The page and accession numbers are located beneath and to the right of the title. Under any one author's name the accession numbers are arranged in sequence.

A

- ABBOT, NEIL C.**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- ABDEL-RAHMAN, A. R. A.**
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans
[HTN-95-A0115] p 199 A95-68898
- ADES, PHILIP A.**
Contribution of body composition and physical activity to age-related decline in peak dot-V_{O2} in men and women
[HTN-95-A0120] p 200 A95-68903
- AHMAIDI, S.**
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
- ALARAINI, SULAIMAN A.**
On the development of an expert system for wheelchair selection
p 214 N95-19627
- ALLAMANDOLA, L. J.**
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices
p 194 A95-66802
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands
p 194 A95-66804
- ANDERSON, D. T.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments
p 191 A95-66777
- ANDERSON, MARK E.**
Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912
- ARMSTRONG, JASON W.**
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading
[HTN-95-A0113] p 198 A95-68896
- ASKEW, E. WAYNE**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929

B

- BAER, P. R.**
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner
[HTN-95-A0154] p 208 A95-68937
- BALCH, SIGNE**
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading
[HTN-95-A0113] p 198 A95-68896
- BALDWIN, KENNETH M.**
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
- BALTSCHIEFFSKY, H.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
- BANIN, A.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
Life on Mars? 1: The chemical environment
p 215 A95-66772
Life on Mars? 2: Physical restrictions
p 215 A95-66773
- BARATTA, G. A.**
Production and evolution of carbonaceous material by ion irradiation in space
p 218 A95-66801
- BATES, JASON H. T.**
Serial distribution of airway mechanical properties in dogs: Effects of histamine
[HTN-95-A0110] p 197 A95-68893
- BAUDRY, NATHALIE**
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle
[HTN-95-A0106] p 196 A95-68889
- BECK, J. SWANSON**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- BEHMANN, RAMIN A.**
Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926
- BEN-JEBRIA, ABDELLAZIZ**
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow
[HTN-95-A0112] p 198 A95-68895
- BERLINER, L.**
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
- BERNATH, P. F.**
Laboratory astrophysics and molecular astronomy of pure carbon molecules
p 189 A95-66753
- BHANSIN, CHARLOTTE A.**
On the development of an expert system for wheelchair selection
p 214 N95-19627
- BHATTACHARYA, JAHAR**
Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909
- BHATTACHARYA, SUNITA**
Diameters of juxtacapillary venules determined by oil-drop method in rat lung
[HTN-95-A0126] p 201 A95-68909
- BIDANI, AKHIL**
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration
[HTN-95-A0124] p 201 A95-68907
- BIEGER-DOSE, A.**
Biochemical constraints for survival under martian conditions
p 192 A95-66778

- BIEGING, J. H.**
Circumstellar chemistry from microwave and mm-wave spectroscopy
p 214 A95-66752
- BISHOP, B.**
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927
- BOLSHIYANOV, D.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments
p 191 A95-66777
- BOON, J. J.**
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission
p 193 A95-66797
- BORZONE, G.**
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
- BOWSER, J. E.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- BRACK, A.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
Were micrometeorites a source of prebiotic molecules on the early Earth?
p 191 A95-66765
- BRANT, ROLLIN**
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910
- BROWER, ROY G.**
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
- BRUSTON, P.**
The low temperature organic chemistry of Titan's geofluid
p 218 A95-66796
- BUETTNER, GARRY R.**
Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892
- BURNETTE, TARA M.**
Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912
- BUSSOLETTI, E.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805

C

- CAIOZZO, VINCENT J.**
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
- CALDERINI, E.**
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
- CALDWELL, JOHN A., JR.**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564
- CARDINAL, DONNA**
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811
- CARNOCHAN, FIONA M. T.**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913

AUTHOR

CAROTENUTO, L.

Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

CESARONE, R. J.

Designing planetary protection into the Mars observer mission p 217 A95-66786

CHANG, SHIH-WEN

Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs [HTN-95-A0155] p 208 A95-68938

CHAPES, STEPHEN K.

Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading [HTN-95-A0113] p 198 A95-68896

CHEN, QIU-HONG

Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906

CHERYCH, N.

A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777

CHESLEY, ALAN

High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

CHEUNG, JEFFERY T.

Oxygen sensor development for life support [BTN-95-EIX95042477107] p 213 A95-68352

CHICARRO, A.

Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

CHUONG, C. J.

Characterization of fracture toughness of renal calculi using a microindentation technique [BTN-94-EIX94361122293] p 209 A95-65873

CLAIREMIDI, J.

Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759

CLANTON, T. L.

Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading [HTN-95-A0135] p 203 A95-68918

CLEMENS, R. E.

Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933

CLEMENTS, J. A.

Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble [HTN-95-A0114] p 198 A95-68897

CLERE, JEAN-MICHEL

Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901

COHEN, MARK E.

Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915

COLANGELI, L.

Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

COLLOMP, K.

Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

COMMANDEUR, J.

Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

CONGER, K. A.

Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698

COOPER, G. W.

Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762

COOPER, S.

Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

COTE, M. G.

Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

CRONIN, J. R.

Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762

CURSTEDT, TORE

Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934

D

D'ANGELO, E.

Chest wall interrupter resistance in anesthetized paralyzed humans [HTN-95-A0142] p 205 A95-68925

DALLAL, MOHAMMED M.

Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs [HTN-95-A0155] p 208 A95-68938

DARLING, STEPHEN R.

Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study [AD-A285050] p 212 N95-19564

DAWSON, CHRISTOPHER A.

Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree [HTN-95-A0121] p 200 A95-68904

DE BERGH, C.

Isotopic ratios in planetary atmospheres p 218 A95-66806

DE BOER, R. W.

(31P)-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937

DELL'AVERSANA, P.

Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

DELSEMMER, A. H.

Cometary origin of the biosphere: A progress report p 189 A95-66757

DEMIDOV, V. V.

Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785

DES MARAIS, D.

Site selection for Mars exobiology p 215 A95-66771

DEVINCENZI, D. L.

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
Exobiological exploration of Mars p 215 A95-66770

DIAZ-VILLAGOMEZ, E.

On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798

DILLARD, ROBERT M.

Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study [AD-A285050] p 212 N95-19564

DILLMANN, R.

Biochemical constraints for survival under martian conditions p 192 A95-66778

DORAN, P.

A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777

DORMAN, L. I.

The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775

DOSE, K.

Biochemical constraints for survival under martian conditions p 192 A95-66778

DUMAY, A. C.

High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537

DYCK, DAVID J.

High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

E

EBADIAN, M. A.

Fluid flow and heat transfer in the crescent-shaped lumen catheter [BTN-94-EIX94361122408] p 209 A95-65904

EHRENFREUND, P.

Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

ENGEL, M. H.

Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763

ENGRAND, C.

Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

ESCOURROU, P.

Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise [HTN-95-A0103] p 196 A95-68886

F

FARMER, J.

Site selection for Mars exobiology p 215 A95-66771

FERGUSON, F.

Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

FESSLER, HENRY E.

Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs [HTN-95-A0136] p 204 A95-68919

FEWELL, JAMES E.

Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats [HTN-95-A0127] p 201 A95-68910

FIKE, CANDICE D.

Developmental differences in vascular responses to hypoxia in lungs of rabbits [HTN-95-A0104] p 196 A95-68887

FLECK, S. J.

Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930

FORTIER, M.

Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

FRIEDL, KARL E.

Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929

FRIEDMANN, E. I.

Life sciences and space research 24 (4): Planetary biology and origins of life: Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751

FRIEDMANN, E. IMRE

A primitive cyanobacterium as pioneer microorganism for terraforming Mars p 193 A95-66784

FRY, A. C.

Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930

FUJIMOTO, KEISAKU

Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906

FUJISHIMA, SEITARO

Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

FULAMBARKER, ASHOK M.

Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915

G

GALCHENKO, V.

A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777

GALLAGHER, C. G.

Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933

GAN, R. Z.

Vascular impedance analysis in dog lung with detailed morphometric and elasticity data [HTN-95-A0125] p 201 A95-68908

GAO, J.

Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats [HTN-95-A0107] p 197 A95-68890

H

- GARDNER, ANDREW W.**
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women
[HTN-95-A0120] p 200 A95-68903
- GARSHNEK, V.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past marian lacustrine environments p 191 A95-66777
- GAUTHIER, J. P.**
Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886
- GAUTIER, D.**
Titan's atmosphere composition: Certainties and speculations p 217 A95-66792
- GE, RI-LI**
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
- GEISER, DENNIS R.**
Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912
- GEN, DING**
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
- GIBBS, JAMES H.**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- GIESBRECHT, GORDON G.**
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910
- GILICHINSKY, D. A.**
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783
- GILLAND, D. R.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- GISOLFI, CARL V.**
Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892
- GLENNY, ROBB W.**
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928
- GOERKE, J.**
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble
[HTN-95-A0114] p 198 A95-68897
- GOLDBERG, A. P.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- GONCHAROV, A. A.**
Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785
- GORDON, CLAIRE C.**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929
- GRABINER, MARK D.**
Maximum rate of force development is increased by antagonist conditioning contraction
[HTN-95-A0134] p 203 A95-68917
- GRAHAM, TERRY E.**
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
- GREELEY, R.**
Site selection for Mars exobiology p 215 A95-66771
- GREENBERG, J. M.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
- GREER, K. L.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- GUTTMAN, JEREMY A.**
Using knowledge exploration tools to study airspace complexity in air traffic control
[HTN-95-90716] p 211 A95-69743
- HABIB, ROBERT H.**
Serial distribution of airway mechanical properties in dogs: Effects of histamine
[HTN-95-A0110] p 197 A95-68893
- HADDAD, FADIA**
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
- HALL, DAVID M.**
Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-heme in blood
[HTN-95-A0109] p 197 A95-68892
- HALSELL, C. A.**
Designing planetary protection into the Mars observer mission p 217 A95-66786
- HAPPEL, J.**
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- HARF, A.**
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans
[HTN-95-A0141] p 205 A95-68924
- HARIK, SAMI I.**
Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926
- HARMAN, S. M.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- HARRIS, T. R.**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
- HARRISON, DAVID K.**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- HASAN, S. U.**
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927
- HASEGAWA, NOKI**
Regional lung hematocrit variation and assessment of acute lung injury
[HTN-95-A0111] p 198 A95-68894
- HASSOUN, JOHN A.**
Use of workload redlines: A KC-135 crew-reduction application
[HTN-95-90717] p 211 A95-69744
- HEATH, J. R.**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
- HEMING, THOMAS A.**
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration
[HTN-95-A0124] p 201 A95-68907
- HERAZO, L. F.**
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936
- HERRICK, ROBERT E.**
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
- HESPEL, PETER J. L.**
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle
[HTN-95-A0105] p 196 A95-68888
- HETHERINGTON, H. P.**
Cumulative effects of repeated brief cerebral ischemia
[AD-A284234] p 210 N95-19698
- HOBISH, M. K.**
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800
- HONDA, Y.**
Chemical studies on the possible existence of life on Mars p 216 A95-66774
- HOPKINS, SUSAN R.**
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928
- HORNECK, G.**
Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789
- HOU, XIN**
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle
[HTN-95-A0106] p 196 A95-68889
- HOYT, REED W.**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929
- HSIA, C. C. W.**
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936
- HU, SHU-CHIEH**
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow
[HTN-95-A0112] p 198 A95-68895
- HUNTINGTON, J. L.**
Exobiology research on Space Station Freedom p 215 A95-66768
- HURLEY, B. F.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- HWANG, H. J.**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754

I

- IRVINE, W. M.**
Organic molecules in the gas phase of dense interstellar clouds p 189 A95-66755
- ISABEY, D.**
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans
[HTN-95-A0141] p 205 A95-68924
- IUCCI, N.**
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization
[HTN-95-60524] p 195 A95-68775
- IVANOV, M. V.**
Biogeochemical evidence of microbial activity on Mars p 192 A95-66780

J

- JACKSON, ANDREW C.**
Serial distribution of airway mechanical properties in dogs: Effects of histamine
[HTN-95-A0110] p 197 A95-68893
- JACKSON, R. JEFFREY**
A multimodal method for assessing and treating airsickness
[HTN-95-90719] p 209 A95-69746
- JAMES, PHILIP B.**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- JANAKUS, JEFFREY**
Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263
Feasibility study of inflatable structures for a lunar base
[BTN-95-EIX95041503792] p 213 A95-69223
- JANJINDAMAI, WARICHA**
Magnesium attenuates pulmonary hypertension due to hypoxia and group B streptococci
[HTN-95-A0129] p 202 A95-68912
- JASZCZAK, R. J.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- JENESON, J. A. L.**
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner
[HTN-95-A0154] p 208 A95-68937
- JOHNSON, PARLEY P.**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564
- JOHNSON, R. L., JR.**
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936
- JOHNSON, SCOTT R.**
Oxygen sensor development for life support
[BTN-95-EIX95042477107] p 213 A95-68352
- JOLLY, S. D.**
Design and construction of shielded lunar outpost
[BTN-95-EIX95032452328] p 213 A95-68265
- JONSSON, JON E.**
Cognitive representations of flight-deck information attributes
[HTN-95-90718] p 212 A95-69745

K

- KAMATH, MANJUNATH**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426
- KANAZAWA, MINORU**
Regional lung hematocrit variation and assessment of acute lung injury
[HTN-95-A0111] p 198 A95-68894
- KANEKO, T.**
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
- KAPLOWITZ, MARK R.**
Developmental differences in vascular responses to hypoxia in lungs of rabbits
[HTN-95-A0104] p 196 A95-68887
- KAWAGOE, YASHUHIRO**
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
- KAWASAKI, Y.**
Studies in the search for life on Mars p 192 A95-66779
- KAYSER, BENGT**
Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923
- KERNELL, D.**
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist
[HTN-95-A0152] p 207 A95-68935
- KERRIDGE, J. F.**
Origin of amino acids in the early solar system p 214 A95-66764
- KERSTHOLT, J. H.**
Learning effects on strategy selection in a dynamic task environment as a function of time pressure
[AD-A285458] p 212 N95-19538
- KHARE, B.**
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797
- KISSEL, J.**
Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758
- KLEIN, H.**
Site selection for Mars exobiology p 215 A95-66771
- KLEIN, H. P.**
Exobiological exploration of Mars p 215 A95-66770
- KOBAYASHI, K.**
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Studies in the search for life on Mars p 192 A95-66779
- KOIKE, J.**
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Studies in the search for life on Mars p 192 A95-66779
- KORTELING, J. E.**
Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework
[AD-A285133] p 212 N95-19583
- KOZIRIS, L. P.**
Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930
- KRAEMER, W. J.**
Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930
- KRENZ, GARY S.**
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree
[HTN-95-A0121] p 200 A95-68904
- KRUEGER, F. R.**
Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758
Carbonaceous matter in cometary dust and coma p 218 A95-66803
- KUBO, KEISHI**
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
- KUO, E. W.**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
- KURAT, G.**
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

L

- LAMANNA, JOSEPH C.**
Hypoxia increases glucose transport at blood-brain barrier in rats
[HTN-95-A0143] p 205 A95-68926
- LANDHEIM, R.**
Site selection for Mars exobiology p 215 A95-66771
- LANGE, F.**
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle
[HTN-95-A0133] p 203 A95-68916
- LAZCANO, A.**
Comets and life in the Universe p 190 A95-66761
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
- LE TRONG, JEAN-LOUIS**
Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923
- LEIN, A. YU.**
Biogeochemical evidence of microbial activity on Mars p 192 A95-66780
- LEWIS, JAMES A.**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study
[AD-A285050] p 212 N95-19564
- LEWIS, MICHAEL I.**
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931
- LIANG, Z.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response
[BTN-94-EIX94361133479] p 209 A95-65976
- LIEBMAN, S. A.**
Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry p 190 A95-66760
- LILLELEHT, L. U.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805
- LIM, JEONG OK**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426
- LIN, JIANMING**
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree
[HTN-95-A0121] p 200 A95-68904
- LINDBERG, C.**
Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789
- LINEHAN, JOHN H.**
Impact of parallel heterogeneity on a continuum model of the pulmonary arterial tree
[HTN-95-A0121] p 200 A95-68904
- LINER, MATS H.**
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891
- LINNARSSON, DAG**
Tissue oxygen and carbon dioxide stores and breath-hold diving in humans
[HTN-95-A0108] p 197 A95-68891
- LLOYD, DAVID**
Extinction of dinosaurs: A possible novel cause p 191 A95-66769
- LOFASO, F.**
Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans
[HTN-95-A0141] p 205 A95-68924
- LOMANGINO, NIOMI A.**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
- LONG, A.**
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
- LORENZ, R. D.**
Raindrops on Titan p 218 A95-66795
- LOWE, J. GRAHAM**
Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation
[HTN-95-A0130] p 202 A95-68913
- LUNINE, J. I.**
Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793

LYNCH, J. M.

- Catecholamine responses to short-term high-intensity resistance exercise overtraining
[HTN-95-A0147] p 206 A95-68930

M

- MACKO, S. A.**
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
- MADEY, GREGORY R.**
On the development of an expert system for wheelchair selection p 214 N95-19627
- MANCINELLI, R. L.**
Life on Mars? 1: The chemical environment p 215 A95-66772
Life on Mars? 2: Physical restrictions p 215 A95-66773
Search for life on Mars: Evaluation of techniques p 216 A95-66776
- MARC, I.**
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure
[HTN-95-A0138] p 204 A95-68921
- MARCHITELLI, LOUIS J.**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929
- MARCINIUK, D. D.**
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease
[HTN-95-A0150] p 207 A95-68933
- MARIETTA, MARTIN**
Minimizing human-machine interface failures in high risk systems
[BTN-95-EIX95042474623] p 213 A95-68277
- MARTINEZ-LOPEZ, LESTER E.**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929
- MARX, KENNETH A.**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426
- MATSUZAWA, YUKINORI**
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude
[HTN-95-A0123] p 201 A95-68906
- MATTHES, RONALD D.**
Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of (dot)NO-herme in blood
[HTN-95-A0109] p 197 A95-68892
- MATTHEWS, C. N.**
Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry p 190 A95-66760
- MAURETTE, M.**
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765
- MCCOOL, F. DENNIS**
Lung volume specificity of inspiratory muscle training
[HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training
[HTN-95-A0132] p 203 A95-68915
- MCCUE, SAMUEL A.**
Interaction of thyroid hormone and functional overload on skeletal muscle isomyosin expression
[HTN-95-A0117] p 199 A95-68900
- MCCUNE, S. A.**
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fa(cp) rats
[HTN-95-A0107] p 197 A95-68890
- MCKAY, C. P.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Planetary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793
- MCKENZIE, DONALD C.**
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation
[HTN-95-A0145] p 206 A95-68928
- MEGIRIAN, DAVID**
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910

- MELCHIOR, FRANCOIS M.**
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg
[HTN-95-A0118] p 199 A95-68901
- MELE, F.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805
- MENNELLA, V.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805
- MEROLA, A. J.**
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading
[HTN-95-A0135] p 203 A95-68918
- MERRILL, R. H.**
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans
[HTN-95-A0115] p 199 A95-68898
- MEYER, M.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments
p 191 A95-66777
- MILIC-EMILI, J.**
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
- MILLER, J. P.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- MILLS, T.**
Comets and life in the Universe p 190 A95-66761
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
- MINIHAN, DANIEL**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426
- MITCHELL, CHARLES E.**
Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263
- MOGFORD, RICHARD H.**
Using knowledge exploration tools to study airspace complexity in air traffic control
[HTN-95-90716] p 211 A95-69743
- MOGHADASI, HAMID**
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration
[HTN-95-A0124] p 201 A95-68907
- MONN, STEPHEN A.**
Interactive effects of emphysema and malnutrition on diaphragm structure and function
[HTN-95-A0148] p 206 A95-68931
- MOOIJ, J. M. V.**
Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration
[HTN-95-A0119] p 200 A95-68902
- MOORE, ROBERT J.**
Lower limit of body fat in healthy active men
[HTN-95-A0146] p 206 A95-68929
- MOREELS, G.**
Possible evidence of hydrocarbons released by the nucleus of Halley's comet
p 190 A95-66759
- MORIN, F. C., III**
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs
[HTN-95-A0144] p 205 A95-68927
- MURPHY, ELIZABETH D.**
Using knowledge exploration tools to study airspace complexity in air traffic control
[HTN-95-90716] p 211 A95-69743
- N**
- NAKAMURA, HIDETOSHI**
Regional lung hematocrit variation and assessment of acute lung injury
[HTN-95-A0111] p 198 A95-68894
- NAVARRO-GONZALEZ, R.**
Chemical studies on the possible existence of life on Mars
p 216 A95-66774
Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799
- NEALY, JOHN E.**
Radiation exposure predictions for long-duration-stay Mars missions
[BTN-95-EIX95032453043] p 209 A95-68268
- NICKLAS, B. J.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- NOUR, MOHAMED A.**
On the development of an expert system for wheelchair selection p 214 N95-19627
- NOWAK, PAUL S.**
Atmospheric pressure within lunar structure
[BTN-95-EIX95032452326] p 213 A95-68263
Feasibility study of inflatable structures for a lunar base
[BTN-95-EIX95041503792] p 213 A95-69223
- NUTH, J.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program
p 195 A95-66805
- O**
- OCAMPO-FRIEDMANN, R.**
A primitive cyanobacterium as pioneer microorganism for terraforming Mars p 193 A95-66784
- OLSON, ROBERT M.**
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness
[AD-A284849] p 210 N95-19714
- ORO, J.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
Comets and life in the Universe p 190 A95-66761
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
- OSEI, K.**
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-fat(cp) rats
[HTN-95-A0107] p 197 A95-68890
- OSHIMA, T.**
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
Studies in the search for life on Mars p 192 A95-66779
- OSIPOV, V. B.**
Modern aspects of planetary protection and requirements to sterilization of space hardware
p 216 A95-66785
- OSTROUMOV, V.**
A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782
- OVERHOLSER, K. A.**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
- P**
- PAILOUS, P.**
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
- PANDE, RAJIV**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems
[BTN-94-EIX94501433007] p 195 A95-67426
- PAPELIER, Y.**
Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise
[HTN-95-A0103] p 196 A95-68886
- PARKER, R. E.**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
- PATEL, KISHOR K.**
Pressure-flow specificity of inspiratory muscle training
[HTN-95-A0132] p 203 A95-68915
- PERMUTT, SOLBERT**
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs
[HTN-95-A0136] p 204 A95-68919
- PERREAU, M.**
Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765
- PESEC-RODRIGUEZ, R. A.**
Organic analysis of hydrogen cyanide polymers: Prebiotic and extraterrestrial chemistry
p 190 A95-66760
- PICOT, M.**
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
- PIZZARELLO, S.**
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite
p 190 A95-66762
- POEHLMAN, ERIC T.**
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women
[HTN-95-A0120] p 200 A95-68903
- PONNAMPERUMA, C.**
Chemical studies on the possible existence of life on Mars p 216 A95-66774
Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800
- POTAMAN, V. N.**
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
- POU, N. A.**
Pulmonary vascular resistance distribution and recruitment of microvascular surface area
[HTN-95-A0139] p 204 A95-68922
- PRANDI, E.**
Chest wall interrupter resistance in anesthetized paralyzed humans
[HTN-95-A0142] p 205 A95-68925
- PRATLEY, R. E.**
Effects of strength training on total and regional body composition in older men
[HTN-95-A0116] p 199 A95-68899
- PREFAUT, C.**
Performance and metabolic effects of benzodiazepine during submaximal exercise
[HTN-95-A0137] p 204 A95-68920
- PREMINGER, G. M.**
Characterization of fracture toughness of renal calculi using a microindentation technique
[BTN-94-EIX94361122293] p 209 A95-65873
- PUTZ, G.**
Surface activity of rabbit pulmonary surfactant subfractions at different concentrations in a captive bubble
[HTN-95-A0114] p 198 A95-68897
- Q**
- QIAN, Y.**
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
- R**
- RAMADURAI, S.**
Extinction of dinosaurs: A possible novel cause
p 191 A95-66769
- RAMANATHAN, M.**
Cardiopulmonary adaptations to pneumonectomy in dogs. 4: Membrane diffusing capacity and capillary blood volume
[HTN-95-A0153] p 208 A95-68936
- RAULIN, F.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992
[ISBN 0-08-042540-2] p 214 A95-66751
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
- REARDON, MATTHEW J.**
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM
[AD-A279193] p 211 N95-19811
- REGGIANI, CARLO**
Myosin isoforms in mammalian skeletal muscle
[HTN-95-A0102] p 195 A95-68885
- REMMERS, JOHN E.**
Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats
[HTN-95-A0127] p 201 A95-68910
- RICE, J.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
- RICHALET, JEAN-PAUL**
Energy balance at high altitude of 6,542 m
[HTN-95-A0140] p 205 A95-68923

RICHTER, ERIK A.
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888

RICKS, WENDELL R.
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745

RISI, S.
Biochemical constraints for survival under martian conditions p 192 A95-66778

ROBERTSON, BENGT
Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934

ROBERTSON, H. THOMAS
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928

RODENBURG, J. B.
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937

ROEDIGER, HENRY, III
Comparing performance on implicit memory tests [AD-A284861] p 212 N95-19740

ROGERS, WILLIAM H.
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745

ROTHSCHILD, L. J.
A "crylic" microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781

ROUSSELOT, P.
Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759

ROWELL, L. B.
Carotid baroreflex control of blood pressure and heart rate in men during dynamic exercise [HTN-95-A0103] p 196 A95-68886

RUBIN, M. A.
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899

RUEB, JUSTIN D.
Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744

RUMMEL, J. D.
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751

RYAN, A. S.
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899

S

SADEH, WILLY Z.
Feasibility study of inflatable structures for a lunar base [BTN-95-EIX95041503792] p 213 A95-69223

SADURSKI, RALPH
Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909

SAGAN, C.
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797

SAITO, T.
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766

SAKAMAKI, FUMIO
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

SALAMA, F.
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804

SANDFORD, S. A.
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802

SARTER, NADINE B.
Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742

SAYAMA, KOICHI
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

SAYKALLY, R. J.
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754

SCARPACE, PHILIP J.
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911

SCATTERGOOD, T. W.
Exobiology research on Space Station Freedom p 215 A95-66768

The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794

SCHIED, P.
Respiratory response to inhaled CO2 during positive inspiratory pressure in humans [HTN-95-A0141] p 205 A95-68924

SCHIAFFINO, STEFANO
Myosin isoforms in mammalian skeletal muscle [HTN-95-A0102] p 195 A95-68885

SCHIDLOWSKI, M.
Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775

Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-66807

SCHOENE, ROBERT B.
Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928

SCHUERCH, DAVID
Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934

SCHUERCH, SAMUEL
Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934

SCHUTTE, W. A.
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802

SCHWARTZ, D. E.
Search for life on Mars: Evaluation of techniques p 216 A95-66776

SCHWEHM, G. H.
Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

SCOON, G. E. N.
Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

SERIES, F.
Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure [HTN-95-A0138] p 204 A95-68921

SEVERINGHAUS, JOHN W.
Exercise O2 transport model assuming zero cytochrome PO2 at dot-VO2(sub max) [HTN-95-A0122] p 200 A95-68905

SHADE, ELIZABETH
Effects of hyperinflation and CPAP on work of breathing and respiratory failure in dogs [HTN-95-A0136] p 204 A95-68919

SHAW, LEONARD
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949

SHERMAN, W. M.
Effects of acute running exercise on whole body insulin action in obese male SHHF/Mcc-1a(cp) rats [HTN-95-A0107] p 197 A95-68890

SHU, YE
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911

SIECK, GARY C.
Interactive effects of emphysema and malnutrition on diaphragm structure and function [HTN-95-A0148] p 206 A95-68931

SILFER, J. A.
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763

SIMONSEN, LISA C.
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268

SNODGRASS, JOAN G.
Facilitation and interference in identification of pictures and words [AD-A285882] p 212 N95-19602

SOINA, V. S.
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

SOLYANINA, I. P.
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767

SORKIN, J.
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899

SPIERO, F.
Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787

SPINELLA, F.
Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801

SPRIET, LAWRENCE L.
High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932

SRIDHAR, G.
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933

SRINIVASAN, R. SRINI
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901

STABENAU, ERICH K.
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO2 equilibration [HTN-95-A0124] p 201 A95-68907

STEPHENS, J. R.
Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

STERNBERG, R.
The low temperature organic chemistry of Titan's geofluid p 218 A95-66796

STRATTON, D. M.
Exobiology research on Space Station Freedom p 215 A95-66768

STRAZZULLA, G.
Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801

STRIDDE, C.
Biochemical constraints for survival under martian conditions p 192 A95-66778

STRIEPE, SCOTT A.
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268

STURE, S.
Design and construction of shielded lunar outpost [BTN-95-EIX95032452328] p 213 A95-68265

SUDANO, JOHN J.
Minimizing human-machine interface failures in high risk systems [BTN-95-EIX95042474623] p 213 A95-68277

SUKI, BELA
Serial distribution of airway mechanical properties in dogs: Effects of histamine [HTN-95-A0110] p 197 A95-68893

SWEETSER, T. H.
Designing planetary protection into the Mars observer mission p 217 A95-66786

SWOAP, STEVEN J.
Interaction of the thyroid hormone and functional overload on skeletal muscle isomyosin expression [HTN-95-A0117] p 199 A95-68900

T

TAKEOKA, MICHIKO
Higher exercise performance and lower dot-VO2(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906

TANAKA, K.
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754

TASAKA, SADATOMO
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

TAVOLA, M.
Chest wall interrupter resistance in anesthetized paralyzed humans [HTN-95-A0142] p 205 A95-68925

TERASHIMA, TAKESHI
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

Y

- THOMPSON, W. R.**
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797
- THULLIER, PHILIPPE H.**
Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901
- TIKINSKY-RUPP, M. H.**
Hyperbaric oxygenation increases arousal and breathing movements in fetal lambs [HTN-95-A0144] p 205 A95-68927
- TOTH, MICHAEL J.**
Contribution of body composition and physical activity to age-related decline in peak dot-VO₂ in men and women [HTN-95-A0120] p 200 A95-68903
- TREUTH, M. S.**
Effects of strength training on total and regional body composition in older men [HTN-95-A0116] p 199 A95-68899
- TRIPATHY, SUKANT K.**
Intelligent materials properties of DNA and strategies for its incorporation into electroactive polymeric thin film systems [BTN-94-EIX94501433007] p 195 A95-67426
- TRIPLETT, N. T.**
Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930
- TROFIMOV, V. I.**
Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785
The experimental study of microbial contamination of the space hardware p 217 A95-66788
- TRUSCOTT, T. GEORGE**
Photobiology in medicine [AD-A284843] p 210 N95-19695
- TSUCHIYA, M.**
Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
- TSUKADA, HIROSHI**
Diameters of juxtacapillary venules determined by oil-drop method in rat lung [HTN-95-A0126] p 201 A95-68909
- TUMER, NIHAL**
Influence of exercise training on myocardial beta-adrenergic signal transduction: Differential regulation with age [HTN-95-A0128] p 202 A95-68911
- TURCOTTE, LORRAINE P.**
Impaired plasma FFA oxidation imposed by extreme CHO deficiency in contracting rat skeletal muscle [HTN-95-A0105] p 196 A95-68888
- TURKINGTON, T. G.**
Quantitative SPECT brain imaging: Effects of attenuation and detector response [BTN-94-EIX94361133479] p 209 A95-65976
- TZELEPIS, GEORGE E.**
Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915
- U**
- ULTMAN, JAMES S.**
Longitudinal distribution of ozone absorption in the lung: Effects of respiratory flow [HTN-95-A0112] p 198 A95-68895
- URANO, TESTUYA**
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894
- V**
- VAN BAAK, M. A.**
Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902
- VAN BORSELEN, F.**
Catecholamine responses to short-term high-intensity resistance exercise overtraining [HTN-95-A0147] p 206 A95-68930
- VAN DER HOEVEN, J. H.**
Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle [HTN-95-A0133] p 203 A95-68916

- VAN ECHELD, C. J. A.**
(31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937
- VAN ORDEN, A.**
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
- VANOYE, CARLOS G.**
Roles of intra- and extracellular carbonic anhydrase in alveolar-capillary CO₂ equilibration [HTN-95-A0124] p 201 A95-68907
- VASIN, V. B.**
The experimental study of microbial contamination of the space hardware p 217 A95-66788
- VEGA, DIEGO L.**
Lung volume specificity of inspiratory muscle training [HTN-95-A0131] p 202 A95-68914
Pressure-flow specificity of inspiratory muscle training [HTN-95-A0132] p 203 A95-68915
- VICAUT, ERIC**
Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle [HTN-95-A0106] p 196 A95-68889
- VIDLICH, MICHAEL A.**
Use of workload redlines: A KC-135 crew-reduction application [HTN-95-90717] p 211 A95-69744
- VILLORESI, G.**
The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775
- VOGEL, JAMES A.**
Lower limit of body fat in healthy active men [HTN-95-A0146] p 206 A95-68929
- VOROBIOVA, E. A.**
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

W

- WAKI, YASUHIRO**
Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894
- WALLIS, MAX**
Extinction of dinosaurs: A possible novel cause p 191 A95-66769
- WANG, LI-HAU**
Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906
- WANG, YAO**
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949
- WECKWERTH, G.**
Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775
- WESTERTERP, KLAAS R.**
Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923
- WHARTON, R. A.**
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
- WHITE, M. R.**
Search for life on Mars: Evaluation of techniques p 216 A95-66776
- WICKRAMASINGHE, N. C.**
Extinction of dinosaurs: A possible novel cause p 191 A95-66769
- WICKRAMASINGHE, N. S. M. D.**
Direct interaction between amino acids and nucleotides as a possible physicochemical basis for the origin of the genetic code p 194 A95-66800
- WOODS, DAVID D.**
Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742
- WOOLLES, W. R.**
Gender-related differences in the baroreceptor reflex control of heart rate in normotensive humans [HTN-95-A0115] p 199 A95-68898
- WOUTERS, LOEK**
Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923
- WRIGHT, F.**
Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920

Z


- ZHAN, WEN-ZHI**
Interactive effects of emphysema and malnutrition on diaphragm structure and function [HTN-95-A0148] p 206 A95-68931
- ZHANG, H. Y.**
Fluid flow and heat transfer in the crescent-shaped lumen catheter [BTN-94-EIX94361122408] p 209 A95-65904
- ZHAO, B.**
Detection of free radicals by electron spin resonance in rat diaphragm after resistive loading [HTN-95-A0135] p 203 A95-68918
- ZHONG, P.**
Characterization of fracture toughness of renal calculi using a microindentation technique [BTN-94-EIX94361122293] p 209 A95-65873
- ZHU, QIN-FAN**
Maximally smooth image recovery in transform coding [BTN-94-EIX94361135047] p 209 A95-65949
- ZIJDEWIND, I.**
Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist [HTN-95-A0152] p 207 A95-68935
- ZINTEL, T. A.**
Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933
- ZVYAGINTSEV, D. G.**
Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783

CORPORATE SOURCE INDEX

AEROSPACE MEDICINE AND BIOLOGY / A Continuing Bibliography (Supplement 401)

May 1995

Typical Corporate Source Index Listing



Listings in this index are arranged alphabetically by corporate source. The title of the document is used to provide a brief description of the subject matter. The page number and the accession number are included in each entry to assist the user in locating the abstract in the abstract section. If applicable, a report number is also included as an aid in identifying the document.

A

- Alabama Univ., Birmingham, AL.**
Cumulative effects of repeated brief cerebral ischemia [AD-A284234] p 210 N95-19698
- Army Aeromedical Research Lab., Fort Rucker, AL.**
Collection of real-time, multichannel EEG data from helicopter pilots in flight: A feasibility study [AD-A285050] p 212 N95-19564
- Army Research Inst. of Environmental Medicine, Natick, MA.**
Distributed medical database system for real-time monitoring of the health and risk exposure of military research study volunteers at USARIEM [AD-A279193] p 211 N95-19811
Annual historical report, calendar year 1993 [AD-A280178] p 211 N95-19827

D

- Department of Energy, Washington, DC.**
Activities of the Division of Energy Biosciences [DE95-001046] p 208 N95-19613

I

- Institute for Human Factors TNO, Soesterberg (Netherlands).**
Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538

J

- Jet Propulsion Lab., California Inst. of Tech., Pasadena, CA.**
Designing planetary protection into the Mars observer mission p 217 A95-66786

K

- Keele Univ. (England).**
Photobiology in medicine [AD-A284843] p 210 N95-19695
- Kent State Univ., OH.**
On the development of an expert system for wheelchair selection p 214 N95-19627

N

- National Aeronautics and Space Administration, Washington, DC.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
Toward the detection of pure carbon clusters in the Interstellar Medium (ISM) p 189 A95-66754
Organic molecules in the gas phase of dense interstellar clouds p 189 A95-66755
Characteristics and formation of amino acids and hydroxy acids of the Murchison meteorite p 190 A95-66762
Stable isotope analysis at the molecular level: A new approach for determining the origins of amino acids in the Murchison meteorite p 190 A95-66763
Site selection for Mars exobiology p 215 A95-66771

- Designing planetary protection into the Mars observer mission p 217 A95-66786
Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793
Analytical pyrolysis experiments of Titan aerosol analogues in preparation for the Cassini Huygens mission p 193 A95-66797
On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
Interleukin-2 therapy reverses some immunosuppressive effects of skeletal unloading [HTN-95-A0113] p 198 A95-68896
Aerospace medicine and biology: A continuing bibliography with indexes (supplement 397) [NASA-SP-7011(397)] p 210 N95-19746

- National Aeronautics and Space Administration, Ames Research Center, Moffett Field, CA.**
Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751
Exobiology research on Space Station Freedom p 215 A95-66768
Exobiological exploration of Mars p 215 A95-66770
Site selection for Mars exobiology p 215 A95-66771
Life on Mars? 1: The chemical environment p 215 A95-66772
Life on Mars? 2: Physical restrictions p 215 A95-66773
Search for life on Mars: Evaluation of techniques p 216 A95-66776
A preliminary comparison of two perennially ice-covered lakes in Antarctica: Analogs of past martian lacustrine environments p 191 A95-66777
A "cryptic" microbial mat: A new model ecosystem for extant life on Mars p 192 A95-66781
Surface-atmosphere interactions on Titan compared with those on the pre-biotic Earth p 193 A95-66793
The physical nature of Titan's aerosols: Laboratory simulations p 193 A95-66794
Very low temperature formaldehyde reactions and the build-up of organic molecules in comets and interstellar ices p 194 A95-66802
Polycyclic aromatic hydrocarbon ions and the diffuse interstellar bands p 194 A95-66804

- Pilot interaction with cockpit automation 2: An experimental study of pilots' model and awareness of the Flight Management System [HTN-95-90715] p 211 A95-69742
- National Aeronautics and Space Administration, Goddard Space Flight Center, Greenbelt, MD.**
Cosmic dust analog simulation in a microgravity environment: The STARDUST program p 195 A95-66805

- National Aeronautics and Space Administration, Langley Research Center, Hampton, VA.**
Radiation exposure predictions for long-duration-stay Mars missions [BTN-95-EIX95032453043] p 209 A95-68268
Cognitive representations of flight-deck information attributes [HTN-95-90718] p 212 A95-69745
- Naval Research Lab., Washington, DC.**
Proceedings of Natural Sciences Symposium [AD-A285452] p 208 N95-19536
- New York Univ., New York, NY.**
Facilitation and interference in identification of pictures and words [AD-A285882] p 212 N95-19602

P

- Physics and Electronics Lab. TNO, The Hague (Netherlands).**
High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537
- Prins Maurits Lab. TNO, Rijswijk (Netherlands).**
Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework [AD-A285133] p 212 N95-19583

R

- Rice Univ., Houston, TX.**
Comparing performance on implicit memory tests [AD-A284861] p 212 N95-19740
- Rothe Development, Inc., San Antonio, TX.**
Echo imaging techniques determine the size of intravascular bubbles in decompression sickness [AD-A284849] p 210 N95-19714


SOURCE

FOREIGN TECHNOLOGY INDEX

AEROSPACE MEDICINE AND BIOLOGY / A Continuing Bibliography (Supplement 401)

May 1995

Typical Foreign Technology Index Listing



Listings in this index are arranged alphabetically by country of intellectual origin. The title of the document is used to provide a brief description of the subject matter. The page number and accession number are included in each entry to assist the user in locating the abstract in the abstract section. If applicable, a report number is also included as an aid in identifying the document.

C

CANADA

- Laboratory astrophysics and molecular astronomy of pure carbon molecules p 189 A95-66753
- Hypoxia similarly impairs metabolic responses to cutaneous and core cold stimuli in conscious rats [HTN-95-A0127] p 201 A95-68910
- Performance and metabolic effects of benzodiazepine during submaximal exercise [HTN-95-A0137] p 204 A95-68920
- Influence of lung volume dependence of upper airway resistance during continuous negative airway pressure [HTN-95-A0138] p 204 A95-68921
- Pulmonary gas exchange during exercise in athletes. 1: Ventilation-perfusion mismatch and diffusion limitation [HTN-95-A0145] p 206 A95-68928
- High physiological levels of epinephrine do not enhance muscle glycogenolysis during tetanic stimulation [HTN-95-A0149] p 207 A95-68932
- Lung volumes and expiratory flow limitation during exercise in interstitial lung disease [HTN-95-A0150] p 207 A95-68933
- Surface activity of lipid extract surfactant in relation to film area compression and collapse [HTN-95-A0151] p 207 A95-68934
- CHINA**
- Higher exercise performance and lower dot-VO₂(sub max) in Tibetan than Han residents at 4,700 m altitude [HTN-95-A0123] p 201 A95-68906

F

FRANCE

- Possible evidence of hydrocarbons released by the nucleus of Halley's comet p 190 A95-66759
- Were micrometeorites a source of prebiotic molecules on the early Earth? p 191 A95-66765

- Titan's atmosphere composition: Certainties and speculations p 217 A95-66792
- The low temperature organic chemistry of Titan's geofluid p 218 A95-66796
- Isotopic ratios in planetary atmospheres p 218 A95-66806
- Carotoid baroreflex control of blood pressure and heart rate in men during dynamic exercise [HTN-95-A0103] p 196 A95-68886
- Nitric oxide-independent response to acetylcholine by terminal arterioles in rat cremaster muscle [HTN-95-A0106] p 196 A95-68889
- Simulation of cardiovascular response to lower body negative pressure from 0 to -40 mmHg [HTN-95-A0118] p 199 A95-68901
- Respiratory response to inhaled CO₂ during positive inspiratory pressure in humans [HTN-95-A0141] p 205 A95-68924

G

GERMANY

- Mass-spectrometric in situ studies of cometary organics for p/Halley and options for the future p 190 A95-66758
- Phosphorus as a potential guide in the search for extinct life on Mars p 216 A95-66775
- Biochemical constraints for survival under martian conditions p 192 A95-66778
- Planetary protection considerations for MarsNet and Mars sample return missions p 217 A95-66789
- Carbonaceous matter in cometary dust and coma p 218 A95-66803
- Isotope fractionations in the terrestrial carbon cycle: A brief overview p 195 A95-66807

I

INDIA

- Extinction of dinosaurs: A possible novel cause p 191 A95-66769

ITALY

- Production and evolution of carbonaceous material by ion irradiation in space p 218 A95-66801
- Myosin isoforms in mammalian skeletal muscle [HTN-95-A0102] p 195 A95-68885
- Chest wall interrupter resistance in anesthetized paralyzed humans [HTN-95-A0142] p 205 A95-68925

J

JAPAN

- Formation of bioorganic compounds in planetary atmospheres by cosmic radiation p 214 A95-66766
- Studies in the search for life on Mars p 192 A95-66779
- Regional lung hematocrit variation and assessment of acute lung injury [HTN-95-A0111] p 198 A95-68894

M

MEXICO

- On the levels of enzymatic substrate specificity: Implications for the early evolution of metabolic pathways p 193 A95-66798
- Role of trace metal ions in chemical evolution. The case of free-radical reactions p 194 A95-66799

N

NETHERLANDS

- Life sciences and space research 24 (4): Planetary biology and origins of life; Topical Meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F3) of the COSPAR Plenary Meeting, 29th, Washington, DC, Aug. 28-Sep. 5, 1992 [ISBN 0-08-042540-2] p 214 A95-66751

- Planetary environment protection id no: F3.3 - M.1.05 implications for the development of a network of surface stations on Mars p 217 A95-66787
- Effect of glucose infusion on endurance performance after beta-adrenoceptor blocker administration [HTN-95-A0119] p 200 A95-68902
- Supernormal muscle fiber conduction velocity during intermittent isometric exercise in human muscle [HTN-95-A0133] p 203 A95-68916
- Energy balance at high altitude of 6,542 m [HTN-95-A0140] p 205 A95-68923
- Index finger position and force of the human first dorsal interosseus and its ulnar nerve antagonist [HTN-95-A0152] p 207 A95-68935
- (31)P-MRS and simultaneous quantification of dynamic human quadriceps exercise in a whole body MR scanner [HTN-95-A0154] p 208 A95-68937
- High performance computing for medical image interpretation [AD-A285457] p 210 N95-19537
- Learning effects on strategy selection in a dynamic task environment as a function of time pressure [AD-A285458] p 212 N95-19538
- Multiple-task performance: A critical review of the literature and a cognitive neuroscience framework [AD-A285133] p 212 N95-19583

R

RUSSIA

- Radiation and thermal stabilities of adenine nucleotides p 191 A95-66767
- Biogeochemical evidence of microbial activity on Mars p 192 A95-66780
- A physical and chemical characterization of Martian permafrost as a possible habitat for viable microorganisms p 216 A95-66782
- Preservation of cell structures in permafrost: A model for exobiology p 192 A95-66783
- Modern aspects of planetary protection and requirements to sterilization of space hardware p 216 A95-66785
- The experimental study of microbial contamination of the space hardware p 217 A95-66788
- The use of cosmic rays for continuous monitoring and prediction of some dangerous phenomena for the earth's civilization [HTN-95-60524] p 195 A95-68775

S

SWEDEN

- Tissue oxygen and carbon dioxide stores and breath-hold diving in humans [HTN-95-A0108] p 197 A95-68891


U

UNITED KINGDOM

- Raindrops on Titan p 218 A95-66795
- Effect of hyperoxia at 1 and 2 ATA on hypoxia and hypercapnia in human skin during experimental inflammation [HTN-95-A0130] p 202 A95-68913
- Photobiology in medicine [AD-A284843] p 210 N95-19695

REPORT NUMBER INDEX

Typical Report Number Index Listing



Listings in this index are arranged alphanumerically by report number. The page number indicates the page on which the citation is located. The accession number denotes the number by which the citation is identified. An asterisk (*) indicates that the item is a NASA report. A pound sign (#) indicates that the item is available on microfiche.

NASA-CR-191326	p 81	N94-13132	*	#
AD-A279193	p 211	N95-19811		
AD-A280178	p 211	N95-19827		
AD-A284234	p 210	N95-19698		#
AD-A284843	p 210	N95-19695		#
AD-A284849	p 210	N95-19714		#
AD-A284861	p 212	N95-19740		
AD-A285050	p 212	N95-19564		#
AD-A285133	p 212	N95-19583		#
AD-A285452	p 208	N95-19536		#
AD-A285457	p 210	N95-19537		#
AD-A285458	p 212	N95-19538		#
AD-A285882	p 212	N95-19602		#
AFOSR-94-0481TR	p 210	N95-19698		#
AFOSR-94-0577TR	p 212	N95-19740		
AFOSR-94-0670TR	p 212	N95-19602		#
AL/CF-TR-1994-0033	p 210	N95-19714		#
BTN-94-EIX94361122293	p 209	A95-65873		
BTN-94-EIX94361122408	p 209	A95-65904		
BTN-94-EIX94361133479	p 209	A95-65976		
BTN-94-EIX94361135047	p 209	A95-65949		
BTN-94-EIX94501433007	p 195	A95-67426		
BTN-95-EIX95032452326	p 213	A95-68263		
BTN-95-EIX95032452328	p 213	A95-68265		
BTN-95-EIX95032453043	p 209	A95-68268		*
BTN-95-EIX95041503792	p 213	A95-69223		
BTN-95-EIX95042474623	p 213	A95-68277		
BTN-95-EIX95042477107	p 213	A95-68352		
CSP-93-1041	p 210	N95-19695		#
DE95-001046	p 208	N95-19613		#
DOE/ER-0621P	p 208	N95-19613		#
FEL-93-B160	p 210	N95-19537		#
HTN-95-A0102	p 195	A95-68885		
HTN-95-A0103	p 196	A95-68886		
HTN-95-A0104	p 196	A95-68887		
HTN-95-A0105	p 196	A95-68888		
HTN-95-A0106	p 196	A95-68889		
HTN-95-A0107	p 197	A95-68890		
HTN-95-A0108	p 197	A95-68891		
HTN-95-A0109	p 197	A95-68892		
HTN-95-A0110	p 197	A95-68893		
HTN-95-A0111	p 198	A95-68894		
HTN-95-A0112	p 198	A95-68895		
HTN-95-A0113	p 198	A95-68896		*
HTN-95-A0114	p 198	A95-68897		
HTN-95-A0115	p 199	A95-68898		
HTN-95-A0116	p 199	A95-68899		
HTN-95-A0117	p 199	A95-68900		
HTN-95-A0118	p 199	A95-68901		
HTN-95-A0119	p 200	A95-68902		
HTN-95-A0120	p 200	A95-68903		
HTN-95-A0121	p 200	A95-68904		
HTN-95-A0122	p 200	A95-68905		
HTN-95-A0123	p 201	A95-68906		
HTN-95-A0124	p 201	A95-68907		
HTN-95-A0125	p 201	A95-68908		
HTN-95-A0126	p 201	A95-68909		
HTN-95-A0127	p 201	A95-68910		
HTN-95-A0128	p 202	A95-68911		
HTN-95-A0129	p 202	A95-68912		
HTN-95-A0130	p 202	A95-68913		
HTN-95-A0131	p 202	A95-68914		
HTN-95-A0132	p 203	A95-68915		
HTN-95-A0133	p 203	A95-68916		
HTN-95-A0134	p 203	A95-68917		
HTN-95-A0135	p 203	A95-68918		
HTN-95-A0136	p 204	A95-68919		
HTN-95-A0137	p 204	A95-68920		
HTN-95-A0138	p 204	A95-68921		
HTN-95-A0139	p 204	A95-68922		
HTN-95-A0140	p 205	A95-68923		
HTN-95-A0141	p 205	A95-68924		
HTN-95-A0142	p 205	A95-68925		
HTN-95-A0143	p 205	A95-68926		
HTN-95-A0144	p 205	A95-68927		
HTN-95-A0145	p 206	A95-68928		
HTN-95-A0146	p 206	A95-68929		
HTN-95-A0147	p 206	A95-68930		
HTN-95-A0148	p 206	A95-68931		
HTN-95-A0149	p 207	A95-68932		
HTN-95-A0150	p 207	A95-68933		
HTN-95-A0151	p 207	A95-68934		
HTN-95-A0152	p 207	A95-68935		
HTN-95-A0153	p 208	A95-68936		
HTN-95-A0154	p 208	A95-68937		
HTN-95-A0155	p 208	A95-68938		
HTN-95-10500	p 214	A95-66751	*	
HTN-95-60524	p 195	A95-68775	*	
HTN-95-90715	p 211	A95-69742	*	
HTN-95-90716	p 211	A95-69743	*	
HTN-95-90717	p 211	A95-69744	*	
HTN-95-90718	p 212	A95-69745	*	
HTN-95-90719	p 209	A95-69746	*	
ISBN 0-08-042540-2	p 214	A95-66751	*	
IZF-1994-B-5	p 212	N95-19583		#
NAS 1.21:7011(397)	p 210	N95-19746	*	
NASA-SP-7011(397)	p 210	N95-19746	*	
NRL/PU/6000-94-252	p 208	N95-19536		#
TDCK-93-2783	p 210	N95-19537		#
TDCK-94-0043	p 212	N95-19583		#
TDCK-94-0055	p 212	N95-19538		#
TNO-TM-1994-B-12	p 212	N95-19538		#
USAARL-94-26	p 212	N95-19564		#
USARIEM-T94-13	p 211	N95-19811		


REPORT

ACCESSION NUMBER INDEX

AEROSPACE MEDICINE AND BIOLOGY / A Continuing Bibliography (Supplement 401)

May 1995

Typical Accession Number Index Listing



Listings in this index are arranged alphanumerically by accession number. The page number indicates the page on which the citation is located. The accession number denotes the number by which the citation is identified. An asterisk (*) indicates that the item is a NASA report. A pound sign (#) indicates that the item is available on microfiche.

A95-65873	p 209	A95-66805 *	p 195
A95-65904	p 209	A95-66806	p 218
A95-65949	p 209	A95-66807	p 195
A95-65976	p 209	A95-67426	p 195
A95-66751 *	p 214	A95-68263	p 213
A95-66752	p 214	A95-68265	p 213
A95-66753	p 189	A95-68268 *	p 209
A95-66754 *	p 189	A95-68277	p 213
A95-66755 *	p 189	A95-68352	p 213
A95-66757	p 189	A95-68775	p 195
A95-66758	p 190	A95-68885	p 195
A95-66759	p 190	A95-68886	p 196
A95-66760	p 190	A95-68887	p 196
A95-66761 *	p 190	A95-68888	p 196
A95-66762 *	p 190	A95-68889	p 196
A95-66763 *	p 190	A95-68890	p 197
A95-66764	p 214	A95-68891	p 197
A95-66765	p 191	A95-68892	p 197
A95-66766	p 214	A95-68893	p 197
A95-66767	p 191	A95-68894	p 198
A95-66768 *	p 215	A95-68895	p 198
A95-66769	p 191	A95-68896 *	p 198
A95-66770 *	p 215	A95-68897	p 198
A95-66771 *	p 215	A95-68898	p 199
A95-66772 *	p 215	A95-68899	p 199
A95-66773 *	p 215	A95-68900	p 199
A95-66774	p 216	A95-68901	p 199
A95-66775	p 216	A95-68902	p 200
A95-66776 *	p 216	A95-68903	p 200
A95-66777 *	p 191	A95-68904	p 200
A95-66778	p 192	A95-68905	p 200
A95-66779	p 192	A95-68906	p 201
A95-66780	p 192	A95-68907	p 201
A95-66781 *	p 192	A95-68908	p 201
A95-66782	p 216	A95-68909	p 201
A95-66783	p 192	A95-68910	p 201
A95-66784	p 193	A95-68911	p 202
A95-66785	p 216	A95-68912	p 202
A95-66786 *	p 217	A95-68913	p 202
A95-66787	p 217	A95-68914	p 202
A95-66788	p 217	A95-68915	p 203
A95-66789	p 217	A95-68916	p 203
A95-66792	p 217	A95-68917	p 203
A95-66793 *	p 193	A95-68918	p 203
A95-66794 *	p 193	A95-68919	p 204
A95-66795	p 218	A95-68920	p 204
A95-66796	p 218	A95-68921	p 204
A95-66797 *	p 193	A95-68922	p 204
A95-66798 *	p 193	A95-68923	p 205
A95-66799	p 194	A95-68924	p 205
A95-66800	p 194	A95-68925	p 205
A95-66801	p 218	A95-68926	p 205
A95-66802 *	p 194	A95-68927	p 205
A95-66803	p 218	A95-68928	p 206
A95-66804 *	p 194	A95-68929	p 206
		A95-68930	p 206
		A95-68931	p 206
		A95-68932	p 207
		A95-68933	p 207
		A95-68934	p 207
		A95-68935	p 207
		A95-68936	p 208
		A95-68937	p 208
		A95-68938	p 208
		A95-69223	p 213
		A95-69742 *	p 211
		A95-69743	p 211
		A95-69744	p 211
		A95-69745 *	p 212
		A95-69746	p 209
		N95-19536 #	p 208
		N95-19537 #	p 210
		N95-19538 #	p 212
		N95-19564 #	p 212
		N95-19583 #	p 212
		N95-19602 #	p 212
		N95-19613 #	p 208
		N95-19627 * #	p 214
		N95-19695 #	p 210
		N95-19698 #	p 210
		N95-19714 #	p 210
		N95-19740	p 212
		N95-19746 *	p 210
		N95-19811	p 211
		N95-19827	p 211

ACCESSION

AVAILABILITY OF CITED PUBLICATIONS

OPEN LITERATURE ENTRIES (A95-60000 Series)

Inquiries and requests should be addressed to NASA Center for AeroSpace Information, 800 Elkridge Landing Road, Linthicum Heights, MD 21090-2934. Orders are also taken by telephone, (301) 621-0390, e-mail, help@sti.nasa.gov, and fax, (301) 621-0134. Please refer to the accession number when request-ing publications.

STAR ENTRIES (N95-10000 Series)

One or more sources from which a document announced in *STAR* is available to the public is ordinarily given on the last line of the citation. The most commonly indicated sources and their acronyms or abbreviations are listed below, and their addresses are listed on page APP-3. If the publication is available from a source other than those listed, the publisher and his address will be displayed on the availability line or in combination with the corporate source line.

Avail: NASA CASI. Sold by the NASA Center for AeroSpace Information. Prices for hard copy (HC) and microfiche (MF) are indicated by a price code following the letters HC or MF in the *STAR* citation. Current values for the price codes are given in the tables on page APP-5.

NOTE ON ORDERING DOCUMENTS: When ordering publications from NASA CASI, use the N accession number or other report number. It is also advisable to cite the title and other bibliographic identification.

Avail: SOD (or GPO). Sold by the Superintendent of Documents, U.S. Government Printing Office, in hard copy.

Avail: BLL (formerly NLL): British Library Lending Division, Boston Spa, Wetherby, Yorkshire, England. Photocopies available from this organization at the price shown. (If none is given, inquiry should be addressed to the BLL.)

Avail: DOE Depository Libraries. Organizations in U.S. cities and abroad that maintain collections of Department of Energy reports, usually in microfiche form, are listed in *Energy Research Abstracts*. Services available from the DOE and its depositories are described in a booklet, *DOE Technical Information Center - Its Functions and Services* (TID-4660), which may be obtained without charge from the DOE Technical Information Center.

Avail: ESDU. Pricing information on specific data, computer programs, and details on Engineering Sciences Data Unit (ESDU) topic categories can be obtained from ESDU International Ltd. Requesters in North America should use the Virginia address while all other requesters should use the London address, both of which are on page APP-3.

Avail: Fachinformationszentrum Karlsruhe. Gesellschaft für wissenschaftlich-technische Information mbH 76344 Eggenstein-Leopoldshafen, Germany.

Avail: HMSO. Publications of Her Majesty's Stationery Office are sold in the U.S. by Pendragon House, Inc. (PHI), Redwood City, CA. The U.S. price (including a service and mailing charge) is given, or a conversion table may be obtained from PHI.

Avail: Issuing Activity, or Corporate Author, or no indication of availability. Inquiries as to the availability of these documents should be addressed to the organization shown in the citation as the corporate author of the document.

Avail: NASA Public Document Rooms. Documents so indicated may be examined at or purchased from the National Aeronautics and Space Administration (JBD-4), Public Documents Room (Room 1H23), Washington, DC 20546-0001, or public document rooms located at NASA installations, and the NASA Pasadena Office at the Jet Propulsion Laboratory.

Avail: NTIS. Sold by the National Technical Information Service. Initially distributed microfiche under the NTIS SRIM (Selected Research in Microfiche) are available. For information concerning this service, consult the NTIS Subscription Section, Springfield, VA 22161.

Avail: Univ. Microfilms. Documents so indicated are dissertations selected from *Dissertation Abstracts* and are sold by University Microfilms as xerographic copy (HC) and microfilm. All requests should cite the author and the Order Number as they appear in the citation.

Avail: US Patent and Trademark Office. Sold by Commissioner of Patents and Trademarks, U.S. Patent and Trademark Office, at the standard price of \$1.50 each, postage free.

Avail: (US Sales Only). These foreign documents are available to users within the United States from the National Technical Information Service (NTIS). They are available to users outside the United States through the International Nuclear Information Service (INIS) representative in their country, or by applying directly to the issuing organization.

Avail: USGS. Originals of many reports from the U.S. Geological Survey, which may contain color illustrations, or otherwise may not have the quality of illustrations preserved in the microfiche or facsimile reproduction, may be examined by the public at the libraries of the USGS field offices whose addresses are listed on page APP-3. The libraries may be queried concerning the availability of specific documents and the possible utilization of local copying services, such as color reproduction.

FEDERAL DEPOSITORY LIBRARY PROGRAM

In order to provide the general public with greater access to U.S. Government publications, Congress established the Federal Depository Library Program under the Government Printing Office (GPO), with 53 regional depositories responsible for permanent retention of material, inter-library loan, and reference services. At least one copy of nearly every NASA and NASA-sponsored publication, either in printed or microfiche format, is received and retained by the 53 regional depositories. A list of the regional GPO libraries, arranged alphabetically by state, appears on the inside back cover of this issue. These libraries are *not* sales outlets. A local library can contact a regional depository to help locate specific reports, or direct contact may be made by an individual.

PUBLIC COLLECTION OF NASA DOCUMENTS

An extensive collection of NASA and NASA-sponsored publications is maintained by the British Library Lending Division, Boston Spa, Wetherby, Yorkshire, England for public access. The British Library Lending Division also has available many of the non-NASA publications cited in *STAR*. European requesters may purchase facsimile copy or microfiche of NASA and NASA-sponsored documents, those identified by both the symbols # and * from ESA — Information Retrieval Service European Space Agency, 8-10 rue Mario-Nikis, 75738 CEDEX 15, France.

STANDING ORDER SUBSCRIPTIONS

NASA SP-7011 supplements and annual index are available from the NASA Center for AeroSpace Information (CASI) on standing order subscription. Standing order subscriptions do not terminate at the end of a year, as do regular subscriptions, but continue indefinitely unless specifically terminated by the subscriber.

ADDRESSES OF ORGANIZATIONS

British Library Lending Division
Boston Spa, Wetherby, Yorkshire
England

National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161

Commissioner of Patents and Trademarks
U.S. Patent and Trademark Office
Washington, DC 20231

Pendragon House, Inc.
899 Broadway Avenue
Redwood City, CA 94063

Department of Energy
Technical Information Center
P.O. Box 62
Oak Ridge, TN 37830

Superintendent of Documents
U.S. Government Printing Office
Washington, DC 20402

European Space Agency-
Information Retrieval Service ESRIN
Via Galileo Galilei
00044 Frascati (Rome) Italy

University Microfilms
A Xerox Company
300 North Zeeb Road
Ann Arbor, MI 48106

Engineering Sciences Data Unit International
P.O. Box 1633
Manassas, VA 22110

University Microfilms, Ltd.
Tylers Green
London, England

Engineering Sciences Data Unit
International, Ltd.
251-259 Regent Street
London, W1R 7AD, England

U.S. Geological Survey Library National Center
MS 950
12201 Sunrise Valley Drive
Reston, VA 22092

Fachinformationszentrum Karlsruhe
Gesellschaft für wissenschaftlich-technische
Information mbH
76344 Eggenstein-Leopoldshafen, Germany

U.S. Geological Survey Library
2255 North Gemini Drive
Flagstaff, AZ 86001

Her Majesty's Stationery Office
P.O. Box 569, S.E. 1
London, England

U.S. Geological Survey
345 Middlefield Road
Menlo Park, CA 94025

NASA Center for AeroSpace Information
800 Elkridge Landing Road
Linthicum Heights, MD 21090-2934

U.S. Geological Survey Library
Box 25046
Denver Federal Center, MS914
Denver, CO 80225

National Aeronautics and Space Administration
Scientific and Technical Information Office
(Code JT)
Washington, DC 20546-0001

NASA CASI PRICE CODE TABLE

(Effective January 1, 1995)

CASI PRICE CODE	NORTH AMERICAN PRICE	FOREIGN PRICE
A01	\$ 6.00	\$ 12.00
A02	9.00	18.00
A03	17.50	35.00
A04-A05	19.50	39.00
A06-A09	27.00	54.00
A10-A13	36.50	73.00
A14-A17	44.50	89.00
A18-A21	52.00	104.00
A22-A25	61.00	122.00
A99	Call For Price	Call For Price

IMPORTANT NOTICE

For users not registered at the NASA CASI, prepayment is required. Additionally, a shipping and handling fee of \$1.00 per document for delivery within the United States and \$9.00 per document for delivery outside the United States is charged.

For users registered at the NASA CASI, document orders may be invoiced at the end of the month, charged against a deposit account, or paid by check or credit card. NASA CASI accepts American Express, Diners' Club, MasterCard, and VISA credit cards. There are no shipping and handling charges. To register at the NASA CASI, please request a registration form through the NASA Access Help Desk at the address or numbers below.

NASA Center for AeroSpace Information
800 Elkridge Landing Road
Linthicum Heights, MD 21090-2934
Telephone: (301) 621-0390
E-mail: help@sti.nasa.gov
Fax: (301) 621-0134

REPORT DOCUMENT PAGE

1. Report No. NASA SP-7011 (401)	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Aerospace Medicine and Biology A Continuing Bibliography (Supplement 401)		5. Report Date May 1995	
		6. Performing Organization Code JT	
7. Author(s)		8. Performing Organization Report No.	
		10. Work Unit No.	
9. Performing Organization Name and Address NASA Scientific and Technical Information Office		11. Contract or Grant No.	
		13. Type of Report and Period Covered Special Publication	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, DC 20546-0001		14. Sponsoring Agency Code	
		15. Supplementary Notes	
16. Abstract This report lists 140 reports, articles and other documents recently announced in the NASA STI Database.			
17. Key Words (Suggested by Author(s)) Aerospace Medicine Bibliographies Biological Effects		18. Distribution Statement Unclassified - Unlimited Subject Category - 52	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 72	22. Price A04/HC

FEDERAL REGIONAL DEPOSITORY LIBRARIES

ALABAMA

AUBURN UNIV. AT MONTGOMERY LIBRARY
Documents Dept.
7300 University Dr.
Montgomery, AL 36117-3596
(205) 244-3650 Fax: (205) 244-0678

UNIV. OF ALABAMA

Amelia Gayle Gorgas Library
Govt. Documents
P.O. Box 870266
Tuscaloosa, AL 35487-0266
(205) 348-6046 Fax: (205) 348-0760

ARIZONA

DEPT. OF LIBRARY, ARCHIVES, AND PUBLIC RECORDS
Research Division
Third Floor, State Capitol
1700 West Washington
Phoenix, AZ 85007
(602) 542-3701 Fax: (602) 542-4400

ARKANSAS

ARKANSAS STATE LIBRARY
State Library Service Section
Documents Service Section
One Capitol Mall
Little Rock, AR 72201-1014
(501) 682-2053 Fax: (501) 682-1529

CALIFORNIA

CALIFORNIA STATE LIBRARY
Govt. Publications Section
P.O. Box 942837 - 914 Capitol Mall
Sacramento, CA 94337-0091
(916) 654-0069 Fax: (916) 654-0241

COLORADO

UNIV. OF COLORADO - BOULDER
Libraries - Govt. Publications
Campus Box 184
Boulder, CO 80309-0184
(303) 492-8834 Fax: (303) 492-1881

DENVER PUBLIC LIBRARY

Govt. Publications Dept. BSG
1357 Broadway
Denver, CO 80203-2165
(303) 640-8846 Fax: (303) 640-8817

CONNECTICUT

CONNECTICUT STATE LIBRARY
231 Capitol Avenue
Hartford, CT 06106
(203) 566-4971 Fax: (203) 566-3322

FLORIDA

UNIV. OF FLORIDA LIBRARIES
Documents Dept.
240 Library West
Gainesville, FL 32611-2048
(904) 392-0366 Fax: (904) 392-7251

GEORGIA

UNIV. OF GEORGIA LIBRARIES
Govt. Documents Dept.
Jackson Street
Athens, GA 30602-1645
(706) 542-8949 Fax: (706) 542-4144

HAWAII

UNIV. OF HAWAII
Hamilton Library
Govt. Documents Collection
2550 The Mall
Honolulu, HI 96822
(808) 948-8230 Fax: (808) 956-5968

IDAHO

UNIV. OF IDAHO LIBRARY
Documents Section
Rayburn Street
Moscow, ID 83844-2353
(208) 885-6344 Fax: (208) 885-6817

ILLINOIS

ILLINOIS STATE LIBRARY
Federal Documents Dept.
300 South Second Street
Springfield, IL 62701-1796
(217) 782-7596 Fax: (217) 782-6437

INDIANA

INDIANA STATE LIBRARY
Serials/Documents Section
140 North Senate Avenue
Indianapolis, IN 46204-2296
(317) 232-3679 Fax: (317) 232-3728

IOWA

UNIV. OF IOWA LIBRARIES
Govt. Publications
Washington & Madison Streets
Iowa City, IA 52242-1166
(319) 335-5926 Fax: (319) 335-5900

KANSAS

UNIV. OF KANSAS
Govt. Documents & Maps Library
6001 Malott Hall
Lawrence, KS 66045-2800
(913) 864-4660 Fax: (913) 864-3855

KENTUCKY

UNIV. OF KENTUCKY
King Library South
Govt. Publications/Maps Dept.
Patterson Drive
Lexington, KY 40506-0039
(606) 257-3139 Fax: (606) 257-3139

LOUISIANA

LOUISIANA STATE UNIV.
Middleton Library
Govt. Documents Dept.
Baton Rouge, LA 70803-3312
(504) 388-2570 Fax: (504) 388-6992

LOUISIANA TECHNICAL UNIV.

Prescott Memorial Library
Govt. Documents Dept.
Ruston, LA 71272-0046
(318) 257-4962 Fax: (318) 257-2447

MAINE

UNIV. OF MAINE
Raymond H. Fogler Library
Govt. Documents Dept.
Orono, ME 04469-5729
(207) 581-1673 Fax: (207) 581-1653

MARYLAND

UNIV. OF MARYLAND - COLLEGE PARK
McKeldin Library
Govt. Documents/Maps Unit
College Park, MD 20742
(301) 405-9165 Fax: (301) 314-9416

MASSACHUSETTS

BOSTON PUBLIC LIBRARY
Govt. Documents
666 Boylston Street
Boston, MA 02117-0286
(617) 536-5400, ext. 226
Fax: (617) 536-7758

MICHIGAN

DETROIT PUBLIC LIBRARY
5201 Woodward Avenue
Detroit, MI 48202-4093
(313) 833-1025 Fax: (313) 833-0156

LIBRARY OF MICHIGAN

Govt. Documents Unit
P.O. Box 30007
717 West Allegan Street
Lansing, MI 48909
(517) 373-1300 Fax: (517) 373-3381

MINNESOTA

UNIV. OF MINNESOTA
Govt. Publications
409 Wilson Library
309 19th Avenue South
Minneapolis, MN 55455
(612) 624-5073 Fax: (612) 626-9353

MISSISSIPPI

UNIV. OF MISSISSIPPI
J.D. Williams Library
106 Old Gym Bldg.
University, MS 38677
(601) 232-5857 Fax: (601) 232-7465

MISSOURI

UNIV. OF MISSOURI - COLUMBIA
106B Ellis Library
Govt. Documents Sect.
Columbia, MO 65201-5149
(314) 882-6733 Fax: (314) 882-8044

MONTANA

UNIV. OF MONTANA
Mansfield Library
Documents Division
Missoula, MT 59812-1195
(406) 243-6700 Fax: (406) 243-2060

NEBRASKA

UNIV. OF NEBRASKA - LINCOLN
D.L. Love Memorial Library
Lincoln, NE 68588-0410
(402) 472-2562 Fax: (402) 472-5131

NEVADA

THE UNIV. OF NEVADA LIBRARIES
Business and Govt. Information Center
Reno, NV 89557-0044
(702) 784-6579 Fax: (702) 784-1751

NEW JERSEY

NEWARK PUBLIC LIBRARY
Science Div. - Public Access
P.O. Box 630
Five Washington Street
Newark, NJ 07101-7812
(201) 733-7782 Fax: (201) 733-5648

NEW MEXICO

UNIV. OF NEW MEXICO
General Library
Govt. Information Dept.
Albuquerque, NM 87131-1466
(505) 277-5441 Fax: (505) 277-6019

NEW MEXICO STATE LIBRARY

325 Don Gaspar Avenue
Santa Fe, NM 87503
(505) 827-3824 Fax: (505) 827-3888

NEW YORK

NEW YORK STATE LIBRARY
Cultural Education Center
Documents/Gift & Exchange Section
Empire State Plaza
Albany, NY 12230-0001
(518) 474-5355 Fax: (518) 474-5786

NORTH CAROLINA

UNIV. OF NORTH CAROLINA - CHAPEL HILL
Walter Royal Davis Library
CB 3912, Reference Dept.
Chapel Hill, NC 27514-8890
(919) 962-1151 Fax: (919) 962-4451

NORTH DAKOTA

NORTH DAKOTA STATE UNIV. LIB.
Documents
P.O. Box 5599
Fargo, ND 58105-5599
(701) 237-8886 Fax: (701) 237-7138

UNIV. OF NORTH DAKOTA

Chester Fritz Library
University Station
P.O. Box 9000 - Centennial and University Avenue
Grand Forks, ND 58202-9000
(701) 777-4632 Fax: (701) 777-3319

OHIO

STATE LIBRARY OF OHIO
Documents Dept.
65 South Front Street
Columbus, OH 43215-4163
(614) 644-7051 Fax: (614) 752-9178

OKLAHOMA

OKLAHOMA DEPT. OF LIBRARIES
U.S. Govt. Information Division
200 Northeast 18th Street
Oklahoma City, OK 73105-3298
(405) 521-2502, ext. 253
Fax: (405) 525-7804

OKLAHOMA STATE UNIV.

Edmon Low Library
Stillwater, OK 74078-0375
(405) 744-6546 Fax: (405) 744-5183

OREGON

PORTLAND STATE UNIV.
Branford P. Millar Library
934 Southwest Harrison
Portland, OR 97207-1151
(503) 725-4123 Fax: (503) 725-4524

PENNSYLVANIA

STATE LIBRARY OF PENN.
Govt. Publications Section
116 Walnut & Commonwealth Ave.
Harrisburg, PA 17105-1601
(717) 787-3752 Fax: (717) 783-2070

SOUTH CAROLINA

CLEMSON UNIV.
Robert Muldrow Cooper Library
Public Documents Unit
P.O. Box 343001
Clemson, SC 29634-3001
(803) 656-5174 Fax: (803) 656-3025

UNIV. OF SOUTH CAROLINA

Thomas Cooper Library
Green and Sumter Streets
Columbia, SC 29208
(803) 777-4841 Fax: (803) 777-9503

TENNESSEE

UNIV. OF MEMPHIS LIBRARIES
Govt. Publications Dept.
Memphis, TN 38152-0001
(901) 678-2206 Fax: (901) 678-2511

TEXAS

TEXAS STATE LIBRARY
United States Documents
P.O. Box 12927 - 1201 Brazos
Austin, TX 78701-0001
(512) 463-5455 Fax: (512) 463-5436

TEXAS TECH. UNIV. LIBRARIES

Documents Dept.
Lubbock, TX 79409-0002
(806) 742-2282 Fax: (806) 742-1920

UTAH

UTAH STATE UNIV.
Merrill Library Documents Dept.
Logan, UT 84322-3000
(801) 797-2678 Fax: (801) 797-2677

VIRGINIA

UNIV. OF VIRGINIA
Alderman Library
Govt. Documents
University Ave. & McCormick Rd.
Charlottesville, VA 22903-2498
(804) 824-3133 Fax: (804) 924-4337

WASHINGTON

WASHINGTON STATE LIBRARY
Govt. Publications
P.O. Box 42478
16th and Water Streets
Olympia, WA 98504-2478
(206) 753-4027 Fax: (206) 586-7575

WEST VIRGINIA

WEST VIRGINIA UNIV. LIBRARY
Govt. Documents Section
P.O. Box 6069 - 1549 University Ave.
Morgantown, WV 26506-6069
(304) 293-3051 Fax: (304) 293-6638

WISCONSIN

ST. HIST. SOC. OF WISCONSIN LIBRARY
Govt. Publication Section
816 State Street
Madison, WI 53706
(608) 264-6525 Fax: (608) 264-6520

MILWAUKEE PUBLIC LIBRARY

Documents Division
814 West Wisconsin Avenue
Milwaukee, WI 53233
(414) 286-3073 Fax: (414) 286-8074

National Aeronautics and
Space Administration
Code JTT
Washington, DC 20546-0001