

FINAL
IN - 10
018 875

FINAL REPORT

NAS5-32782

"Wind Variabilty in Intermediate Luminosity B Supergiants"

Submitted to: Steve Kramer
Code 216
NASA/Goddard Space Flight Center
Greenbelt, MD 20771

Submitted by: Dr. Derck Massa
Applied Research Corporation
8201 Corporate Drive
Landover, MD 20785
(301)459-8833

ARC Report #: R96-250

Date: April 26, 1996

Final Report

Contract: NAS5-32782

PI: Derck Massa

Title: Wind Variability in Intermediate Luminosity B Supergiants

1. Introduction

The object of this study was to determine the ubiquity and nature of wind variability in intermediate luminosity B supergiants. These stars were selected for study because, unlike the more frequently studied O and WR stars, they contain spectral diagnostics which present distinct advantages for addressing many observational issues concerning wind physics. Some of these advantages are:

1. **BIs have well-developed, but unsaturated wind lines that span a wide range of ionization stages.** These can run from C II to C IV for B2-5 and from Si III to N V at B0-1, and often include adjacent stages of ionization for Si (Si III $\lambda 1206$ and Si IV $\lambda\lambda 1400$). Furthermore, many of these ions are much closer to the stages expected to be dominant in the wind. The fact that the lines are frequently unsaturated allows more reliable optical depths to be determined and the fact that they are closer to the dominant ionization stage allows more accurate estimates of mass loss, \dot{M} , to be made since smaller adjustments are required for the ionization fraction, q , of the observed ionic species.
2. **BIs frequently have more than one unsaturated wind line.** This situation occurs often and opens up the possibility of studying how the ionization ratios in the winds change with respect to the physical parameters of the star and with time, in the case of variable winds.
3. **After hydrogen, the strongest photospheric lines of BIs are the UV silicon lines.** The strong UV photospheric silicon lines provide sensitive temperature and surface gravity diagnostics. The strongest photospheric lines also show wind effects and serve as excellent probes of the wind-photosphere interface.

4. **Abundance anomalies can be quantitatively measured in BI winds.** Because the photospheric UV silicon spectra of BIs make it possible to place their spectra onto a 2-dimensional silicon based grid unambiguously, one can study their wind lines independently. Typically, wind lines from carbon and nitrogen (when present) follow the trends indicated by the photospheric line strengths *and* the silicon wind lines. There are, however, distinct exceptions. The vast majority of these are related to abundance anomalies.
5. **BIs have longer wind flow times than O stars.** Because they have lower terminal speeds and larger radii than O stars, their wind flow times ($t_{flow} \propto R/v_{\infty}$ for a steady wind) are longer. This presents two distinct advantages. First, the spectroscopic signatures of their wind variability evolve more slowly. Thus, wind variations in BIs are easier to observe, especially at low velocities where the accelerations are largest. Second, the lower terminal velocities allow the two components of the important Si IV $\lambda\lambda 1400$ doublet to become largely uncoupled.

2. Work Completed

The following work was completed under this contract:

1. A detailed analysis of wind variability in the B0.5 Ib star HD 64760 (Massa et al. 1995b, enclosure 1) which demonstrated:
 - The presence of massive ejections of material into the wind which involve the photosphere.
 - The first clear demonstration of the long sought after “photospheric connection” between wind and atmospheric lines.
 - The development of new analysis tools to examine the ionization in the wind and a new model fitting procedure.
 - A clear demonstration of whole scale shifts in the ionization of the normal stellar wind.
2. Organized and lead the large, international effort, dubbed the “*IUE* MEGA campaign” (Massa et al. 1995a, enclosure 2) which obtained unprecedented temporal coverage of wind variability in rapidly rotating, early-type stars. The results of this aspect of the study included

- The demonstration of recurring wind features which repeated with a period equal to an one fourth of the rotation period of a B supergiant. This result provided the first clear indication of wind enhancements whose origin is rooted in the stellar photosphere (Prinja, Massa & Fullerton 1995, enclosure 3).
 - The periodic statistical behavior of wind enhancements in an O supergiant (Howarth et al. 1995, enclosure 4).
3. A detailed analysis of wind variability in the rapidly rotating B1 Ib, γ Ara (Prinja, Massa & Fullerton 1996, enclosure 5). This analysis demonstrated
 - The existence of a two component wind – with the denser, more slowly evolving component most probably associated with a wind compressed disk.
 - The discovered a stellar wind in a normal B supergiant which has distinctly different mean states from one epoch to the next.
 4. Organized a follow on campaign to the MEGA project, designed to study slowly rotating O and B supergiants. This campaign was deemed a program of lasting value by the joint ESA/NASA review board, and will obtain 30 days of *IUE* observations of 4 OB stars. This program will be performed in May-June of 1996.
 5. Performed a global survey of all the available *IUE* time series for B supergiants (Massa, Prinja, & Fullerton 1995, BAAS, 27, 1311, enclosure 6), and identified recurring spectroscopic signatures which were tentatively identified with different physical phenomena.

3. Work in Progress

The following work is still in progress.

- **Completion of item 5, the overall survey:** This final bit of work is still in progress, and will help to place the previous results into context. This work will be pursued this summer with Raman Prinja at University College, London. This work will be done in conjunction with a NATO visiting Scientist grant which will pay my living and travel expenses.

- **The follow on to the IUE MEGA campaign (item 4):** The observations from this project will be obtained this summer and will represent the longest time series ever obtained for stellar winds (a record which, unfortunately, will probably last for many years because *IUE* is scheduled to be decommissioned this fall) and it is extremely important to make these data available to the community. This will be one of my objectives during my stay in London this summer.

Publications resulting from this contract:

- “Stellar Wind Variability: the Things B Supergiants Do”, D. Massa, R.K. Prinja, & A. Fullerton 1995, BAAS, 27, 1311
- “Wind Variability in B Supergiants: II. The Two Component Stellar Wind of γ Ara (B1 Ib)”, R.K. Prinja, D. Massa, & A. Fullerton 1996, in press
- “The *IUE* MEGA Campaign: Wind Variability and Rotation in Early-Type Stars” D. Massa, et. al. 1995 ApJ, 452, L53
- “The *IUE* MEGA Campaign: Modulated Structure in the Wind of HD 64760 (B0.5 Ib)” R.K. Prinja, D. Massa, & A.W. Fullerton 1995 ApJ, 452, L61
- “The *IUE* MEGA Campaign: The Rotationally Modulated Wind of ζ Puppis” I.D. Howarth, R.K. Prinja, & D. Massa 1995 ApJ, 452, L65
- “Wind Variability in B Supergiants: I. The Rapid Rotator HD 64760 (B0.5 Ib)”, D. Massa, A. Fullerton & R.K. Prinja 1995, ApJ, 452, 842.

Enclosure 1

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 1996	3. REPORT TYPE AND DATES COVERED Contractor Report	
4. TITLE AND SUBTITLE Wind Variability in Intermediate Luminosity B Supergiants			5. FUNDING NUMBERS Code 681 Contract: NAS5-32782	
6. AUTHOR(S) PI: Dr. Derck Massa				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Applied Research Corporation 8201 Corporate Drive Landover, MD 20785			8. PERFORMING ORGANIZATION REPORT NUMBER R96-250	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA Aeronautics and Space Administration Washington, D.C. 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER CR-203673	
11. SUPPLEMENTARY NOTES Technical Monitor: D. West, Code 681				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category: 89 Report available from the NASA Center for AeroSpace Information, 800 Elkridge Landing Road, Linthicum Heights, MD 21090; (301) 621-0390.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This study used the unique spectroscopic diagnostics of intermediate luminosity B supergiants to determine the ubiquity and nature of wind variability. Specifically, 1) A detailed analysis of HD 64760 demonstrated massive ejections into its wind, provided the first clear demonstration of a "photospheric connection" and ionization shifts in a stellar wind. 2) The international "IUE MEGA campaign" obtained unprecedented temporal coverage of wind variability in rapidly rotating stars and demonstrated regularly repeating wind features originating in the photosphere. 3) A detailed analysis of wind variability in the rapidly rotating B1 Ib, gamma Ara demonstrated a two component wind with distinctly different mean states at different epochs. 4) A follow-on campaign to the MEGA project to study slowly rotating stars was organized and deemed a key project by ESA/NASA, and will obtain 30 days of IUE observations in May-June 1996. 5) A global survey of archival IUE time series identified recurring spectroscopic signatures, identified with different physical phenomena. Items 4 and 5 above are still in progress and will be completed this summer in collaboration with Raman Prinja at University College, London.				
14. SUBJECT TERMS Early-type Stars, Stellar Winds, Wind Variability, IUE			15. NUMBER OF PAGES 62	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	