

**STUDY OF ELONGATED SUPERBUBBLE AROUND CYG OB1**

NASA Grant No. NAG5-12612

**Final Report**

For Period 1 January 2003 through 31 December 2004

Principal Investigator  
Joy Nichols

October 2004

Prepared for:

National Aeronautics and Space Administration  
Goddard Space Flight Center  
Greenbelt, MD 20771

Smithsonian Institution  
Astrophysical Observatory  
Cambridge, Massachusetts 02138

The Smithsonian Astrophysical Observatory  
is a member of the  
Harvard-Smithsonian Center for Astrophysics

The NASA Technical Officer for this Grant is George Sonneborn, Code 681, NASA-Goddard Space Flight Center, Greenbelt, MD 20771

# Study of the Elongated Superbubble Around Cyg OB1

Joy Nichols, PI, FUSE Cycle 3

## 1 Introduction

OB association superbubbles and supernova remnants are believed to be the primary sources of the hot component of the interstellar medium. Superbubbles are also thought to “blow out” of the galactic plane and are thus the potential source of galactic coronal gas. Quantifying the energetics and accurately modeling the evolution of superbubbles is thus an important task in the understanding of the evolution of our Galaxy.

The superbubble surrounding the Cyg OB1 association (Figure 1) is an excellent example for study of the early stages of superbubble evolution. This superbubble, discovered in IRAS data (Saken et al. 1992, Nichols-Bohlin & Fesen 1993), is an elongated structure 2x5 degrees (26x65 pc at distance=1.5 kpc), with at least 2 known supernova remnants believed to be associated with the superbubble. The elongated shape is more extended than can be explained by differential galactic rotation. A major problem in understanding the evolution of the Cyg OB1 superbubble is the discrepancy between the age of the superbubble and the derived age of the parent star association. The superbubble is believed to be no more than 1 Myr old, while the association is about 4-5 Myr old (Saken et al. 1992).

## 2 Observations and Analysis

Four spectra of hot stars in the line of sight to the IR-detected Cyg OB1 superbubble were obtained with FUSE using the LWRS aperture and TTAG mode. Table 1 lists the star name, spectral type, total counts, and total exposure time for each observation.

Table 1. FUSE Observations of Cyg OB1 Stars

Star	Sp. Type	Total Counts	Total Exposure Time (sec)
HD 192639	O7.5 IIIf	603525	3117
HD 192641	WC7	353624	4517
HD 193077	WN6+abs	1828858	21650
HD 192303	B1 III	178558	3635

Figure 1 shows a 60/100  $m\mu$  IRAS image of the Cyg OB1 superbubble, with the positions of our program stars indicated. Two of the stars are near the center of the superbubble, while 2 are near the northern and southern edge, respectively.

The spectra were first reprocessed using CalFUSE 3.0.2 to take advantage of the improvements in this version of the pipeline processing, particularly the intermediate data products. The data were then analyzed using IDL tools distributed by the FUSE Science Center.

We examined the interstellar absorption lines for evidence of high velocity features which could be attributed to the superbubble expansion. We found multiple high velocity features in the O VI lines, Fe II, Cl II, N II, P V, and S III.

### 3 Results

In the attached proceedings publication, we present the preliminary results for analysis of the O VI high velocity features of the four stars observed.

In a paper about to be submitted to ApJ, we report the velocity and column density measurements of all the interstellar high velocity features we were able to identify. We have used these FUSE data, as well as IUE data for C IV and N V, to constrain shock models of the superbubble using 2D MHD code.

The conclusions of this work are that the Cyg OB1 superbubble was created by non-coeval supernovae and massive star wind generation within a multicluster region, including Cyg OB1, OB3, and OB9. These clusters must share a peculiar space velocity of approximately +60 km/s for the energy budget and the shock speed to be consistent with the FUSE data. We find that the superbubble is still being formed from star formation regions along its rim, and that the multiple superbubbles have merged and swept out the interior dust, which results in the low IR emission seen in Figure 1.




Figure 1: Cyg OB1 superbubble as seen in 60/100  $m\mu$  IRAS ratio image. Background stars observed with FUSE are indicated on the figure.

## FUSE Analysis of the Cyg OB1 Superbubble

J. Nichols and P. Mendygral

*Harvard-Smithsonian Center for Astrophysics, 60 Garden St.,  
Cambridge, MA 02138*

**Abstract.** FUSE observations of four stars in the line of sight to the Cyg OB1 IR-detected superbubble have been analyzed for high-velocity features in the O VI interstellar lines, which might be attributed to the shock structure of the superbubble. Multiple components were detected in the spectra of all four stars, with a velocity range of  $-85$  km/s to  $+29$  km/s. As many as four separate velocity components were identified in each spectrum, implying multiple shock structures in the superbubble. Derived column densities of the O VI components indicate shock velocities of 160-190 km/s according to steady state shock theory.

### 1. Background

The Cyg OB1 superbubble is an elongated structure that overlaps the boundaries of the associations Cyg OB1, OB3, and OB9 (Saken et al. 1992). The size is  $2 \times 4.5$  degrees; at an estimated distance of 1.5 kpc the physical size is  $52 \times 118$  pc. This superbubble is not the same as the well-known Cygnus superbubble which is much larger and includes the association Cyg OB2, but it may be an extension or "blister" of this larger superbubble.

The Cyg OB1 superbubble is best identified in the infrared (Saken et al. 1992; Nichols-Bohlin & Fesen 1993), although evidence of portions of this superbubble have been detected in the optical (Lozinskaya & Repin 1990). Several theories have been proposed to explain the elongated shape of this superbubble. It could be due to non-coeval evolution of the Cyg OB1 association (Saken et al. 1992), or possibly multiple superbubbles that have merged.

The Cyg OB1 superbubble is believed to be in the early stages of evolution (Saken et al. 1992). Nichols-Bohlin and Fesen (1993) estimated the total energy output of Cyg OB1 and, assuming the superbubble is a result of the winds of the member stars and the estimated 4 supernovae that would have occurred during the 2 million year lifetime of Cyg OB1, calculated that the superbubble is about  $2 \times 10^6$  yr old and has an expansion velocity of about 80 km/sec.

Of the 22 stars in the direction to the Cyg OB1 superbubble having IUE high dispersion observations available, 18 show high-velocity components in both low and high ionization interstellar lines (Nichols-Bohlin & Fesen 1993). The velocity of the components detected in the IUE spectra span the range  $\pm 90$  km/sec, although negative velocities predominate the detections. The multiple components to the interstellar lines, coupled with the lack of geometric velocity gradient across the superbubble, imply that there are several overlapping superbubbles in this line of sight, including a probable multi-association superbubble.

## 2. Observations

Four stars in the line of sight to the Cyg OB1 superbubble have been observed with FUSE. The observations were made using the LWRS in TTAG mode. The data were reprocessed with CalFUSE 3.0.2 and smoothed with a 3-point median filter. Only the LiF1A spectra were included in this analysis in order to obtain the best spectral resolution. Gaussian profiles were fit to the components of the O VI absorption line at 1031 Å. The four stars observed are all members of the Cyg OB1 association, at an estimated distance of 1.5 kpc. Two of the stars (HD 192639 and HD 193077) are near the center of the superbubble, and the other two are near the northern and southern edges of the superbubble, as mapped in the infrared data. Figure 1 shows the smoothed FUSE spectra in the O VI region, with the high-velocity features marked.




Figure 1. FUSE spectra of four stars observed in the line of sight to Cygnus OB1 superbubble, with indications of the high-velocity components in the O VI interstellar lines

## 3. Comparison to Shock Models

The O VI column density is very sensitive to shock speed in the steady state shock models in the range of 140-190 km/sec. By comparison of the O VI column density to the steady state shock model predictions, we can evaluate the consistency of each component, both low- and high-velocity, to the expected column density. Table 1 lists each component detected in the O VI line for each

star, spectral type, radial velocity, and equivalent width of the components, the derived column density, and the model prediction of the shock speed.

Table 1. O VI Column Densities

Star	Sp. Type	Velocity of Component (km/sec)	Equivalent Width (mÅ)	N(O VI)	Model Shock Speed (km/sec)
HD 192639	O7.5 IIIf	-83	103	$8.23 \times 10^{13}$	169
		-48	43	$3.43 \times 10^{13}$	159
		-16	211	$1.68 \times 10^{14}$	185
		+27	57	$4.60 \times 10^{13}$	162
HD 192641	WC7	-69	98	$7.83 \times 10^{13}$	169
		-40	22	$1.76 \times 10^{13}$	154
		-14	189	$1.51 \times 10^{14}$	184
HD 193077	WN6+abs	-74	88	$7.03 \times 10^{13}$	169
		-28	123	$9.82 \times 10^{13}$	171
HD 192303	B1 III	-85	62	$4.95 \times 10^{13}$	188
		-19	253	$2.02 \times 10^{14}$	166

#### 4. Results

- High-velocity components in the O VI lines have been found in all four stars observed toward the Cygnus OB1 superbubble
- As in the IUE data, no clear geometric effects are apparent in the data between stars projected near the edge of the superbubble and stars projected near the center. Further analysis may reveal such a gradient.
- Steady state shock theory predicts shock speeds of 160-190 km/sec for the O VI column densities we have measured. These shock speeds are plausible if the Cyg OB1 association as a large differential galactic velocity, as has been proposed.
- Multiple components to the O VI lines imply multiple shocks along the line of sight to Cyg OB1.

**Acknowledgments.** We acknowledge the support of NASA Grant NAG5-12612.

#### References

- Esipov, V. F., Lozinskaya, T. A., Mel'nekov, V. V., Pravdikova, V. V., Sitnik, T. G., and Nichols-Bohlin, J. 1996, *AstL*, 22, 571
- Lozinskaya, T. A. and Repin, S. V. 1990, *AZh*, 67, 1152L
- Nichols-Bohlin, J. and Fesen, R. A. 1993, *AJ*, 672, 302
- Saken, J. M., Shull, J. M., Garmany, C. D., Nichols-Bohlin, J., and Fesen, R. A. 1992, *ApJ*, 397, 537