

7. Impacts of a Changing Climate and Land Use on Reindeer Pastoralism: Indigenous Knowledge & Remote Sensing

Maynard, N.G.¹, Oskal², A., Turi², J.M., Mathiesen^{2,3}, S.D., Eira³, I.M.G., Yurchak¹, B., Etylin⁴, V., Gebelein⁵, J.

¹NASA Goddard Space Flight Center, Cryospheric Sciences Branch, Code 614.1, Greenbelt, MD 20771 USA

²International Centre for Reindeer Husbandry, Boaranjarga 1, N-9521, Guovdageaidnu/Kautokeino, Norway 9520

³Sami Allaskuvla/Sámi University College, Guovdageaidnu/Kautokeino, Norway 9520.

⁴Chukotka Branch of the North-Eastern Research Institute. Russian Academy of Science, Anadyr, Chukotka, Russia

⁵Department of International Relations & Geography, Florida International University, Miami, FL. 33199, USA

7.1. Introduction

7.1.1 Reindeer Pastoralism and Arctic Changes

The Arctic is home to many indigenous peoples, including those who depend on reindeer herding for their livelihood, in one of the harshest environments in the world. For the largely nomadic peoples, reindeer not only form a substantial part of the Arctic food base and economy, but they are also culturally important, shaping their way of life, mythologies, festivals and ceremonies. Reindeer pastoralism or husbandry has been practiced by numerous peoples all across Eurasia for thousands of years and involves moving herds of reindeer, which are very docile animals, from pasture to pasture depending on the season. This means herders lead either a nomadic life living in a tent on the Arctic tundra year-round as a family unit, or a semi-nomadic life, having permanent residences for parts of the year and having fewer family members herding on a daily basis. Thus, herders

must adapt on a daily basis to find optimal conditions for their herds according to the constantly changing conditions.

Having learned over generations to live with uncertainties in an Arctic environment, societies that practice traditional reindeer husbandry are good examples of sustainable human communities that are highly interconnected with the ecosystems in which they live. Climate change and variability plus rapid development are increasingly creating major changes in the physical environment, ecology, and cultures of these indigenous reindeer herder communities in the North, and climate changes are occurring significantly faster in the Arctic than the rest of the globe, with correspondingly dramatic impacts. (Oskal, 2008)

Fig. 1 (Reindeer peoples of the world)

In response to these changes, Eurasian reindeer herders have created the EALAT project, a comprehensive new initiative to study these impacts and to develop local adaptation strategies based upon their traditional knowledge of the land and its uses – in targeted partnership with the science and remote sensing community - involving extensive collaborations and co-production of knowledge to minimize the impacts of the various changes. This unprecedented new reindeer herder-led initiative has resulted in the development of an international, interdisciplinary collaboration with scientists through the International Polar Year (IPY) EALAT Consortium (IPY Project # 399 “EALAT, Reindeer Pastoralism in a Changing Climate”) and directly addresses the herders’ need for additional data and information for responding to the global and environmental changes through a variety of different projects. The EALAT initiative is considered unique because the study was conceived and implemented by indigenous reindeer herders who continue to lead this large multidisciplinary group of invited scientists and other collaborators in this complex study of the multiple challenges facing Arctic reindeer herding communities.

This chapter provides background on climate and development challenges to reindeer husbandry across the Arctic and an overview of the EALAT initiative, with an emphasis on indigenous knowledge, remote sensing, Geographic Information Systems (GIS), and other scientific data to “co-produce” datasets for use by herders for improved decision-making and herd management. It also provides a description of the EALAT monitoring data integration and sharing system and portal being developed for reindeer pastoralism. In addition, the chapter provides some preliminary results from the EALAT Project, including some early remote sensing research results.

7.1.2 Reindeer Pastoralism across the Arctic – Background and Challenges

Reindeer husbandry has a long history in the Arctic. There are more than 20 different indigenous peoples in the Arctic who are reindeer herders. Reindeer husbandry is practiced in Norway, Sweden, Finland, Russia, Mongolia, China, Alaska, Canada and Greenland. This livelihood involves some 100,000 herders and approximately 2.5 million semi-domesticated reindeer, which graze approximately 4 million square kilometers in Eurasia. While reindeer husbandry is spread across the Arctic and across many cultures, its organization is remarkably similar everywhere, consisting of a nomadic livelihood with family-based working communities and a typical indigenous way of life. For hundreds of years, reindeer herders have managed vast barren circumpolar areas of land that hold little value for others. Herding represents a model for sustainable management of these areas that has developed through generations. In recent years, however, as noted earlier, Arctic reindeer herders increasingly face major challenges, such as climate change, loss of grazing land due to development by humans, and effects of global change in their local societies. (Oskal, 2008)

Reindeer/caribou are the very base of the traditional economy for many indigenous northern peoples across the Arctic. *Rangifer tarandus*, called reindeer or caribou, is the most common large land mammal of the Arctic and SubArctic, which gathers in large herds of tens to hundreds of thousands of animals on their calving grounds during the Arctic summer, and scattering widely in smaller groups for the remainder of the year (Hall, 1989). In Russia, the total number of domesticated reindeer in the region has decreased significantly within the last 100 years with particularly marked change from approximately 2.5 million in 1969 to 1.2 million in 2000 (Jernsletter and Klokov, 2002). This decline in numbers of reindeer in Northern Russia, which has by far the largest share of pasture lands (87%) and about 67% of all reindeer, is causing a serious decline in the reindeer husbandry industry, and, in turn is directly affecting the health and well-being of the indigenous peoples associated with reindeer husbandry (Schindler, D.L., 1994; Jernsletten and Klokov, 2002; Nuttal et al, 2005). This decline is not only causing poverty in the Russian indigenous communities associated with reindeer herding, but also, because reindeer husbandry is the very core of their traditional way of life, the decline is causing serious damage to the ethnic traditions and to the families of nomadic reindeer herders. (Abrjutina, 2003; Jernsletten and Klokov, 2002; Glazovsky et al, 2004; Nuttal et al, 2005; Klokov, 2000).

In Norway, reindeer husbandry is one industry in which the number of people involved has increased over the past 50 years (Eira, A.J., 2001).

However, while approximately 40% of mainland Norway is designated reindeer pastureland, there are serious threats to those lands from not only climate changes, but also loss of pastures by increasing encroachment from development, tourism, damming of rivers, cultivation, oil and gas development, and roads and power lines, accompanied by similar impacts as are observed in Russia (Eira, A.J., 2001). In Finnmark, which is the northernmost, largest and least populated county in Norway, there are approximately 2000 registered reindeer owners which represent 73% and 75% of semi-domesticated reindeer and Sami reindeer owners in Norway, respectively (Tyler et al, 2007).

In recent years, a number of studies have been published on historical and future challenges which relate directly to reindeer herding in Eurasia including – as a few examples – Eira et al (2008), Nuttal et al (2008; 2005), Klein et al (2005), a diverse collection of multidisciplinary articles summarizing results from the RENMAN Project (“The challenges of modernity for REiNdeer MANagement: integration and sustainable development in Europe’s subarctic and boreal regions” by Forbes et al, 2006), the ENSINOR Project, (Stammler and Burgess, 2007), Walker et al (this volume), Jia et al (2003), Tommervik et al (2004), Hinzman et al (2005), Kittl et al (2006), Kumpala et al (2006), Tape et al (2006), Stammler and Peskor (2008).

7.1.3 Reindeer, Climate Change and Development

As mentioned in Chapter 1, climate change is happening faster than in any other regions of the world. The changes in ice cover and increases in temperature have already impacted reindeer husbandry and will continue to do so both directly, for example through changes in food availability, and indirectly such as through changes in human land use. (Oskal, 2008) Temperature changes have begun to cause some rivers to freeze later in the autumn and melt earlier in the spring, resulting in challenges for the annual migration of reindeer between different seasonal pastures. Warming-induced changes in freeze-thaw cycles are also creating problems. For example, as river and lake ice thaws earlier in the spring along migration routes, newborn calves can no longer cross the ice surface, but have to attempt crossing open waters, and large numbers of calves have been swept away by currents (Klein et al, 2005; Nuttal et al, 2005). Another change that has already been observed is increasing climate variability at a local level. This is especially true during the critical wintertime, where increasingly periods of mild weather accompanied by rain will be followed by colder periods, form ice layers in the snow and block the reindeers’ access to food on the ground. As reindeer live only on natural pastures, this often

represents a “worst-case scenario” from the reindeer herders’ perspective. Increasing precipitation in the form of snow can add to these challenges, while warming would shorten the period of snow cover in any particular year. (Oskal, 2008)

Fig. 2. Future warming in Norway....

A deeper snow pack in winter can also make the reindeer more vulnerable to predator attacks (e.g., wolves) because the lighter wolves can travel on thinner snow crusts that reindeer sink through (Brotton and Wall, 1997). Increased insect harassment, accompanying warmer temperatures, is a second major factor shown to interfere with foraging. (Kitti et al, 2006) The outcome of this harassment is increased energy requirements, and results in a significant decline in body fat and lactation and decrease in calving success (Walsh et al 1992; Brotton and Wall, 1997; Gunn and Skogland, 1997). One example of recent climate impact is the unusually warm winter of 1996-1997, which was associated with a deep snow pack and icing, and which caused about 10,000 reindeer to die of starvation on Russia’s far northeast Chukotsk Peninsula. (Malcolm, 1996; Nuttall et al, 2005)

Reindeer herders have also observed major changes in biodiversity. A significant example of this is repeated occurrences of certain species replacing others, such as the spreading of shrubs into the barren tundra-areas (Jia et al, 2003; Hinzman et al, 2005; Tape et al, 2006). Shrubs contribute to a hard packing of snow during the tough winter months, thus making access to food a challenge for reindeer. In addition, important food resources for the reindeer, such as lichens and reindeer preferred species of grasses, in time may disappear partially if not fully due to this shrub encroachment. Changes and/or increases in insect populations could also change reindeer behaviour during the summer by not allowing them to feed long enough in summer pastures due to increased harassment. (Oskal, 2008; Kitti et al, 2006)

Fig. 3 Climate variability in the Arctic....

Indirect effects of climate change are also being observed, with major implications for reindeer pasture availability and migration routes. (Kitti et al, 2006) Due to the sea ice melting and longer summers, increased accessibility of the Arctic regions for human activities is a growing threat to reindeer herders. Human development and activities represent disturbances with negative effects for the semi-domesticated reindeer herds (Kitti et al, 2006) and irreversible loss of marginal pasture resources – a serious challenge for reindeer husbandry. In particular, female reindeer and calves will stay away from humans, physical installations and general human activity. In the last 50 years, for example, approximately 25 % of the

reindeer pastures of the Euro-Arctic Barents Region have in effect been lost due to human development (Tyler et al, 2007).

Of particular relevance today is the fact that the Arctic is estimated to contain approximately 25 % of the world's remaining undeveloped petroleum resources (Forbes, 2000). For instance, Yamal in Western Siberia holds about 90 % of Russia's gas reserves, while also being the largest reindeer herding area of the world. Activities to access these resources would reduce the grazing lands, and are viewed as another human activity in the Arctic contributing to the reduction of the "available room for adaptation" for reindeer husbandry (Forbes, 2000; Nuttal et al, 2005). In fact, industrial development (e.g., pipelines and oil and gas infrastructure) has increased across reindeer migration routes in Northern Russia, blocking pathways to summer pasturelands (Forbes et al, 2006; see also Walker et al, this volume)).

It is also expected that there will be a sharp increase in the near future in oil and gas development, mining, and other forms of development in the Russian North – accompanied by infrastructure, pollution, and other manifestations of human presence – which will increase future pressure on available pasturelands for the reindeer and the indigenous communities associated with them. (Forbes, 2000; Forbes et al, 2006; Jernsletten and Klovov, 2002; Derome and Lukina, this volume) Furthermore, future reductions in sea ice from global warming recently projected are very likely to increase the amount of marine traffic and general access to the Arctic and, as a result, significantly increase development as well as serious problems related to sovereignty, social, cultural and other environmental issues, which will directly impact the indigenous reindeer herding community. (McCarthy et al, 2005)

7.1.4 Socioeconomic, Political and Other Pressures

Parts of the Arctic are unique in terms of the political settlements and land claims that have been achieved over the last thirty years or so. The extent of vulnerability and resilience to climate change not only depends on cultural aspects and ecosystem diversity, but on the political, legal and institutional rules which govern social-economic systems and social-ecological systems. (Nuttal et al, 2005, 2008) On the one hand, climate change has the potential to enhance economic development, but with further climate change, the climate in the Arctic is predicted to become more variable and extreme weather events more frequent and severe, which on the other hand, can undermine economic activities. Thus, it seems particu-

larly important that attention be given to the management of resources and to the effectiveness of governance institutions, and critical questions must be asked as to whether they can create additional opportunities to increase resilience, flexibility and the ability to deal with change. (Nuttal et al, 2005)

Compounding the problem for the reindeer herding community in Russia, the transition of Russia to a market economy has, over the past few years, resulted in considerable disorder in many parts of the supply and transport systems in remote northern areas. This has resulted in serious disruption of any system of goods, services, and health care to northern Russian indigenous Peoples. (Nuttal et al, 2005; Abrjutina, 2003; Jernsletten and Klovov, 2002; Klovov, 2002) Basic commodities such as paraffin lighting, fabrics, and vegetables or other foods are no longer easily available. The reindeer herders have also been cut off from any health care services at all, and, as a result of these factors combined, there are rapidly deteriorating health and living conditions in the reindeer herder communities, with growing death rates and serious health impacts. (Nuttal et al, 2005; Abrjutina, 2003; Jernsletten and Klovov, 2002; Klovov, 2000)

For all of these reasons, the Arctic Council has called for the full attention of the international community to the situation in the reindeer herding industry and the critical state of the indigenous peoples of the North in Russia. The 2nd World Reindeer Herders' Congress (2003) reported that there is a "real threat of the complete loss of reindeer husbandry in large parts of eastern Russia" and "indigenous peoples connected with reindeer husbandry here face an ethnic disaster". These concerns were echoed once again at the 3rd World Reindeer Herders' Congress in March 2005. Since that time, the reindeer herding communities have increasingly continued to develop new partnerships and organizations to improve their collective abilities to respond to the challenges of climate change and development. Indeed, as a result, the interdisciplinary multiparty EALAT Project was successfully launched by the Association of World Reindeer Herders in partnership with the Russian Union of Reindeer Herders, the Sami Reindeer Herders Association of Norway, and the Sami Council to address these many threats to reindeer herding through collaborative efforts to help prepare reindeer herders in Eurasia, their societies, institutions, and management for change, and accordingly, begin to reduce their vulnerability to these changes.

7.2 IPY EALAT Project: “Reindeer Pastoralism in a Changing Climate”

7.1.3 IPY EALAT: Overview

Developed under the leadership of indigenous reindeer herders, the IPY EALAT Project (“Reindeer Pastoralism in a Changing Climate”) is an interdisciplinary, intercultural study that is assessing the vulnerability of reindeer herding - a coupled human-ecological system - to changes in key aspects of the natural and human environments, actively involving reindeer herders, linguists, remote sensing scientists, meteorologists, lawyers, anthropologists, biologists, geographers, philosophers (the ethical dimension) as well as indigenous institutions and organizations, relevant industrial enterprises and management authorities. The name of the project, EALAT, which comes from the word "pasture" in the Sámi language, reflects the emphasis of the project on the close connection these cultures have to the environment in which they live. It focuses on the adaptive capacity of reindeer pastoralism to climate variability and change and, in particular, on the integration of reindeer herders’ knowledge with scientific research and analysis of their ability to adapt to environmental variability and change. (www.EALAT.org)

Fig. 4 Picture 4 reindeer feeding (heads in snow)

The IPY EALAT Project was initiated by the Association of World Reindeer Herders (WRH), a circumpolar indigenous peoples’ organization with observer status in the Arctic Council. The project leaders believe that valuing traditional and scientific knowledge equally and, hence, integrating herders’ experience and competence within the scientific method will enable it to contribute towards reducing the vulnerability of reindeer husbandry to the effects of climate change, which are likely to be pronounced over reindeer pastures in the north. *The EALAT-Network* study has adopted a multicultural approach in a multidisciplinary field that includes monitoring, research, outreach and communication.

To accomplish the goals of the EALAT Project, there are seven core “Work Packages”, which are self-contained areas of research and investigation that cover the wide range of issues that EALAT addresses. They are:

1. Identification of local climate conditions important for reindeer herding and development of basic climate scenarios
2. Customization of pasture conditions for reindeer pastoralism

3. Reindeer herders' knowledge: codifying and communicating coping mechanisms
4. Social and economic adaptation, - institutions and governance as constraints and opportunities
5. Reindeer: consequences of climate variability and change
6. Reindeer welfare and nutrition: herders' observations and scientific data
7. Synthesis: assessing vulnerability

EALAT-Outreach has as its objective to communicate Arctic reindeer herders' traditional knowledge and scientific knowledge related to climate change to herders and the mainstream society in the Arctic. *The Reindeer Portal* (www.reindeerportal.org) is a multiplatform web portal into the world of reindeer husbandry. Developed by the International Centre for Reindeer Husbandry in Kautokeino, the goal of the Reindeer Portal is to be a one stop shop information site for reindeer herders, students, administrators, politicians, indigenous people, business interest, the general public and anyone that is remotely interested in reindeer and the peoples that herd them.

In the IPY EALAT Consortium, *EALAT-Monitoring* is already endorsed as a future expert monitoring network in the Circumpolar Biodiversity Monitoring Program of the CAFF Working Group of the Arctic Council (CAFF, 2006). The data collected in IPY will be the start of a future place-based monitoring system of reindeer herders' pastures and societies, while at the same time representing a unique opportunity for validation of satellite imagery in cooperation with the NASA LCLUC program.

The legacy of IPY EALAT will be continued through a UArctic Institute for Reindeer Husbandry (UArctic EALAT) hosted in Kautokeino, Norway, as a pilot institute for research, outreach and education. For detailed information on EALAT, the reader is referred to www.EALAT.org.

7.2.2 IPY EALAT: Goals

The primary goals of the IPY EALAT Project are to assess the vulnerability of reindeer herding - a coupled human-ecological system - to climate and other changes in key aspects of the natural and human environments and to build optimal adaptive strategies through the integration of indigenous reindeer herder knowledge with scientific data and analyses. The IPY EALAT project partners believe that it is critical to empower indigenous peoples in Eurasia with the best technologies available to combine with indigenous knowledge for achieving a truly sustainable development of the Arctic. The EALAT team also believes it is important that

indigenous peoples' traditional knowledge must be a critical component of the future management and monitoring of the reindeer pastures and their societies. Therefore, another important goal of EALAT is to build competence locally about land cover land use change, including tools such as remote sensing observations and geographic information systems (GIS) as one important factor which could increase future adaptive capacity locally in Eurasian reindeer herding societies. Thus, developing training programs as well as EALAT monitoring systems for Eurasian reindeer herders, which will extend well beyond the international Polar Years (IPY) are a high priority.

For adaptation questions, *EALAT-Research* uses a vulnerability framework to assess the degree to which reindeer pastoralism is likely to experience harm as a result of exposure to multiple and interacting forces of change. The goal of the *EALAT-Outreach* component of the project is to communicate Arctic reindeer herders' traditional knowledge and scientific knowledge related to climate change to herders and the mainstream society in the Arctic.

In recent years, traditional knowledge has increasingly become an integral part scientific studies and a number of projects have included indigenous scientists and reindeer herders as full participants in project planning, implementation and dissemination of studies based on indigenous knowledge of modern reindeer management. (Heikkila, 2006; Hukkinen et al, 2006; Jaasko et al, 2006; Kitti et al, 2006; Magga, 2006; Muller-Willie et al, 2006; Stammer and Peskov, 2008) However, the EALAT project is considered unique because indigenous reindeer herders organized and are leading this complex, interdisciplinary, intercultural study, inviting scientists and other colleagues to collaborate. In this effort to inform the Arctic nations both about the changes to which they are subjected and give some concrete examples how herders' traditional knowledge relates to adaptation to changing conditions, another important goal of the study includes the challenge of taking reindeer herders' knowledge into action for sustainable development of the Arctic.

7.2.3 IPY EALAT Project: Study Sites

The IPY EALÁT project has research, information and outreach activities in five different reindeer herding societies across Eurasia; Nenets, Yamal-Nenets Autonomous Okrug, Sahka-Yakuta (64° 04'20.35" N 135° 58'03.24" E) and Chukotka (65°36.055' N 175°51.499') in Russia and Saami in Northern Europe. Researchers will concentrate initially on the two largest reindeer herding cultures in the world: The Sámi, who inhabit Northern Europe and the Kola Peninsula in North West Russia, with a fo-

cus on the Norwegian county of Finnmark, and the Nenets, focusing in particular on herders in the Yamal-Nenets Autonomous Okrug in Russia. Future research in the other regions where IPY EALAT has carried out information activities, will be implemented as part of the IPY EALAT legacy.

Fig. 5. Map of the 5 EALAT Eurasian study sites...

7.3 EALAT Studies

7.3.1 EALAT: Results from Early Studies

7.3.1.1 Early EALAT Results: SAR Studies for Pasture Quality

This section describes some results from Reindeer Mapper, an early EALAT pilot project of the NASA Land Cover Land Use Change Program, to investigate the possible use of SAR (Synthetic Aperture Radar) for characterizing pasture quality as an alternative to sensors relying on the visible part of the electromagnetic spectrum and the resulting problems due to low illumination and cloud cover in the Arctic. (Maynard et al, 2005; Yurchak and Maynard, 2005) This work, part of a larger study called Reindeer Mapper, preceded the EALAT program as a pilot for the EALAT project to study remote sensing technologies for reducing the threats to reindeer husbandry from climate and land changes by creating a source of usable, timely satellite data that could be combined with traditional, local and other data and information for improved decision-making. (Maynard, 2003 a,b; 2004)

Based upon discussions among Reindeer Mapper team members from within the reindeer husbandry community, including discussions and publications such as the Yakutsk Declaration from the Third World Reindeer Herders' Congress in March 2005, a preliminary list of the highest priority environmental measurements for remotely sensed data was generated. These requirements constituted the primary elements determining pasture quality and state, the most important overall set of parameters for reindeer herders. A summary of these data requirements is listed in figure 6.

Fig. 6. High priority remote sensing observations.....

This preliminary study of the use of SAR for characterizing the quality of reindeer pasture was initiated because SAR does not rely on the visible part of the spectrum and, therefore, has the ability to provide data regardless of weather or light conditions. (Yurchak and Maynard, 2005) These early studies focused initially on the highest priority measurements/data products identified by the reindeer herders on the team. (Box 1). The applications of SAR for characterization of vegetation and measuring snow

parameters are not as well-developed as optical sensors. Initial studies of seasonal changes in SAR backscatter from different kinds of land features in two locations (Anadyr River Research Area (ARRA) and Vaegi Village Research Area (VVRA) in Chukotka, Russia, were carried out for the four seasons of the period between the years 2000 and 2004. Site selection was done based on data availability from the Alaska Satellite Facility (ASF) and on the location of typical tundra landscapes on reindeer pasture areas. Based on these criteria, two sites within the Anadyr district of Chukotskiy Autonomous Okrug (ChAO) were selected. The first site is a nature conservation area north of “Krasnoe” lake along the Anadyr river (Anadyr river research area - ARRA); the second site, a fire risk area south of Vaegi Village (Vaegi Village research area -VVRA). (Maynard et al, 2005; Yurchak and Maynard, 2005)

Results from the study show that the SAR data can detect fire scars very well and could be used for fire scar inventory mapping in conjunction with other systems such as the MODIS Rapid Response System and an analysis of tundra lakes’ radar properties suggested the possibility for remote assessment of the depth of lakes (Yurchak and Maynard, 2005). It was also possible to observe the snow masking effect (Ulaby et al., 1994) and wet snow (Bagdadi et al., 1997). Studies showed the capability of SAR to delineate different types of tundra species as well as demonstrate seasonal changes in radar backscatter from tussock and mountain tundra in time series studies. The sensitivity of SAR data to vegetation and snow cover over plains and mountain tundra is demonstrated on time series study of a selected area in the north of ARRA, fig.7 and fig.8.

Figs. 7 & 8 (A classified part of SAR image.....)

The results showed clear seasonal changes in tundra radar backscatter. For tussock tundra the backscatter was higher in summer months and dropped to the lowest value in the fall due to decrease of soil (vegetation) moisture because of freezing. The subsequent backscatter increase in the winter could be related to snow cover impact. For mountain tundra, summer backscatter behavior is opposite to that of tussock: it is the lowest. Also, the range of winter-summer decrease is rather high: ~ 60 DN. The reason for such behavior, probably, is different local incidence angles for tussock tundra (~ 23⁰) and for the mountain slope (~0⁰). Further field validation work was planned for this study. In addition, SAR data were shown to be capable of delineating detailed geobotanic polygons. SAR data were compared with ground-based geobotanic maps and were found to provide a higher resolution set of polygons than aerial surveys. These preliminary results suggested that further development of the methodology as well as its validation and calibration may result in a reliable method for SAR applications to these important environmental parameters.

In summary, the study showed that SAR data can detect fire scars very well and could be used for fire scar inventory mapping in conjunction with other systems such as the MODIS Rapid Response System. Studies showed the capability of SAR to delineate different types of tundra species as well as demonstrate seasonal changes in radar backscatter from tussock and mountain tundra in time series studies. In addition, SAR data were shown to be capable of delineating detailed geobotanic polygons. SAR data were compared with ground-based geobotanic maps and were found to provide an even higher resolution set of polygons than aerial surveys. An analysis of tundra lakes' radar properties suggests that SAR may provide a useful means of remotely assessing the state of lakes. As temperature increases cause earlier melting of lakes along migration routes in springtime, this technique for lake assessment could prove to be very valuable for herders on a real-time basis.

7.3.2: EALAT On-going Studies

7.3.2.1 Indigenous Linguistics Studies of Reindeer Herding Language

One of the main objectives within the IPY EALÁT project is documenting indigenous knowledge about snow conditions and indigenous perceptions about how they are adapting to changing conditions. A great deal of insight can be gained from centuries-old knowledge within reindeer herding societies such as the Sáami and the Nenets. With the numbers of those from traditional reindeer herding communities following more traditional ways of life on the decline, it is important to document this knowledge as much as possible while it still exists. The study of the language itself is important because it is through language that traditional knowledge becomes available and, particularly, through specific terminology.

Some early results from an EALAT project on the language of reindeer husbandry have recently been summarized by Eira et al (2008) and they demonstrate the importance and the richness of the use of all kinds of information and knowledge to address the dramatic changes occurring in the Arctic today. This EALAT study is focusing on how to empower indigenous reindeer herders with the best information – indigenous knowledge as well as scientific/technical knowledge – for addressing the increasing challenges from climate change and loss of grazing lands. Language is a very important part of this and one of the main objectives of the overall EALAT project is to document indigenous knowledge about reindeer herding with the traditional language of reindeer herders in the key role structuring their knowledge and knowledge-sharing.

In the EALAT linguistics study, one of the key early investigations has been the collection and analysis of the existing linguistic concepts that are used in Saami reindeer herding language. There are, for example, in the dialect of Guovdageaidnu/Kautokeino over 1000 individual terms describing reindeer and, especially their appearance including 50 words describing the shape of the antlers. (Magga, 2005; Eira et al, 2008) This early investigation has been studying techniques for the use of language in communicating traditional knowledge about reindeer herding among reindeer herders as well as to non-indigenous scientific and the broader world communities. Results are being documented and publications prepared at this time. (Eira et al, 2008)

7.3.2.2 Indigenous and Scientific Snow Studies

A joint indigenous and scientific snow project now underway addresses a key EALAT objective to document knowledge about snow conditions and how reindeer herders are adapting to the changing conditions in relation to snow and pasture availability and mobility of the herds. As noted earlier, for indigenous peoples of the North, a clear understanding of real-time snow, precipitation, and ice in their area has been critical to their survival. Current EALAT studies are focusing on Saami snow terminology and herding strategies during the winter to find the best forage for the herd. Section 7.3.2.1 describes a key EALAT indigenous linguistics study of reindeer herding language, which highlights centuries-old knowledge from herder societies. In addition, historical observations are being documented which include oral histories and stories from parents, grandparents and other elders about extreme snow events and other phenomena.

The study described in this section is a joint observational study by EALAT and NASA indigenous and scientific collaborators, which is a data collection project with a focus on climate and development changes (snow and vegetation especially). The study is being led by a reindeer herder and Ph.D. student at the Sami University College and Co-PI of EALAT. (Eira et al, 2008) She is coordinating the collection of a comprehensive set of observations in Northern Norway, by 6 reindeer herders over several years, each of whom provide a detailed set of observations of snow, vegetation, meteorological conditions, herd behavior, and other data as they traverse their seasonal migration routes throughout complete seasons. The study area is located across the Saami migration pastures of Northern Norway in EALAT Study area #1.

Fig. 9. Location of EALAT-NASA project site....

This project includes continued data collection with the EALAT-NASA thermochrons which was initiated in 2007. (See Section 7.3.2.3) Each

reindeer herder is recording on special data log sheets the GPS location/time/date of their daily observations data on eleven weather parameters (e.g., wind, cloud cover, precipitation, temperatures), Saami snow terms, physical measurements of snow depth, type, and description, herd behavior, snow conditions as they pertain to the ability of the reindeer to reach the lichens beneath the snow, and thermochron location data.

The herder observations will also be supported by remote sensing and meteorological data from the same time period wherever possible. Remote sensing data to be combined with the herder archives are presently being inventoried for land cover/land use change assessment information including Landsat, MODIS, AMSR-E, and high resolution commercial satellite imagery. A series of cloud-free Landsat scenes between 1972 and 2007 over the field areas of interest have been obtained and are presently being processed. These will be combined with Eurasian GIS data layers, meteorological data, *in situ* field data, and the wealth of indigenous knowledge described in other sections of this chapter. Analyses of these data are under way and, pending research funding, results should be available in 2009-2011.

7.3.2.3 Indigenous and Scientific Studies of Pasture Icing or “Lock-Out”

The increasing temperature variations which are accompanying climate change in the Arctic are causing more freeze-thaw-freeze cycles, resulting in icing of the lichens or ice layers within and on the snow pack in reindeer pastures and pasture “lock-out”. EALAT Project members are carrying out field research to obtain indigenous and scientific data that can improve the capability of reindeer herders to predict and adapt to these adverse weather conditions and climate changes, especially with climate change making weather conditions more variable and less predictable than before.

For example, the traditional practice of the Sáami reindeer herders in Northern Scandinavia was to allow their reindeer to graze on the tundra in coastal areas during the summer months, where they can graze on abundant grasses, bushes, mushrooms and daffodils. However, these traditional practices have been disrupted somewhat due to the presence of modern national borders, development, and climate changes. After the annual slaughter and the first snowfall, the herders bring their reindeer to overwintering pastures in the mountains and tundra in the interior part of upper Scandinavia, where reindeer dig through the snow to get to lichens, the primary staple of the winter diet of reindeer. However certain meteorological conditions can sometimes create conditions that “lock out” winter grazing pastures. If a warm period that partially melts the snow is followed by

rain and then the temperature drops below freezing, this can create a thick coating of ice in the winter pastures that makes it impossible for reindeer to access their primary food source. This can lead to illness and starvation for the reindeer, which translates into serious losses for herders.

Accordingly, the IPY EALÁT project is developing a new adaptive strategy for “lock out” prediction to avoid this increasingly difficult problem. Researchers are using indigenous knowledge together with scientific data to better predict when and where adverse winter grazing conditions might occur so that eventually a service could be set up that would help herders know where winter pastures with bad grazing conditions are so they can avoid them. Meteorologists from the Norwegian Meteorological Institute in Oslo are providing data for models that try to predict snow conditions in Finnmark by looking at temperature gradients throughout the snow pack.

Fig. 10. Photo of NASA thermochrons in snowpack over pasture

These models are being combined with real-time field observations by herders to verify the predictions the models make, including, the use of some NASA technologies. Starting in October 2007, a team of researchers from the Sámi University College placed NASA thermochrons (small devices that record daily temperature readings at regular intervals) along several reindeer migration routes at various depths between the ground and the top of the snow pack. In May 2008, the team removed the first set of thermochrons and temperature data is being analyzed and compared with Norwegian Meteorological Institute (NMI) model predictions. Observations are being compared with remote sensing data from NASA and the European Space Agency (ESA) and data shared with collaborators in the NASA Global Snowflake Network (GSN) and History of Winter (HOW). <http://education.gsfc.nasa.gov/how/> This process is being repeated for four successive winters in order to get an adequate data set to compare with the model.

Fig. 11. Temperatures (°C) measured by NASA thermochrons...

Figure 11 shows the temperatures (°C) measured by NASA thermochrons in one of the reindeer observations stations. The temperatures were measured on the ground (at the bottom of the snowpack) and in the air (at 1.5 meters above ground). The thermochron data show rapid changes in temperature during this period. According to daily observations recorded by the reindeer herders, the snow conditions became worse and *cearga* conditions were formed from April 20th. (*Cearga* in Saami means snow-drift which is so hard that it bears the weight of the reindeer, i.e., a very hard snowpack) Prior to April 15th, cold temperatures and strong winds developed conditions conducive to formation of the *cearga*. On April 18th, frequent temperatures above 0° C created even worse grazing conditions

during this period. This snow condition could be characterized as a kind of “lock-out” condition because it prevents the reindeers’ access through the snow to the vegetation.

New data sets will be produced for snow type and distribution as a result of the thermochron study, which include NASA remote sensing data such as MODIS and AMSR-E as well as in situ data from the Norwegian Meteorological Institute and NOAA/NCEP and integrated into indigenous knowledge and observations for that area.

The Polar View Consortium, which runs a variety of earth-observation services for its end-users all over the world, will also be contributing to the study by providing researchers with snow maps created using satellite data. These maps help in giving an overview of the amount of snow cover and snow cover type in the regions being studied in the EALÁT project. These observations will provide useful information on the snow temperature from the ground to the surface of the snow pack because the influence of the temperature of the ground on the snow above is not fully-understood. Researchers are also examining historical meteorological data taken over the past several decades as well as satellite data, concentrating in particular in years when many pastures were “locked out” due to ice cover.

The combined set of observations will help better understand what happens to the snow above if warming occurs on the ground as well as the questions: what will happen to the snow above it? Will it get wetter? Will it get drier? These answers will help the understanding of how the energy exchange occurs between the ground and the snow above.

Data from the snow study will be integrated by Saami reindeer herders into the system which will become part of a special service to be produced by the combination of all observations and a model to predict whether certain pastures will be locked out due to ice in the snow pack, so that herders can avoid these areas. These will constitute a unique dataset for the land-use and land-cover studies for this part of Eurasia and the first “adaptation early warning system” created by and for reindeer herders. All data such as these will be distributed through the International Centre for Reindeer Husbandry as part of this special service to reindeer herders.

7.4 EALAT Monitoring and Information Integration System – Adaptation and Planning for the Future

To achieve optimal adaptation strategies for reindeer husbandry, it is imperative that governments, local reindeer herders, management, policy,

and decision-makers include reindeer herders and their traditional local and scientific knowledge in future decision-making which impacts the herding community. To enable the efficient, timely collection and integration of their data into these decision- and policy-making processes, EALAT Project reindeer herders are developing their own system to monitor changes based upon traditional knowledge and modern technologies. This new system is based on the principles of the UN Convention of Biological Diversity Art 8, UN Agenda 21 Declaration Ch 26, ILO-169 Convention on Rights of Indigenous Peoples, UN Declaration concerning the rights of Indigenous Peoples 2007, UNESCO's Convention on Protection and Development of Cultural Diversity, and the Yakutsk Declaration from Third World Reindeer Herders Congress in 2005. In fact, the Yakutsk Declaration explicitly stated that reindeer herders should develop their own system to monitor changes of the Arctic natural resources, based on traditional knowledge and modern technology.

EALAT/Monitoring is developing an observation program or monitoring system for reindeer pastoralism in place-based studies in the Saami area (Norway, Sweden, Finland and NW-Russia) and Yamal-Nenets AO, and, later possibly, Nenets AO, the Republic of Sakha (Yakutia), and Chukotka AO. The EALAT/Reindeer Mapper System - being developed at the International Center for Reindeer Husbandry - is a data integration and sharing system to integrate traditional indigenous knowledge together with physical, scientific, and technical data into a common GIS database for improved decision-making and herd management. The system is a follow-on to an early EALAT pilot project of the NASA Land Cover Land Use Change called Reindeer Mapper, which developed a preliminary version of this system designed to bring remote sensing, ground measurements, and information technologies together with indigenous traditional and local knowledge for herder use in management of Northern Eurasian reindeer herds. (Maynard, 2003a,b,c; 2004; Maynard et al, 2005)

Fig. 12. Reindeer Mapper system concept...

This EALAT georeferenced data sharing system will use secure Intranet connections for data collection, management, transmission, analysis, access, and dissemination. The system will function as a portal to link data from a variety of sources and provide that information to multiple herders. Observations and information are being integrated into a central GIS data base so that data from all sources such as NASA products, reindeer herder knowledge, observations and maps, ground-based measurements and observations, herd movements can all be inputted, managed, transmitted, accessed and disseminated in real time for herd management. The EALAT/Reindeer Mapper Information System will assist in the ongoing analysis of trends and detection of emerging events and conditions, which

affect humans, agriculture, and the environment to enhance early warning and management of responses and adaptation.

It is intended that EALAT will be able to provide reindeer herders with an efficient tool for managing the real-time movements and migrations of their herds through enabling improved efficiency in linking different members of the herder settlements or communities and providing real-time local, satellite or other data (e.g., ice melt in lakes and rivers, weather events), thus enabling real time adjustments to herd movements to avoid problems such as changing weather/climate conditions, freeze-thaw “lock-out” problems, or take advantage of availability of better pasturelands along migration routes. The system is being designed to incorporate local data to allow users to bring their own data into the system for analysis in addition to the data provided by the system itself. With the local information of the population, up to date environmental data and habitat characteristics, the system could generate maps depicting important features of interest for reindeer managers.

One of the products derived from the planned system will be a web-based graphic display that allows analysts to quickly pinpoint areas of interest such as those with large concentrations of reindeer and provide surrounding environmental information. The system could be automatically updated with near-real-time information such as hourly precipitation and snowfall rate and accumulation, daily surface and air temperatures, and vegetation cover conditions. The system could bring attention to the proximity of human and animal populations as part of the need for control response. A local GIS will bring these many layers together with several supporting models, showing only a straightforward graphic of the real-time situation in the field. Because the system proposed will be operating in the Internet environment, it should be virtually accessible from any network computers and wireless remote access from the field. The International Center for Reindeer Husbandry in Kautokeino, Norway, is providing regional and international coordination of and access to data sets and expertise, and will act as overall clearinghouse for EALAT information.

7.4 Reindeer Pastoralism and the Future: UArctic International Centre for Reindeer Husbandry

For a sustainable future, reindeer herders themselves are having to define and anticipate risks related to rapid change in their local communities and plan for optimal adaptation strategies. Reindeer herders in Eurasia from the Bering Strait in the east to the Atlantic Ocean in the west will face many challenges related to changes in their grazing lands and their societies due to climate variability and change and Arctic industrial devel-

opment. Reindeer herders therefore are having to prepare themselves, their societies, and management authorities to reduce their vulnerability to change, including, empowering themselves with new technologies to monitor their local communities based on the best knowledge available.

The most recent development in direct response to this need is the initiative from the Association of the World Reindeer Husbandry (WRH) and the International Centre for Reindeer Husbandry and Saami University College in association with the Arctic Center, University of Lapland, Finland to the establishment of the UArctic Institute for Reindeer Husbandry (“UArctic EALAT”) to ensure EALAT has a legacy after the IPY years. As stated by the International Centre for Reindeer Husbandry:

“The new UArctic Institute will be hosted in Kautokeino, Norway, as a pilot institute for research, outreach and education within the objectives of the UArctic Strategic Plan 2008-2013, in accordance with the Yakutsk Declaration (2005) from the Third Congress of world Reindeer Herders and in agreement with the Fairbanks Declaration from the Eighth conference of Parliamentarians of the Arctic Region, Fairbanks, the United States of America, 12-14 August 2008.

“The Fairbanks Declaration states:

“Further build capacity in Arctic communities to adapt to climate change, including the development of new education programmes and skills training initiatives, to allow. Encourage the University of the Arctic to build practical capacity in the north to address the challenges of adaptation to climate change, and to solve the Arctic’s needs for energy, from technical, cultural, economic as well as environmental perspectives, and to provide further education of health care personnel with special focus on Arctic conditions”.

“The UArctic EALAT Institute will provide a unique opportunity in building competence locally in reindeer herders society not presently available in the UArctic. The EALAT network is today responsible for the UArctic thematic network: Adaptation to globalization of the Arctic. The EALAT network is based on the unique cooperative network established by the Association of World Reindeer Herders (WRH) through the Eurasian North.

The Institute is a very significant vehicle for building the capacity of Arctic countries and, in particular, their indigenous peoples, to adapt to climate change, industrial development, and globalization across the Arctic as well as to reduce their vulnerabilities through empowerment with the best indigenous, scientific and technological knowledge available – including, remote sensing.

References

- Abrjutina L.I., 2003. State policy and health of small nationalities of the north of Russia.
- ACIA, 2004. Impacts of a Warming Climate: Arctic Climate Impact Assessment. Cambridge
- Anisimov, O. and B. Fitzharris, "Polar Regions (Arctic and Antarctic)," p.p. 801-841 In: McCarthy, J. et al. Climate Change 2001: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Third Assessment Report of the IPCC, Cambridge University Press, 2001.
- D. Arseneault, N. Villeneuve, C. Boismenu, Y. Leblanc, and J. Deshayé, "Estimating lichen biomass and caribou grazing on the wintering grounds of northern Quebec: an application of fire history and Landsat data," *Journal of Applied Ecology*, vol. 34, p.p. 65-78, 1997.
- A. Auclair, "The role of fire in lichen-dominated tundra and forest-tundra," p.p. 235-256 in R.W. Wein and D.A. MacLean, eds. *The role of fire in northern circumpolar ecosystems*. John Wiley and Sons, Ltd., Chichester, 1983.
- N. Bagdadi, Y. Gauthier, and M. Bernier, "Capability of Multitemporal ERS-1 SAR Data for Wet-Snow Mapping," *Remote Sens. Environ.*, vol. 60, p.p. 174-186, 1997.
- G.I. Belchansky, G.K. Ovchinnikov, and D.C. Douglas, "Comparative evaluation of ALMAZ, ERS-1, JERS-1 and Landsat-TM for discriminating wet tundra," *Geoscience and Remote Sensing Symposium*, 1995. IGARSS '95. "Quantitative Remote Sensing for Science and Applications", Firenze, Italy, 07/10/1995 -07/14/1995, vol.1, p.p. 309-311, 1995.
- J. Brotton, and G. Wall, "Climate change and the Bathurst Caribou Herd in the Northwest Territories, Canada," *Climatic Change*, vol. 35, p.p. 35-52, 1997.

- A. Colpaert, J. Kumpula and M. Nieminen, "Reindeer Pasture Biomass Assessment Using Satellite Remote Sensing," *Arctic*, vol. 56, No. 2, p.p. 147-158, 2003
- D.J. Craighead and J.J. Craighead, "Tracking Caribou Using Satellite Telemetry," *National Geographic Research*, vol. 3(4), p.p. 462-479, 1987.
- P.T. Doran, C.P. McKay, W.P. Adams, M.C. English, R.A. Wharton, and M.A. Meyer, "Climate forcing and thermal feedback of residual lake-ice covers in the high Arctic," *Limnology and Oceanography*, vol. 41, p.p. 839-848, 1996.
- C.R. Duguay, W.R. Rouse, P.M. Lafleur, L.D. Boudreau, Y. Crevier, and T.J. Pultz, "Analysis of Multi-Temporal ERS-1 SAR Data of Subarctic Tundra and Forest in the Northern Hudson Bay Lowland and Implications for Climate Studies," *Canadian Journal of Remote Sensing*, vol. 25, No. 1, p.p. 21-33, 1999.
- Eira, A.J. 2001. "Reindeer Husbandry in Norway". 2nd World Reindeer Congress. P. 42
- Forbes, B., 2000. Reindeer herding and petroleum development on Poluostrov Yamal: Sustainable development or mutually incompatible uses. The Arctic Is: A web resource on human-environment relationships in the Arctic, <http://www.thearctic.is/>.
- **Glazovsky, N.F.;, D.S. Ojima, and N.G. Maynard. (2004?) "Land Use Interactions: Societal-Ecosystem Linkages. Chapter 3.4 of NEESPI Science Plan.....
- T.H. George, W.J. Stringer, and J.N. Baldrige, "Reindeer range in western Alaska from computer-aided digital classification of Landsat data," *Proceedings of the Eleventh International Symposium on Remote Sensing of Environment*. Vol. I., 25-29 April, p.p. 671-682 1977.
- A. Gunn, and T. Skogland, "Responses of caribou and reindeer to global warming," in Walter C. Oechl, et.al.(eds.), *Global Change and Arctic Terrestrial Ecosystems*, Springer-Verlag, New York, 191 p., 1997.
- D.K. Hall, D.B. Fagre, F. Klasner, G. Linebaugh, and G.E. Liston, "Analysis of ERS-1 synthetic aperture radar data of frozen lakes in northern Montana and implications for climate studies," *Journal of Geophysical Research*, vol. 99, C11, p.p. 22473-22482, 1994.
- J.L. Jernsletten, and K. B. Klovov, "Sustainable Reindeer Husbandry," Arctic Council. University of Tromso. 157 p., 2002.
- Johansen B., Karlsen S.R., 2002. Finnmarksvidda – changes in lichen cover 1987-2000. 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.
- C.O. Justice, and S. Korontzi, "A review of satellite fire monitoring and the requirements for global environmental change research," In: Ahern F., Goldammer, G., Justice C.O. 2 (Eds). *Global and Regional Vegeta-*

- tion Fire Monitoring From Space: Planning a Coordinated International Effort, SPB Academic Publishing, The Hague, The Netherlands, p.p. 1-18, 2001.
- C.O. Justice, J.R.G. Townshend, E.F. Vermote, E. Masuoka, R.E. Wolfe, N. El Saleous, D.P. Roy, and J.T. Morisette, "An overview of MODIS Land data processing and product status," *Remote Sensing of Environment*, vol. 83, No. 1-2, p.p. 3-15, 2002.
- R.E.J. Kelly, A.T.C. Chang, J.L. Foster, and D.K. Hall, "Using remote sensing and spatial models to monitor snow depth and snow water equivalent," in R.E.J. Kelly, N.A. Drake and S. Barr (eds.) *Spatial Modelling of the Terrestrial Environment*, Chichester: John Wiley and Sons Ltd., 2004.
- Kitti, H., and Kumpula, T., 2002. "Classification of reindeer pastures: Mapping based on Traditional Ecological Knowledge (TEK) and Remote sensing based mapping," 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.
- Klein, 2005. ACIA reference
- Klokov, K. 2000. Nenets reindeer herders on the lower Yenisei River: traditional economy under current conditions and responses to economic change. *Polar Research* 19 (1), 39-47.
- Kumpula, T., Manderscheid, A., and Colpaert, A., 2002. Evidence of different pasture use from satellite images: Cases from Lapland and the Tibetan plateau. 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.
- S. Li, R. Guritz, T. Logan, M. Shindle, J. Groves, C. Olmsted, F. Carsey, and J. Macmahon, "Summer environmental mapping potential of a large-scale ERS-1 SAR mosaic of the state of Alaska," *Int. J. Remote Sensing*, vol. 20, No. 2, p.p. 387- 401, 1999.
- S.C. Liew, L.K. Kwoh, O.K. Lim, and H. Lim, "Remote sensing of fire and haze," in "Forest fires and regional haze in Southeast Asia", ed. P. Eaton and M. Radojevic. New York: Nova Science Publishers, Chapter 5, p.p. 67-89, 2001.
- Maynard, N.G., B.S. Yurchak, J.M. Turi, S.D. Mathiesen, and R.L. Aissi-Wespi. (2004). "Integrating Indigenous Traditional, Local and Scientific Knowledge for Improved Management, Policy and Decision-Making in Reindeer Husbandry in the Russian Arctic". IASSA 5th International Congress of Arctic Social Sciences (ICASS V). May 19-23, 2004. Fairbanks, AK.
- Maynard, N.G., B.S. Yurchak, and J.M. Turi. (2003). "Shared Knowledge for Decision-Making on Environment and Health Issues in the Arctic". SEARCH (A Study of Environmental Arctic Change) Open Science Meeting. October 27, 2003. Seattle, Washington.

Maynard, N.G., B.S. Yurchak, and J.M. Turi. (2003). "Shared Knowledge for Decision-Making on Environment and Health Issues in the Arctic". HARC: Human/Environment Interactions Research: Patterns, Connections, and Methods Workshop. October 25-26, 2003. Seattle, Washington.

Maynard, N.G., and B.S. Yurchak. (2003). Integrated Scientific, Local and Indigenous Knowledge for Management and Decision-Making on Environment, Climate and health Issues. 2003 Open Meeting Human Dimensions of Global Environmental Change. Montreal, October 16-18, 2003.

Mironenko, O., 2000. "The ways of optimizing of land use and organization systems in reindeer trade-husbandry economies," *Novosti Olenovodstva (Reindeer Husbandry News, in Russian)*

Nieminen, M., Kumpula, J., and Colpaert, A. 2002. "Winter pasture resources of wild forest reindeer (*Rangifer tarandus fennicus*) in Salama-jarvi area in central Finland," 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.

Nuttal et al 2005 - ACIA

Nuttal, M., P.A. Forest, and S.D. Mathiesen. 2008. Background Paper on Adaptation to Climate change in the Arctic. In: Snellmann, O., L. Kullerud, G. Lindstrom, and B.W. Ropstad. Proceedings of the Joint Seminar UArctic Rectors Forum and the Standing committee of Parliamentarians of the Arctic Regions. UArctic publications Series No. 2. Pp. 17-21. ISBN 978-952-484-232-7.

Oskal, A. "Old Livelihoods in New Weather. Arctic Indigenous Reindeer Herders Face the Challenges of Climate Change". Development Outreach. World Bank Institute Special Paper. p.p. 22-25. 2008.

W.G. Rees, M. Williams, and P. Vitebsky, "Mapping land cover change in a reindeer herding area of the Russian Arctic using Landsat TM and ETM+ imagery and indigenous knowledge," *Remote Sensing of Environment*, vol. 85, p.p. 441-452, 2003.

P. Sandstrom, T. Pahlen, L. Edenius, H. Tommervik, O. Hagner, L. Remberg, H. Bisson, K. Baer, T. Stenlund, L. Brandt, and M. Egberth, "Conflict Resolution by Participatory Management: Remote Sensing and GIS as Tools for Communicating Land-use Needs for Reindeer Herding in Northern Sweden," *Ambio*, vol. 32, No. 8, p.p. 557 – 567, 2003.

A.B. Tait, D.K. Hall, J.L. Foster, and R.L. Armstrong, "Utilizing Multiple Datasets for Snow-Cover Mapping," *Remote Sens. Environ.*, vol. 72, p.p. 111-126, 2000.

Tamstorf, M.K., and Aastrup, P., 2002. "Vegetation mapping of West-greenland caribou ranges," 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.

- F.T. Ulaby, E.H. Stiles, and M.A. Abdelrazik, "Snowcover influence on backscattering from terrain," *IEEE Transactions on Geoscience and Remote Sensing*, vol. GE-22, p.p. 126-133, 1984.
- UNEP, 1999. Levine, J.S., Bobbe, T., Ray, N., Singh, A. and R.G. Witt. "Wildland Fires and the Environment a Global Synthesis," UNEP/DEIAEW/TR.99-1.
- Vistnes, I., Nellemann, C., Jordhøy, P., and Strand, O., 2002. "Infrastructure as barriers to wild reindeer migration," 12-th Nordic Conference on Reindeer Research in Kiruna, Sweden, 11-13 March 2002.
- N.E. Walsh, S.G. Fancy, T.R. McCabe, and L.F. Pank, "Habitat use by the Porcupine caribou herd during predicted insect harassment," *Journal of wildlife management*, vol. 56 (3), p.p. 465-473, 1992.
- C. Wang, J. Qi, S. Moran, and R. Marsett, "Soil moisture estimation in a semiarid rangeland using ERS-2 and TM imagery," *Remote Sensing of Environment*, vol. 90, p.p. 178-189, 2004.
- B.S. Yurchak and N.G. Maynard, "Time-Series SAR Observations of Chukotka Sub-Arctic Lakes and Forest-Tundra Fire Scars," *Proceedings of the 31st International Symposium on Remote Sensing of Environment. Global Monitoring for Sustainability and Security. ("ISRSE 2005")*. Saint Petersburg, June 20-24, 2005.