



Wind Observations of Wave Heating and/or Particle Energization at Supercritical Interplanetary Shocks

Lynn Bruce Wilson III (1), Adam Szabo (1), Andriy Koval (5,1), Cynthia A. Cattell (2), Paul J. Kellogg (2), Keith Goetz (2), Aaron Breneman (2), Kris Kersten (2), Justin C. Kasper (3), and Marc Pulupa (4)

(1) Goddard Space Flight Center, Heliospheric Physics Laboratory, Greenbelt, United States, (2) Department of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota, United States, (3) Harvard-Smithsonian Center for Astrophysics, Harvard University, Cambridge, Massachusetts, United States, (4) Space Sciences Lab, University of California at Berkeley, Berkeley, California, United States, (5) Goddard Earth Sciences and Technology Center, University of Maryland Baltimore County, Baltimore, Maryland, United States

We present the first observations at supercritical interplanetary shocks of large amplitude (> 100 mV/m pk-pk) solitary waves, ~ 30 mV/m pk-pk waves exhibiting characteristics consistent with electron Bernstein waves, and > 20 nT pk-pk electromagnetic lower hybrid-like waves, with simultaneous evidence for wave heating and particle energization. The solitary waves and the Bernstein-like waves were likely due to instabilities driven by the free energy provided by reflected ions [Wilson III *et al.*, 2010]. They were associated with strong particle heating in both the electrons and ions. We also show a case example of parallel electron energization and perpendicular ion heating due to a electromagnetic lower hybrid-like wave. Both studies provide the first experimental evidence of wave heating and/or particle energization at interplanetary shocks. Our experimental results, together with the results of recent Vlasov [Petkaki and Freeman, 2008] and PIC [Matsukiyo and Scholer, 2006] simulations using realistic mass ratios provide new evidence to suggest that the importance of wave-particle dissipation at shocks may be greater than previously thought.