

CASW

Ronald Cohen

Tuesday, August 30, 2011

31 - Physical properties and seasonal behavior of H₂O, HDO, CO₂ and trace gases on Mars: Quantitative mapping from Earth-based observatories

Robert E Novak¹, rnovak@iona.edu, Michael J Mumma², Geronimo L Villanueva². (1) Department of Physics, Iona College, New Rochelle, New York 10801-1830, United States, (2) Solar System Exploration Division, NASA Goddard Space Flight Center, Greenbelt, Maryland 20771, United States

Since 1997, we have used high-resolution ($R > 40000$) spectrometers on ground based-telescopes to study molecules that have astrobiological significance in Mars' atmosphere. We have used the NASA-IRTF, Keck II, and VLT telescopes in the 1.0-5.0 micron range. The spectrometer is set at a wavelength to detect specific molecules. Spectral/spatial images are produced. Extracts from these images provide column densities centered at latitude/longitude locations (resolution $\sim 400\text{km}$ at sub-Earth point). We have mapped the O₂ singlet-Delta emission (a proxy for ozone), HDO, and H₂O for seasonal dates throughout the Martian year. Previously undiscovered isotopic bands of CO₂ have been identified along with isotopic forms of CO. We are searching for other molecules that have astrobiological importance and have successfully measured methane in Mars' atmosphere.

We acknowledge support from NSF (AST-0805540) and NASA (Planetary Astronomy (344-32-51-96) and Astrobiology (344-53-51)) and thank the Director and Staff of the telescopes mentioned for observing time.

~~Tuesday, August 30, 2011 10:50 AM~~

~~[Chemistry as a Tool for Space Exploration and Discovery at Mars \(08:30 AM - 11:50 AM\)](#)~~

~~Location: Colorado Convention Center~~

~~Room: 102~~

~~[Close Window](#)~~