

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

Planetary and Space Science

journal homepage: www.elsevier.com/locate/pss

Jovian plasma torus interaction with Europa. Plasma wake structure and effect of inductive magnetic field: 3D hybrid kinetic simulation

A.S. Lipatov^{a,c,*}, J.F. Cooper^b, W.R. Paterson^b, E.C. Sittler Jr.^b, R.E. Hartle^b, D.G. Simpson^b

^a Goddard Planetary Heliophysics Institute, UMBC/NASA GSFC, MD 20771, USA

^b NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

^c Department of Problems of Physics and Power Engineering, Moscow Institute of Physics and Technology, Russia

ARTICLE INFO

Article history:

Received 27 November 2011

Received in revised form

11 October 2012

Accepted 8 January 2013

Available online 4 February 2013

Keywords:

Europa

Jovian magnetosphere

Plasma

Magnetic fields

Ion composition

ABSTRACT

The hybrid kinetic model supports comprehensive simulation of the interaction between different spatial and energetic elements of the Europa moon–magnetosphere system with respect to a variable upstream magnetic field and flux or density distributions of plasma and energetic ions, electrons, and neutral atoms. This capability is critical for improving the interpretation of the existing Europa flyby measurements from the Galileo Orbiter mission, and for planning flyby and orbital measurements (including the surface and atmospheric compositions) for future missions. The simulations are based on recent models of the atmosphere of Europa (Cassidy et al., 2007; Shematovich et al., 2005). In contrast to previous approaches with MHD simulations, the hybrid model allows us to fully take into account the finite gyroradius effect and electron pressure, and to correctly estimate the ion velocity distribution and the fluxes along the magnetic field (assuming an initial Maxwellian velocity distribution for upstream background ions). Photoionization, electron-impact ionization, charge exchange and collisions between the ions and neutrals are also included in our model. We consider the models with O^{++} and S^{++} background plasma, and various betas for background ions and electrons, and pickup electrons. The majority of O_2 atmosphere is thermal with an extended non-thermal population (Cassidy et al., 2007). In this paper, we discuss two tasks: (1) the plasma wake structure dependence on the parameters of the upstream plasma and Europa's atmosphere (model I, cases (a) and (b) with a homogeneous Jovian magnetosphere field, an inductive magnetic dipole and high oceanic shell conductivity); and (2) estimation of the possible effect of an induced magnetic field arising from oceanic shell conductivity. This effect was estimated based on the difference between the observed and modeled magnetic fields (model II, case (c) with an inhomogeneous Jovian magnetosphere field, an inductive magnetic dipole and low oceanic shell conductivity).

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

The interaction of the Jovian plasma torus with Europa and other moons is a fundamental problem in magnetospheric physics (see e.g., Goertz, 1980; Southwood et al., 1980; Southwood and Dunlop, 1984; Wolf-Gladrow et al., 1987; Ip, 1990; Schreier et al., 1993; Lellouch, 1996). The plasma environment near Europa was studied by flyby observations during the Galileo prime mission

and the extended Galileo Europa mission (Kivelson et al., 1997; Khurana et al., 1998; Kivelson et al., 1999; Paterson et al., 1999).

Europa, one of the icy moons of Jupiter, was encountered by the Galileo satellite three times during its primary mission, seven times during its Galileo Europa Mission (GEM), and once during Galileo Millennium Mission (GMM). Europa is located at a radial distance of $9.4R_J$ (Jovian radii, 71,492 km) from Jupiter and has a radius of 1560 km ($1R_E$).

The interaction of Europa with the magnetized plasma of the Jovian plasma sheet gives rise to a so-called Alfvén wing, which has been extensively studied in the case of Io (e.g., Neubauer, 1980; Southwood et al., 1980; Herbert, 1985; Lipatov and Combi, 2006). Neubauer (1998, 1999) has shown theoretically how an Alfvén wing is modified by an induced magnetic field, such as that found at Europa (Kivelson et al., 2000). Observations by Kivelson et al. (1992) show the generation of ultra-low frequency electromagnetic waves in Europa's wake. These waves have frequencies near and below the gyrofrequencies of the ion species in the

* Corresponding author. NASA Goddard Space Flight Center, Greenbelt, Code 673, Bld. 21, Rm. 247, 8800 Greenbelt Rd., Greenbelt, MD 20771, USA.

Tel.: +1 301 286 0906; fax: +1 301 286 1648.

E-mail addresses: Alexander.Lipatov-1@nasa.gov, alipatov@umbc.edu (A.S. Lipatov), John.F.Cooper@nasa.gov (J.F. Cooper), William.R.Paterson@nasa.gov (W.R. Paterson), Edward.C.Sittler@nasa.gov (E.C. Sittler Jr.), Richard.E.Hartle@nasa.gov (R.E. Hartle), David.G.Simpson@nasa.gov (D.G. Simpson).

plasma torus (e.g., ionized sulfur, oxygen, and protons). Ion cyclotron waves grow when ion distribution functions are sufficiently anisotropic, as occurs when ion pickup creates a ring distribution of ions (in velocity space). The analysis of these waves has been done by Huddleston et al. (1997) (Io), Volwerk et al. (2001) and Kivelson et al. (2009) (Europa). They found intensive wave power at low frequencies (near and below the cyclotron frequencies of heavy ions) in Europa's wake during the E11 and E15 flybys. However, our current 3D hybrid modeling cannot yet produce these waves due to insufficient spatial grid resolution.

The most general and accurate theoretical approach to this problem would require the solution of a nonlinear coupled set of integro-MHD/kinetic-Boltzmann equations which describe the dynamics of Jupiter's corotating magnetospheric plasma, pickup ions, and ionosphere, together with the neutrals from Europa's atmosphere. To first order, the plasma and neutral atoms and molecules are coupled by charge exchange and ionization. The characteristic scale of the ionized components is usually determined by the typical ion gyroradius, which for Europa is much less than characteristic global magnetospheric scales of interest, but which may be comparable to the thickness of the plasma structures near Europa. Kinetic approaches, such as Direct Simulation Monte Carlo, have been applied to the understanding of global aspects of the neutral atmosphere (Marconi et al., 1996; Austin and Goldstein, 2000). Plasma kinetic modeling is, however, much more complicated, and even at the current stage of computational technology requires some approximations and compromises to make some initial progress. Several approaches have been formulated for including the neutral component and pickup ions self-consistently in models that describe the interaction of the plasma torus with Europa.

There have been recent efforts to improve and extend the pre-Galileo models for Europa, Io and Ganymede, in terms of the MHD (Kabin et al., 1999; Combi et al., 1998; Linker et al., 1998; Kabin et al., 2001; Jia et al., 2008), the electrodynamic (Saur et al., 1998, 1999; Schilling et al., 2008), and hybrid kinetic (Lipatov and Combi, 2006; Lipatov et al., 2010) approaches. These approaches are distinguished by the physical assumptions that they include. MHD and hybrid kinetic models cannot, at least yet, include the charge separation effects which are likely to be important very close to the moon where the neutral densities are large and the electric potential can introduce non-symmetric flow around the body. MHD models for Io either include constant artificial conductivity (Linker et al., 1998) or assume perfect conductivity (Combi et al., 1998). Comparisons of the sets of published results do not indicate that this choice has any important consequences. The MHD model of Europa developed by Kabin et al. (1999) includes an exospheric mass loading, ion-neutral charge exchange, and recombination. Further development of this model by Liu et al. (2000) already includes a possible intrinsic dipole magnetic field of Europa. Schilling et al. (2007, 2008) found that for the conductivity of Europa's ocean values of 500 mS/m or large combined with ocean thickness of 100 km and smaller to be most suitable to explain the magnetic flyby data. They also found that the influence of the fields induced by the time variable plasma interaction is small compared to the induction caused by the time-varying background field.

Hybrid kinetic models can include the finite ion gyroradius effects, non-Maxwellian velocity distribution for ions, and correct flux of pickup ions along the magnetic field. Hybrid modeling of Io has demonstrated several features. The kinetic behavior of ion dynamics reproduces the inverse structure of the magnetic field (due to drift current) which cannot be explained by standard MHD or electrodynamic modeling which do not account for anisotropic ion pressure. The diamagnetic effect of non-isotropic

gyrating pickup ions broadens the B-field perturbation and produces increased temperatures in the flanks of the wake, as observed by the Galileo spacecraft, but had not been explained by previous models. The temperatures of the electrons which are created and cooled by collisions with neutrals in the exosphere and inside the ionosphere may strongly affect the pickup ion dynamics along the magnetic field and consequently the pickup distribution across the wake. The physical chemistry in Io's corona was considered in the paper by Dols et al. (2008). They couple a model of the plasma flow around Io plus a multi-species chemistry model and compare the model results to the Galileo observation in Io's wake.

Galileo flyby measurements E4, E6 (plasma only), E11, E12, E14, E15, E19, and E26 demonstrate several features in the plasma environment: Alfvén wing formation and an induced magnetosphere, possible existence of the dipole-type induced magnetic field, and variation of the magnetic field in the plasma wake due to diamagnetic currents. The measurements also demonstrate mass loading of the plasma torus plasma by pickup ions and the interaction of the ions with the surface of Europa. For an interpretation of these data, we need to use a kinetic model because of effects of the finite ion gyroradius.

Hybrid models have been shown to be very useful in studying the complex plasma wave processes of space, astrophysical, and laboratory plasmas. These models provide a kinetic description of plasmas in local regions, together with the possibility of performing global modeling of the whole plasma system. Revolutionary advances in computational speed and memory are making hybrid modeling of various space plasma problems a much more effective general tool.

In this paper, we apply a time-dependent Boltzmann equation (a "particle in cell" approach) together with a hybrid kinetic plasma (ion kinetic) model in three spatial dimensions (see, e.g. Lipatov and Combi, 2006; Lipatov et al., 2010), using a prescribed but adjustable neutral atmosphere model for Europa. A Boltzmann simulation is applied to model charge exchange between incoming and pickup ions and the immobile atmospheric neutrals. In this paper, we discuss the results of the hybrid kinetic modeling of Europa's environment—namely the global plasma structures (formation of the magnetic barrier, Alfvén wing, pickup ion tail, etc.). The results of these kinetic modeling are compared with the Galileo E4 flyby observational data. Currently, we are working on the hybrid model of the E12 flyby. The remarkable aspect of this flyby is a strong variation in the upstream plasma density profile approximately from 400 cm^{-3} to 80 cm^{-3} . The results of this modeling will be discussed in future publications.

The paper is organized as follows: in Section 2 we present the computational model and a formulation of the problem. In Section 3 we present the results of the modeling of the plasma environment near Europa and the comparison with observational data. Finally, in Section 4 we summarize our results and discuss the future development of our computational model.

2. Formulation of the problem and mathematical model

To study the interaction of the plasma torus with the ionized and neutral components of Europa's environment, we use a quasineutral hybrid model for ions and electrons. The model includes ionization (which in the Europa environment is dominated by electron impact ionization, not photoionization) and charge exchange. The atmosphere is considered to be an immobile component in this paper.

In our hybrid modeling, the dynamics of upstream ions and implanted ions are described in a kinetic approach, while the dynamics of the electrons are described in a hydrodynamical

approximation. The details of this plasma-neutral approach were developed early for the study of the Io–Jovian plasma interaction (Lipatov and Combi, 2006).

The single ion particle motion is described by the equations (see, e.g. Eqs. (1) and (14) from Mankofsky et al., 1987)

$$\frac{d\mathbf{r}_{s,l}}{dt} = \mathbf{v}_{s,l}, \quad \frac{d\mathbf{v}_{s,l}}{dt} = \frac{e}{M_i} \left(\mathbf{E} + \frac{\mathbf{v}_{s,l} \times \mathbf{B}}{c} \right) - \frac{m_e v_{ie}}{M_i} (\mathbf{v}_{s,l} - \mathbf{U}_i) - \frac{m_e v_{io}}{M_i} \mathbf{J} - v_{io} \mathbf{v}_{s,l}. \quad (1)$$

Here we assume that the charge state is $Z_i = 1$. \mathbf{U}_i , and \mathbf{J} denote the charge-averaged velocity of all (incoming and pickup) ions and the total current, Eq. (5). The subscript s denotes the ion population ($s=1,2$ for incoming ions and $s=3,4$ for pickup ions) and the index l is the particle index. v_{ie} and v_{io} are collision frequencies between ions and electrons, and ions and neutrals that may include Coulomb collisions and collisions due to particle–wave interaction.

For a plasma, the thermal velocity, v'_α ($\alpha = i, e$), is assumed greater than the drift velocity, so we take

$$v_{\alpha,0} = n_\alpha \sigma^{\alpha,z} v'_{\alpha,z}, \quad (2)$$

where the cross section $\sigma^{\alpha,z}$ is typically about $5 \times 10^{-15} \text{ cm}^2$ (see, e.g., Eq. (17) from Mankofsky et al., 1987).

For massless electrons the equation of motion of the electron fluid takes the form of the standard generalized Ohm's law (e.g. Braginskii, 1965)

$$\mathbf{E} = \frac{1}{en_e c} (\mathbf{J}_e \times \mathbf{B}) - \frac{1}{en_e} \nabla p_e - \frac{m_e}{e} \left[\sum_s v_{e,s} \left[(\mathbf{U}_i - \mathbf{U}_s) - \frac{\mathbf{J}}{ne} \right] + v_{a,eo} \mathbf{U}_e \right], \quad (3)$$

where $p_e = nm_e \langle v_e'^2 \rangle / 3 = n_e k_B T_e$, and v'_e are the scalar electron pressure and the thermal velocity of electrons, and the electron current is estimated from Eq. (5).

The induction equation (Faraday's law) has a form

$$\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} + \nabla \times \mathbf{E} = 0. \quad (4)$$

The total current is given by

$$\mathbf{J} = \mathbf{J}_e + \mathbf{J}_i, \quad \mathbf{J}_i = \sum_{s=1}^2 en_s \mathbf{U}_s = en_i \mathbf{U}_i, \quad (5)$$

where \mathbf{U}_s is the bulk velocity of ions of the type s .

Since we suppose that electron heating due to collisions with ions is very small, the electron fluid is considered adiabatic. For simplicity, we assume that the total electron pressure may be represented as a sum of partial pressures of all electron populations

$$p_e \propto \frac{(\beta_e n_{i,up}^{5/3} + \beta_{e,pl} n_{i,pl}^{5/3})}{\beta_e}, \quad (6)$$

where β_e and $\beta_{e,pl}$ denote electron upwind and pickup betas, respectively. Note that $\beta_{e,k} = p_{e,k} / (B^2 / 8\pi)$, where k is a population of electrons. We also assume here that $n_{e,up} = n_{i,up}$, $n_{e,pl} = n_{i,pl}$.

The neutral atmosphere of Europa serves as a source of new ions, mainly by electron impact ionization from corotating (or nearly corotating) plasma and also by photoionization. The neutral atmospheric molecules also serve as collisional targets for charge exchange by corotating ions. The impacting ions consist of both upstream torus ions and newly implanted ions which are picked up by the motional electric field.

In the current model, we assume that the background plasma contains only the ions with molecular mass/charge of 8 and 16 corresponding to O^{++} and S^{++} , respectively.

We assume that Europa has a radius $R_E = 1560 \text{ km}$. We have also adopted a two-species description for the neutral O_2

exosphere of exponential form (Shematovich et al., 2005)

$$n_{\text{neutral},k} \approx n_{\text{atmos},k} \exp[-(r - r_{\text{exobase},k}) / h_{\text{atmos},k}], \quad (7)$$



where $n_{\text{atmos},k}$ denotes the maximum value of the neutral density extrapolated to the exobase ($n_{\text{atmos},1} = 3 \times 10^4 \text{ cm}^{-3}$; $n_{\text{atmos},2} = 8.5 \times 10^7 \text{ cm}^{-3}$; $r_{\text{exobase},1} \approx 1700 \text{ km}$; $r_{\text{exobase},2} \approx 1560 \text{ km}$), and index k denotes either non-thermal ($k=1$) or thermal ($k=2$) species. Here the scale heights $h_{\text{atmos},1} = 200 \text{ km}$ and $h_{\text{atmos},2} = 30 \text{ km}$.

The production rate of new ions from the exosphere near Europa corresponds to

$$G_{\text{exo},k} \propto v_{i,k} n_{\text{atmos},k} \exp[-(r - r_{\text{exobase},k}) / h_{\text{atmos},k}], \quad (8)$$

where $n_{\text{atmos},k}$ denotes the value of the neutral component density at $r = r_{\text{exobase},k}$ and $v_{i,k}$ is the effective ionization rate per atom or molecule of species k . $v_{i,k}$ includes the photoionization v_{ph} , and the electron impact ionization by the magnetospheric electrons $v_{e,im}$. We assume that our model of the atmosphere mainly consists of O_2 , and we use the effective photoionization rate $1.7 \times 10^{-8} \text{ s}^{-1}$ (Johnson et al., 2009). We also adopt the effective electron impact ionization rates of $2.4 \times 10^{-8} \text{ cm}^3/\text{s}$ (for 20 eV electrons) and $1.1 \times 10^{-7} \text{ cm}^3/\text{s}$ (for 250 eV electrons) (see e.g. Johnson et al., 2009). Since the hot electrons represent only 5% of the total electron density (see Voyager 1 plasma science (PLS) measurements analyzed by Sittler and Strobel, 1987; Bagenal, 1994) we use the same composition for computing the impact ionization rate. We assume that the Sun is located in the direction opposite the x -axis.

The interaction of ions with neutral particles by charge exchange (see Eqs. (12)–(15) from Lipatov and Combi, 2006) currently includes for the following reactions:



The effective cross section for charge exchange ($\sigma_{c,ex} = 2.6 \times 10^{-19} \text{ m}^2$) was the same as that used in the hybrid modeling of Io's plasma environment (see Lipatov and Combi, 2006; McGrath and Johnson, 1989). A more complete list of reactions will be considered in future modeling. Of course, this also requires the addition of Monte Carlo computations. However, this approach is beyond the scope of this paper.

Our code solves Eqs. (1)–(9).

We discuss two models of the interaction between the Jovian magnetosphere and Europa. In Section 3.1 we discuss the interaction model for the cases with different ions and electron betas, different pickup ion production rates near the surface of Europa, and homogeneous global Jovian magnetic field (model I, cases (a) and (b)), whereas in Section 3.2 we consider the model II, case (c) with realistic global Jovian magnetic field and the internal dipole magnetic field placed in the center of Europa. To study the interaction of the plasma torus with the ionosphere of Europa, the following set of Jovian plasma torus and ionosphere parameters were adopted in accordance with the Galileo Europa E4 flyby observational data (Paterson et al., 1999; Khurana et al., 1998; Kivelson et al., 1997, 1999): magnetic field, $B_0 = 469 \text{ nT}$ and $\mathbf{B} = (77.6, -140.7, -441.3) \text{ nT}$; torus plasma speed relative to Europa (Paterson et al., 1999), $U_0 = 105 \text{ km/s}$; upstream ion densities, $\rho_{O^{++}} = 10 \text{ cm}^{-3}$; $\rho_{S^{++}} = 10 \text{ cm}^{-3}$ and ion temperature, $T_i = 25 - 100 \text{ eV}$ (Paterson et al., 1999); electron temperature for suprathermal population, $T_e = 20 \text{ eV}$ (Sittler and Strobel, 1987); ratio of specific heats, $\gamma = 5/3$; Alfvén and sonic Mach numbers, $M_A = 0.25$, $M_S = 3.66$.




Fig. 1. Europa's environment and system of coordinates.

Initial conditions: Initially, the computational domain contains only supersonic and sub-Alfvénic plasma torus flow with a homogeneous spatial distribution and a Maxwellian velocity distribution; the pickup ions have a weak density and spherical spatial distribution. The magnetic and electric fields are $\mathbf{B} = \mathbf{B}_0$ and $\mathbf{E} = -\mathbf{U}_0 \times \mathbf{B}_0$. Inside Europa the electromagnetic fields are $\mathbf{E} = 0$ and $\mathbf{B} = \mathbf{B}_0$, and the bulk velocities of ions and electrons are also equal to zero. Here the x -axis is directed in the corotation direction, the y -axis is directed toward Jupiter, and the z -axis is directed to the north, as shown in Fig. 1. In model I, cases (a) and (b) we use a homogeneous magnetic field for the initial and boundary conditions (see paragraph above). In model II, case (c) we use an extrapolation of the magnetic field profile along E4 trajectory (see, Kivelson et al., 1999, 2009) onto the computation domain for the initial and boundary conditions. The effect of global variation in the magnetic field in the rest of Europa was not taken into account directly in the modeling but it was included in to the modeling as an internal magnetic dipole (see, Schilling et al., 2007, 2008).

At $t > 0$ we begin to inject the pickup ions with a spatial distribution according to Eq. (8). Far upstream ($x = -15 R_E$), the background ion flux is assumed to have a Maxwellian velocity distribution.

Boundary conditions: On the side boundaries ($y = \pm DY/2$ and $z = \pm DZ/2$), periodic boundary conditions were imposed for incoming flow particles. The pickup ions exit the computational domain when they intersect the side boundary surfaces $y = DY/2 - 5 \times \Delta y$, $y = -DY/2 + 5 \times \Delta y$, $z = DZ/2 - \times \Delta z$, $z = -DZ/2 + 5 \times \Delta z$. Thus there is no influx of pickup ions at the side boundaries.

At the side boundaries, we also use a damping boundary condition for the electromagnetic field (see e.g., Lipatov and Combi, 2006; Umeda et al., 2001). This procedure allows us to reduce outgoing electromagnetic perturbations, which may be reflected at the boundaries.

Far downstream ($x = 12 R_E$), we adopted a free escape condition for particles and the ‘‘Sommerfeld’’ radiation condition for the magnetic field (see e.g., Tikhonov and Samarskii, 1963) and a free escape condition for particles with re-entry of a portion of the particles from the outflow plasma.

At Europa's surface, $r = R_E \approx 1560$ km, the particles are absorbed. In model I, there is no boundary condition at Europa's surface for the electromagnetic field; we also use our equations for the electromagnetic field, (see, Eqs. (2), (4) and (9) from Lipatov and Combi, 2006) inside Europa but using the low internal conductivity (Reynolds number, $Re = 0.5$) and very small value for bulk velocity that are calculated from the particles. In model II, we also use an inductive magnetic dipole

$(0, 0, -72.5) \text{ nTR}_E^3$ for the boundary condition at Europa's surface that simulates the effect of nonstationarity of Jovian magnetic field at the position of Europa. In this way, the jump in the electric field is due to the variation of the value of the conductivity and bulk velocity across Europa's surface (note that the center of Europa is at $x = 0, y = 0, z = 0$).

The three-dimensional computational domain has dimensions $DX = 27 R_E$, $DY = 30 R_E$ and $DZ = 30 R_E$. We used mesh of $301 \times 301 \times 271$ grid points, and 5×10^8 and 5×10^8 particles for ions and pickup ions, respectively, for a homogeneous mesh computation. The particle time step Δt_p and the electromagnetic field time step Δt_{EB} satisfy the following condition: $v_{max} \Delta t_p \leq \min(\Delta x, \Delta y, \Delta z)/8$ and $v_{max} \Delta t_{EB} \leq \min(\Delta x, \Delta y, \Delta z)/256$.

The global physics in Europa's environment is controlled by a set of dimensionless independent parameters such as M_A , β_i , β_e , M_i/M_p , ion production and charge exchange rates, diffusion lengths, and the ion gyroradius $\epsilon = \rho_{ci}/R_E$. Here $\rho_{ci} = U_0 / (eB/M_i c) = M_{AC}/\omega_{pi}$ and the ion plasma frequency $\omega_{pi} = \sqrt{4\pi n_0 e^2/M_i}$. M_i and M_p denote the ion and proton masses. For real values of the magnetic field, the value of the ion gyroradius is about 80 km, which is calculated from the local bulk velocity. The dimensionless ion gyroradius and grid spacing have the values $\epsilon = 0.05$ and $\Delta x/R_E = 0.1$.

In order to study ion kinetic effects (e.g. excitation of low-frequency oscillations ($\omega \ll \Omega_b$) by the mass loading), we must satisfy the condition $\Delta \leq (10-20)c/\omega_{pb}$, where Ω_b and ω_{pb} denote the gyrofrequency and the plasma frequency for upstream ions (Winske et al., 1985). The above estimation of the plasma parameters shows that we have good resolution for the low-frequency waves (see also Lipatov et al., 2012).

There is another problem—numerical resolution of the gyroradius on the spatial grid. It becomes very important near Europa's surface where the MHD model cannot to be used and we have to use a kinetic model to study a trajectory of the heavy ions and their interaction with the surface of Europa. Our current model still does resolved the last effect and we expect to improved the model by use a spherical system of coordinates in future research.

3. Results of Europa's environment simulation

3.1. Effects of plasma betas on the plasma wake structure

In order to study the effect of plasma parameters on the structure of the plasma wake and the Alfvén wing, we have performed a modeling (model I) for two cases (a) and (b) with different values of the upstream background ion temperatures, pickup electron temperatures, and value of the pickup production rate near the surface of Europa.

The following plasma parameters are chosen the same for both models: full magnetosphere corotation speed is $U_0 = 105$ km/s; upstream densities are $\rho_{O^{++}} = 10 \text{ cm}^{-3}$, $\rho_{S^{++}} = 10 \text{ cm}^{-3}$; magnetic field is $B_0 = 469$ nT; $\mathbf{B} = 77.6, -140.7, -441.3$ nT; Alfvénic Mach number $M_A = 0.25$; magnetosonic Mach number $M_s = 3.66$. The model of O_2 atmosphere was taken from Cassidy et al. (2007), Shematovich et al. (2005) and Smyth and Marconi (2006). In model I, cases (a) and (b), Europa's interior is represented as low conducting body with Reynolds number $Re = 0.5$.

Model I, case (a): Upstream ion temperatures are $T_{O^{++}} = 25$ eV; $T_{S^{++}} = 25$ eV and upstream electron temperature is $T_{e,0} = 20$ eV. Temperatures of electrons connected with non-thermal and thermal O_2^+ pickup ions are $T_{e,non-thermal} = 20$ eV; $T_{e,thermal} = 20$ eV.




Fig. 2. 2D cuts of non-thermal pickup ion O_2^+ density profile. Model I, case (a) (top) and case (b) (bottom). x - y cuts (left column) are located at $z=0$, y - z cuts are located at $x/R_E = 7$, and x - z cuts (right column) are located at $y=0$.




Fig. 3. 2D cuts of the thermal pickup ion O_2^+ density profile. Model I, case (a) (top) and case (b) (bottom). x - y cuts (left column) are located at $z=0$, y - z cuts are located at $x/R_E = 7$, and x - z cuts (right column) are located at $y=0$.

Model I, case (b) (reduced density for thermal O_2 by factor 60 near surface and higher electron temperatures; increased upstream ion temperatures, $T_{O^{++}} = 100$ eV; $T_{S^{++}} = 100$ eV); upstream

electron temperature is $T_{e,0} = 20$ eV; temperatures of electrons connected with non-thermal and thermal O_2^+ pickup ions $T_{e,non-thermal} = 200$ eV; $T_{e,thermal} = 200$ eV.




Fig. 4. 2D cuts of the background O^{++} ion density profiles. Model I, case (a) (top) and case (b) (bottom). x - y cuts (left column) are located at $z=0$, y - z cuts are located at $x/R_E = 7$, and x - z cuts (right column) are located at $y=0$.

We have performed a several hybrid modeling with different ions and electron betas, and different rate productions for O_2^+ pickup ions but we discuss here only the modeling, which can fit the observations.

The initial thermal velocities of O_2^+ non-thermal and thermal ions are chosen as the following: $v_{th,non-thermal} = 3.0$ km/s (2 eV) and $v_{th,thermal} = 0.5$ km/s (0.05 eV). The initial bulk velocity of O_2^+ pickup ions is about 1 km/s. Eq. (8) gives the following total pickup ion production rate: $Q_{O_2^+,thermal} = 0.825 \times 10^{28} \text{ s}^{-1}$ and $Q_{O_2^+,non-thermal} = 1.95 \times 10^{26} \text{ s}^{-1}$.

Let us consider first the global picture of the interaction of the plasma torus with Europa. The results of this modeling are shown in Figs. 2–4. Figs. 2 and 3 demonstrate 2D cuts for non-thermal and thermal O_2^+ pickup ion density profiles. One can observe the asymmetrical distribution of the pickup ion density (top, case (a)) and (bottom, case (b)) in the x - y , y - z ($x = 5R_E$) and z - x planes. The pickup ion motion is determined mainly by the electromagnetic drift. The motion along the magnetic field is due to the thermal velocity and the gradient of the electron pressure. A more wider density profile of the pickup ions was observed in the case (b), Figs. 2 and 3 (bottom).

The figures demonstrate a strong structuring in the non-thermal and thermal O_2^+ ion density profiles. While case (a) produces a much higher peak in the thermal O_2^+ ion density as was seen in E4 observations, case (b) produces much better agreement with observation for the thermal O_2^+ ion density as shown in Figs. 2 and 3.

The modeling also demonstrates the asymmetrical distribution of the background O^{++} ion density in the x - y , y - z ($x = 5R_E$) and z - x planes, Fig. 4. The asymmetrical distribution of the background ions in the x - y plane may be explained by the existence of a strong B_z component in the upstream magnetic field. One can also see an increase in the plasma density near Europa due to the formation of a magnetic barrier (not shown here). In case (b) this




Fig. 5. 1D cuts of the background O^{++} ion density profile. The cuts are located at $y=0$, $z=0$. Model I, case (a) (top) and case (b) (bottom).

effect is stronger than in case (a). The density profiles for SO^{++} background ions are close to the density profiles for O^{++} ions.

The inclination of the magnetic field results in an asymmetrical boundary condition for ion dynamics (penetration and




Fig. 6. 1D cuts of the background O^{++}/S^{++} , and pickup non-thermal/thermal (O_2^+) ion densities from simulation. $Y(Re)$ denotes a projection of satellite trajectory onto the y axis. Model I, case (a) (left) and case (b) (right). Bottom—E4 observation of the total ion density (Paterson et al., 1999).

reflection) in Europa's ionosphere and an asymmetrical Alfvén wing.

Note that the background ion flow around the effective obstacle that is produced by pickup ions and the ionosphere. The pickup ions flow from the “corona” across the magnetic field due to electromagnetic drift, whereas the motion along the magnetic field is determined by the thermal velocity of ions and the electron pressure.

Fig. 5 demonstrates the 1D cuts ($y=0, z=0$) of the background density O^{++} for case (a) (top) and case (b) (bottom). Strong jumps in the plasma density with $N_{O^{++},max} = 80 \text{ cm}^{-3}$ (case (a)) and $N_{O^{++},max} = 17 \text{ cm}^{-3}$ (case (b)) are observed on the day-side of the ionosphere, whereas a reduction in the plasma density is observed in the plasma wake. Note that the jump in the plasma density profile is stronger in case (a) than it is observed in case (b). Both jumps are located near the surface of Europa.

Fig. 6 shows 1D density profiles of the background and pickup ions along the E4 trajectory of the Galileo spacecraft. One can see a strong plasma void in the center of the plasma wake. There is also a sharp boundary with an overshoot in the density profiles on the side of the plasma wake in the Jupiter-direction, and a smooth boundary layer on the side in the anti-Jupiter direction, Fig. 6 (top). The density profile for O^{++} is similar the density profiles for the S^{++} upstream ions. Fig. 6 (middle and bottom) also shows the density profiles for the non-thermal (top) and thermal (bottom) O_2^+ pickup ions. One can see the split structure of the

plasma tail. The effect of splitting of the plasma tail was also observed in the hybrid modeling of weak comets (see, e.g., Lipatov et al., 1997; Lipatov, 2002). The general feature of this plasma density is due to the effect of the finite heavy gyroradius. The total ion density profile observed in E4 pass is shown in Fig. 6 (bottom). The observed value of the density in these peaks is lower than in modeling and it may be explained by an overestimated density of O_2^+ pickup ions for case (a). In the case (b), disagreement is not as strong, an improvement of the atmosphere model is still required.

The modeling gives the following total fluxes for the O_2^+ pickup ions (case (a)): $1.4 \times 10^{22} \text{ mol/s}$ (non-thermal) and $1.75 \times 10^{25} \text{ mol/s}$ (thermal); (case (b)): $0.8 \times 10^{22} \text{ mol/s}$ (non-thermal) and $1.0 \times 10^{25} \text{ mol/s}$ (thermal) across the back boundary $x = 12R_E$.

Let us consider a global distribution of the electric and magnetic field in Europa's environment. Fig. 7 shows B_x, B_z magnetic and E_y electric field profiles for case (a) (left) and case (b) (right). The $y-z$ cuts (top and middle) are located at $x/R_E = 7$, and $x-y$ cuts (bottom) are located at $y=0$. The figure demonstrates perturbations in the magnetic B_x and electric E_y field profiles, which are due to the formation of an Alfvén wing. The increase in the magnetic field B_z indicates the formation of an asymmetrical magnetic barrier, Fig. 7 (bottom).

The asymmetry of the modeling distributions in \mathbf{B} appears to be caused by the finite gyroradius effects of incoming and pickup ions. A weak perturbation of the magnetic field was observed near




Fig. 7. 2D cuts of the B_x , B_z magnetic and E_y electric field profiles. Model I, case (a) (left) and case (b) (right). y - z cuts (top and middle) are located at $x/R_E = 7$, and y - x cuts (bottom) are located at $z/R_E = 0$.

the ionosphere of Europa: compression of the upstream magnetic field and decompression in the plasma wake.

The modeling also shows the formation of an Alfvén wing in the direction of the main magnetic field. The formation of the Alfvén wing in a sub-Alfvénic flow near Europa is similar to a formation near Io, which was first studied analytically by Neubauer (1980). The pickup ions play important role on the fine structure of the Alfvén wing due to effects of mass loading. In particular, the scale of the front of the Alfvén wing must be determined by the gyroradius of pickup ions. Unfortunately, in our 3D hybrid kinetic simulation we cannot yet resolved this spatial scales.

3.2. Effects of inductive Europa's magnetic field

In the first set of models (Section 3.1, model I, cases (a) and (b)), we used a homogeneous global magnetic field as an initial condition. These models do not produce agreement between the simulated and observed magnetic fields.

In the second set of modeling, we take into account the gradient of the global Jovian magnetic field for an initial magnetic field distribution. In the paper by Kivelson et al. (1999, 1997, 2000), it has been shown that the B_y component of the magnetospheric magnetic field has strong time variations at the position of Europa. In the MHD-fluid approximation, the effects of such

magnetic field variations are estimated in Schilling et al. (2007, 2008). The initial plasma density and bulk velocity distribution in our modeling were taken from the E4 flyby data (Paterson et al., 1999).

We created the following model II, case (c), for simulation: the density for thermal O_2 is the same as for model I, case (b), and the pickup electron temperature is lower than in model I, case (b). The plasma density and bulk velocity distribution in our modeling were taken from the E4 flyby data (Paterson et al., 1999): full magnetosphere corotation speed $U_0 = 105$ km/s; upstream densities are $\rho_{O^{++}} = 10$ cm $^{-3}$; $\rho_{S^{++}} = 10$ cm $^{-3}$; upstream ion and electron temperatures, $T_{O^{++}} = 100$ eV; $T_{S^{++}} = 100$ eV; $T_{e,0} = 20$ eV. The temperatures of electrons connected with non-thermal and thermal O_2^+ pickup ions are $T_{e,non-thermal} = 100$ eV; $T_{e,thermal} = 100$ eV.

In our hybrid kinetic modeling (model II) we use a simple magnetic dipole model of the induced oceanic magnetic field from the 10-hour corotation variation of the background Jovian magnetic field at Europa (see paragraph “Boundary Conditions”,

Section 2). And, finally, we fit the results of modeling to the components of the measured magnetic field.

This is not yet a fully self-consistent approach but provides a first approximation. Also, the ocean may not be exactly a spherically symmetric conducting shell and may ultimately require a higher-order multipole model for the induced fields.

Fig. 8 demonstrates the 2D cuts for non-thermal and thermal O_2^+ pickup ion densities. The figure does not show any extension of the pickup ion profile in the y and z directions. The plasma wake is narrower in y and z directions in compare with that was produced by model I, cases (a) and (b). The reason of this effect is due to lower temperature of electrons connected with pickup O_2^+ ions than it was in a case (b) and lower pickup ions production rate near the surface of Europa than it was in a case (a).

Fig. 9 shows the distribution of the O^{++} ion density in the x - y , y - z ($x = 5 R_E$) and z - x planes. The narrow plasma wake may be explained by smaller temperature of the electrons connected with O_2^+ pickup ions, and, hence, with a smaller polarization electric field which is responsible for an expansion of Europa's ionosphere.




Fig. 8. 2D cuts of the pickup ion O_2^+ density profile. Non-thermal O_2^+ (top), thermal O_2^+ (bottom). Model II, case (c). x - y cuts (left column) are located at $z=0$, y - z cuts are located at $x/R_E = 7$, and x - z cuts (right column) are located at $y=0$.




Fig. 9. 2D cuts of the background O^{++} ion density profiles. Model II, case (c). x - y cuts (left column) are located at $z=0$, y - z cuts are located at $x/R_E = 7$, and x - z cuts (right column) are located at $y=0$.

One can also see an increase in the plasma density near Europa due to the formation of a magnetic barrier (not shown here). The density profile for SO^{++} background ions is close to the density profile for O^{++} ions as in model I, cases (a) and (b).

Fig. 10 shows a 1D cut of the background O^{++} density along the x-axis ($y=0, z=0$). One can see jump in the background plasma density with $N_{O^{++},max} = 90 \text{ cm}^{-3}$ (model II, case (c)) on the day-side of the ionosphere and depletion in the plasma density in Europa's plasma wake. Note that the jump in the plasma density profile is stronger in model II, case (c), than it is observed in model I, case (a). The jump is located near the surface of Europa as it was observed in model I, cases (a) and (b).

Fig. 11 shows 1D density profiles of the background and pickup ions along the E4 trajectory of the Galileo spacecraft. One can see a strong plasma void in the center of the plasma wake. There is also a sharp boundary with an overshoot in the density profiles on the left side of the plasma wake, and a smooth boundary layer on the right side of the plasma wake, Fig. 11 (top). The density profile for S^{++} is similar to the density profile for O^{++} background ions. Fig. 10 (middle) shows the density profiles for non-thermal and thermal O_2^+ pickup ions. The total ion density profile observed during the E4 pass is shown in Fig. 11 (bottom). Again, one can see two peaks in the total ion density profile. However, the observed value of the density in these peaks is lower than predicted by the model; this may be explained by an overestimated density of O_2^+ pickup ions for model II, case (c).

The modeling shows that the shaping of Europa's global plasma environment depends on a combination of the upstream plasma parameters and pickup ions and electron parameters. For example, the reducing in the temperature of electrons connected with pickup ions results in the higher density of the thermal O_2^+ pickup ions at the trajectory of a spacecraft, compare Fig. 6 (right) and Fig. 11. This effect is connected with the polarization electric field which is proportional to the gradient of the electron pressure. The reducing in the temperature of the background upstream ions results in the widening of the plasma wake, compare Fig. 6 (left and right, top) and Fig. 11 (top). These effects were earlier demonstrated in the 3D hybrid simulation of Io's plasma environment (Lipatov and Combi, 2006). We have found the similarities between the plasma environments of these objects. Indeed, Io and Europa have sufficiently thin exospheres and strong magnetic fields resulting in a small value of the ion gyroradius.

Let us consider the global distribution for the electromagnetic field of model II, case (c). Fig. 12 shows 2D cuts for the magnetic B_x , B_z and electric E_y field profiles. The distributions for the B_z , E_y field shown in the figure are close to the distributions for model I, case (b). However, there are significant differences between the B_x profiles for model I, case (a) and case (b), and model II, case (c). The differences between the B_x profiles for cases




Fig. 10. 1D cuts of the background O^{++} ion density profile. The cut is located at $y=0, z=0$. Model II, case (c).




Fig. 11. Background (S^{++} , O^{++}), non-thermal (O_2^+) and thermal (O_2^+) pickup ion densities from simulation. $Y(Re)$ denotes a projection of the spacecraft position onto the y axis. Model II, case (c). Bottom—E4 observation of the total ion density (Paterson et al., 1999).

(a) and (b), Fig. 7 (top) are due to much higher density of the thermal O_2^+ pickup ions in the plasma wake, whereas the differences between the B_x profiles for cases (b) and (c) are due to the nonlinear interaction of the Alfvén wing with nonhomogeneous Jovian magnetic field in model II, case (c).

Fig. 13 shows the magnetic field components (solid line) B_x , B_y , B_z , and $|B|$ along the E4 trajectory of the Galileo spacecraft. The magnetic field components of the inductive magnetic dipole that simulates the effect of the nonstationarity of the Jovian magnetic field are shown by a dotted line (---). The circles (○) denote observational data from Kivelson et al. (1997) and the initial Jovian magnetospheric field at the position of Europa (+++). The simulation produces a satisfactory agreement with the observational data for the B_y magnetic field component, but not for the




Fig. 12. Model II, case (c). 2D cuts of the magnetic B_x (top), electric E_y (top) and magnetic B_z (bottom) field profiles. y - z cuts (top) are located at $x/R_E = 7$, and y - x cuts (bottom) are located at $z/R_E = 0$.

B_x and B_z magnetic field components. A multipole model for the oceanic magnetic field may address this issue. We will need to improve the model of the O_2 atmosphere, the resolution of the ion trajectory, and the gradient in the atmosphere/ionosphere density profiles near the surface of Europa to obtain better agreement in the B_x and B_z magnetic field components

4. Conclusions

Hybrid modeling of Europa's plasma environment for E4 encounter with three ion species demonstrated several features:

- The modeling shows a strong phase mixing in the plasma wake. The plasma wake demonstrates the formation of time-dependent structuring in the pickup ion tails (see, e.g., McKenzie et al., 2001 for weak comet case) and the splitting of the pickup ion tails. The splitting of the plasma wake has the same nature as the splitting of the weak comet's plasma wake or the splitting of Titan's plasma wake. Such finite gyroradius effects were also observed in 2.5D hybrid and bi-fluid

modeling of a weak comet (see, e.g., Lipatov et al., 1997; Sauer et al., 1996, 1997; Lipatov, 2002) and in 3D hybrid modeling of Titan's plasma environment (Lipatov et al., 2011, 2012). The further investigation of these fine structure needs an additional modeling with much better resolution.

- The model shows the magnetic field barrier formation at the day-side portion of the ionosphere. The formation of an Alfvén wing in the plane of the external magnetic field was also observed. Note that the Alfvén wing was earlier observed in a hybrid simulation of the plasma environment of Io and Europa by Lipatov and Combi (2006) and by Lipatov et al. (2010). An MHD simulation of the plasma environment of Io and Europa also produces the formation of an Alfvén wing (Saur et al., 1999, 1998; Liu et al., 2000; Schilling et al., 2008).
- The ion and electron temperatures play an important role in plasma structure formation, and in creating the ion fluxes inside the ionosphere. These effects were observed earlier in a 3D hybrid simulation of Io's plasma environment (Lipatov and Combi, 2006). Hybrid model produces a correct pickup ion flux along the magnetic field in opposite the MHD models which operate with pickup ions with a Maxwellian velocity distribution. In the current paper, we have presented only three runs




Fig. 13. Magnetic field component profiles along the E4 trajectory after fitting with inductive dipole magnetic field. Solid line—modeling, (–) denotes dipole field, and (+) is the Jovian magnetic field at the position of Europa. ○—Galileo's E4 flyby measurements (Kivelson et al., 1997). $X(\text{Re})$ denotes a projection of the spacecraft position onto the x -axis.

with different combinations of the upstream ion and pickup electron temperatures.

- The model's total ion density in the plasma wake does not satisfactory match the observed density.
- The constant induced dipole moment (model II, case (c)) improves a fit the magnetic field B_y component to the E4 trajectory. However, a fit for the magnetic field B_x component is still not satisfactory due to the not perfect model of the atmosphere/ionosphere and not satisfactory numerical resolving of the gyroradius on the grid cell.
- Use of an inhomogeneous background magnetic field provides a good agreement between the observed and simulated magnetic fields. However, we still need to improve the resolution of the gradient in the atmosphere density, the gyroradius of pickup ions, and the fields in the internal non-conduction ice shell and conduction ocean layers of Europa.

In our future computational models, we plan to include a nonstationary boundary condition for the magnetic field in order to take into account the spatially inhomogeneous and nonstationary background Jovian magnetic field. This model will also

respect to a potentially nonspherical ocean shell. We also plan the use of a varying atmospheric density, a varying electron temperature (that plays key-role in the pickup ion dynamics), and sputtering processes (Johnson, 1990; Johnson et al., 1998) at the surface of Europa. We also plan to use a composite grid structure using the “cubed sphere” technique (see, e.g. Koldoba et al., 2002) to improve the resolution of the small scales near the surface of Europa and to increase the size of the computational domain.

The composite grid structure will allow us to estimate the inductive magnetic field from the ocean as a part of the total current closure that also includes the external plasma currents. This technique will allow us to study wave–particle interaction effects in the far plasma wake, such as ion cyclotron waves that have been observed in the Galileo flyby mission (see e.g. Volwerk et al., 2001; Kivelson et al., 2009). These models must include the induced magnetic field from a putative subsurface ocean, and will also include particle trajectory tracing for test particles, e.g. electrons and high-energy ions.

Note that the larger computational domain allows us to use the upstream parameters for the plasma and electromagnetic field instead of the use of the “damping” boundary condition. However, in the outer region of the computational domain (large cell size) we have to use a drift-kinetic approach (see e.g. Lipatov et al., 2005) for ion dynamics since we cannot approximate the ion trajectory there. We can also use a complex particle kinetic technique (see e.g. Lipatov, 2012) which provides a flexible fluid/kinetic description and may significantly save computational resources.

Acknowledgments

A.S.L. was supported in part by the Project/Grant 00004129, and 00004549 between the GPI UMBC and NASA GSFC. J.F.C. was supported as Principal Investigator by the NASA Outer Planets Research Program. Computational resources were provided by the NASA Ames Advanced Supercomputing Division (SGI—Columbia, Project SMD-09-1110).

References

- Austin, J.V., Goldstein, D.B., 2000. Rarefied gas model of Io's sublimation-driven atmosphere. *Icarus* 148, 370–383.
- Bagenal, F., 1994. Empirical model of the Io plasma torus: Voyager measurements. *Journal of Geophysical Research* 99, 11043–11062.
- Braginskii, S.L., 1965. Transport processes in a plasma. In: Leontovich, M.A. (Ed.), *Reviews of Plasma Physics*. Consultants Bureau, New York, pp. 205–240.
- Cassidy, T.A., Johnson, R.E., McGrath, M.A., Wong, M.C., Cooper, J.F., 2007. The spatial morphology of Europa's near-surface O_2 atmosphere. *Icarus* 191, 755–764.
- Combi, M.R., Kabin, K., Gombosi, T., De Zeeuw, D.L., Powell, K., 1998. Io's plasma environment during the Galileo flyby: global three-dimensional MHD modeling with adaptive mesh refinement. *Journal of Geophysical Research* 103, 9071–9081.
- Dols, V., Delamere, P.A., Bagenal, F., 2008. A multispecies chemistry model of Io's local interaction with the Plasma Torus. *Journal of Geophysical Research (Space Physics)* 113, 9208–, doi:10.1029/2007JA012805.
- Goertz, C.K., 1980. Io's interaction with the plasma torus. *Journal of Geophysical Research* 85, 2949–2956.
- Herbert, F., 1985. “Alfvén wing” models of the induced electrical current system at Io: a probe of the ionosphere of Io. *Journal of Geophysical Research* 90, 8241–8251.
- Huddleston, D.E., Strangeway, R.J., Warnecke, J., Russel, C.T., Kivelson, M.G., 1997. Ion cyclotron waves in the Io torus during the Galileo encounter: warm plasma dispersion analysis. *Geophysical Research Letters* 24 (17), 2143–2146.
- Ip, W.-H., 1990. Neutral gas–plasma interaction: the case of the Io plasma torus. *Advances in Space Research* 10 (1), 15–18.
- Jia, X., Walker, R.J., Kivelson, M.G., Khurana, K.K., Linker, J.A., 2008. Three-dimensional MHD simulation of Ganymede's magnetosphere. *Journal of Geophysical Research* 113, A06212.
- Johnson, R.E., 1990. *Energetic Charge–Particle Interaction with Atmospheres and Surfaces*. Springer-Verlag, New York.

- Johnson, R.E., Killen, R.M., Waite, J.H., Lewis, W.S., 1998. Europa's surface composition and sputter-produced ionosphere. *Geophysical Research Letters* 25, 3257–3260.
- Johnson, R.E., Burger, M.H., Cassidy, T.A., Leblanc, F., Marcony, M., Smyth, W.H., 2009. Composition and detection of Europa's sputter-induced atmosphere. In: Pappalardo, R.T., McKinnon, W.B., Khurana, K.K. (Eds.), *Europa*. University of Arizona Press, Tucson, pp. 507–527.
- Kabin, K., Combi, M.R., Gombosi, T.I., Nagy, A.F., DeZeeuw, D.L., Hansen, K.S., Powell, K.G., 1999. On Europa's magnetosphere interaction: a MHD simulation of the E4 flyby. *Journal of Geophysical Research* 104 (A9), 19983–19992.
- Kabin, K., Combi, M.R., Gombosi, T.I., DeZeeuw, D.L., Hansen, K.S., Powell, K.G., 2001. Io's magnetospheric interaction: an MHD model with day-night asymmetry. *Planetary and Space Science* 49, 337–344.
- Khurana, K.K., Kivelson, M.G., Stevenson, D.J., Schubert, G., Russell, C.T., Walker, R.J., Polansky, C., 1998. Induced magnetic fields as evidence for subsurface oceans in Europa's and Callisto. *Nature* 395, 777–780.
- Kivelson, M.G., Khurana, K.K., Means, J.D., Russell, C.T., Snare, R.C., 1992. The Galileo magnetic field investigation. *Space Science Reviews* 60, 357–383.
- Kivelson, M.G., Khurana, K.K., Joy, S., Russell, C.T., Southwood, D., Walker, R.J., Polansky, C., 1997. Europa's magnetic signature: report from Galileo's pass on 19 December 1996. *Science* 276, 1239–1241.
- Kivelson, M.G., Khurana, K.K., Stevenson, D.J., Benett, L., Joy, S., Russell, C.T., Walker, R.J., Polansky, C., 1999. Europa and Callisto: induced or intrinsic fields in periodically varying plasma environment. *Journal of Geophysical Research* 104, 4609–4625.
- Kivelson, M.G., Khurana, K.K., Russell, C.T., Volwerk, M., Walker, R.J., Zimmer, C., 2000. Galileo magnetometer measurements strengthen the case for a subsurface ocean at Europa. *Science* 289, 1340–1343.
- Kivelson, M.G., Khurana, K., Volwerk, M., 2009. Europa's interaction with the Jovian magnetosphere. In: Pappalardo, R.T., McKinnon, W.B., Khurana, K.K. (Eds.), *Europa*. University of Arizona Press, Tucson, pp. 545–570.
- Koldoba, A.V., Romanova, M.M., Ustyugova, G.V., Lovelace, R.V.E., 2002. Three dimensional MHD simulation of accretion to an inclined rotator: the “cubed sphere” method. *Astrophysical Journal* 576, L53–L56.
- Lellouch, E., 1996. Urey Prize Lecture. Io's atmosphere: not yet understood. *Icarus* 124, 1–21.
- Linker, J.A., Khurana, K.K., Kivelson, M.G., Walker, R.J., 1998. MHD simulation of Io's interaction with the plasma torus. *Journal of Geophysical Research* 103 (E9), 19867–19877.
- Lipatov, A.S., Sauer, K., Baumgärtel, K., 1997. 2.5-D hybrid code simulation of the solar wind interaction with weak comets and related objects. *Advances in Space Research* 20 (2), 279–282.
- Lipatov, A.S., 2002. The hybrid multiscale simulation technology. In: *An Introduction with Application to Astrophysical and Laboratory Plasmas*. Springer-Verlag, Berlin, Heidelberg, New York, p. 413.
- Lipatov, A.S., Motschmann, U., Bagdonat, T., Grießmeier, J.-M., 2005. The interaction of the stellar wind with an extrasolar planet—3D hybrid and drift-kinetic simulations. *Planetary and Space Science* 53, 423–432.
- Lipatov, A.S., Combi, M.R., 2006. Effects of kinetic processes in shaping Io's global plasma environment: a 3D hybrid model. *ICARUS* 180, 412–427.
- Lipatov, A.S., Cooper, J.F., Paterson, W.R., Sittler Jr., E.C., Hartle, R.E., 2010. Jovian plasma torus interaction with Europa: 3D hybrid kinetic simulation. *First Results, Planetary and Space Science* 58 (13), 1681–1691.
- Lipatov, A.S., Sittler Jr., E.C., Hartle, R.E., Cooper, J.F., Simpson, D.G., 2011. Background and pickup ion velocity distribution dynamics in Titan's plasma environment: 3D hybrid simulation and comparison with CAPS observations. *Advances in Space Research* 48, 1114–1125.
- Lipatov, A.S., Sittler Jr., E.C., Hartle, R.E., Cooper, J.F., Simpson, D.G., 2012. Saturn's magnetosphere interaction with Titan for T9 encounter: 3D hybrid modeling and comparison with CAPS observations. *Planetary and Space Science* 61, 66–78.
- Lipatov, A.S., 2012. Merging for particle–mesh complex particle kinetic modeling of the multiple plasma beams. *Journal of Computational Physics* 231, 3101–3118.
- Liu, Y., Nagy, A.F., Kabin, K., Combi, M.R., DeZeeuw, D.R., Gombosi, T.I., Powell, K.G., 2000. Two species, 3D MHD simulation of Europa's interaction with Jupiter's magnetosphere. *Geophysical Research Letters* 27, 1791–1794.
- Mankofsky, A., Sudan, R.N., Denavit, J., 1987. Hybrid simulation of ion beams in background plasma. *Journal of Computational Physics* 70, 89–116.
- Marconi, M.L., Dagum, L., Smyth, W.H., 1996. Hybrid fluid/kinetic approach to planetary atmospheres: an example of an intermediate-mass body. *Astrophysical Journal* 469, 393–401.
- McGrath, M.A., Johnson, R.E., 1989. Charge exchange cross sections for the io plasma torus. *Journal of Geophysical Research* 94 (A3), 2677–2683.
- McKenzie, J.F., Sauer, K., Dubinin, E.M., 2001. Stationary waves in a bi-ion plasma transverse to the magnetic field. *Journal of Plasma Physics* 65 (3), 197–212.
- Neubauer, F.M., 1980. Nonlinear standing Alfvén wave current system at Io: Theory. *Journal of Geophysical Research* 85, 1171–1178.
- Neubauer, F.M., 1998. The sub-Alfvénic interaction of the Galilean satellites with the Jovian magnetosphere. *Journal of Geophysical Research* 103, 19834–19866.
- Neubauer, F.M., 1999. Alfvén wing and electromagnetic induction in the interiors: Europa and Callisto. *Journal of Geophysical Research* 104, 28671–28684.
- Paterson, W.R., Frank, L.A., Ackerson, K.L., 1999. Galileo plasma observation at Europa: ion energy spectra and moments. *Journal of Geophysical Research* 104 (A10), 22779–22791.
- Sauer, K., Bogdanov, A., Baumgärtel, K., Dubinin, E., 1996. Plasma environment of comet Wirtanen during its low-activity stage. *Planetary and Space Science* 44 (7), 715–729.
- Sauer, K., Lipatov, A.S., Baumgärtel, K., Dubinin, E., 1997. Solar wind–pluto interaction revised. *Advances in Space Research* 20 (2), 295.
- Saur, J., Strobel, D.F., Neubauer, F.M., 1998. Interaction of the Jovian magnetosphere with Europa: constraints on the neutral atmosphere. *Journal of Geophysical Research* 103 (E9), 19947–19962.
- Saur, J., Neubauer, F.M., Strobel, D.F., Summers, M.E., 1999. Three-dimensional plasma simulation of Io's interaction with the Io plasma torus: asymmetric plasma flow. *Journal of Geophysical Research* 104, 25105–25126.
- Schilling, N., Neubauer, F.M., Saur, J., 2007. Time-varying interaction of Europa with the Jovian magnetosphere: constraints on the conductivity of Europa's subsurface ocean. *Icarus* 192, 41–55.
- Schilling, N., Neubauer, F.M., Saur, J., 2008. Influence of the internally induced magnetic field on the plasma interaction of Europa. *Journal of Geophysical Research* 113, A03203, <http://dx.doi.org/10.1029/2007JA012842>.
- Schreier, R., Eviatar, A., Vasyliūnas, V.M., Richardson, J.D., 1993. Modeling the Europa plasma torus. *Journal of Geophysical Research* 98, 21231–21243.
- Shematovich, V.I., Johnson, R.E., Cooper, J.F., Wong, M.C., 2005. Surface-bounded atmosphere of Europa. *Icarus* 173, 480–498.
- Sittler Jr., E.C., Strobel, D.F., 1987. Io plasma torus electrons: Voyager 1. *Journal of Geophysical Research* 92, 5741–5762.
- Smyth, W.H., Marconi, M.L., 2006. Europa's atmosphere, gas tori, and magnetospheric implications. *Icarus* 181, 510–526.
- Southwood, D.J., Kivelson, M.G., Walker, R.J., Slavin, J.A., 1980. Io and its plasma environment. *Journal of Geophysical Research* 85, 5959–5968.
- Southwood, D.J., Dunlop, M.W., 1984. Mass pickup in sub-Alfvénic plasma flow: a case study for Io. *Planetary and Space Science* 32, 1079–1089.
- Tikhonov, A.N., Samarskii, A.A., 1963. *Equations of Mathematical Physics*. Mac Millan, New York, p. 765.
- Umeda, T., Omura, Y., Matsumoto, H., 2001. An improved masking method for absorbing boundaries in electromagnetic particle simulations. *Computer Physics Communications* (137), 286–299.
- Volwerk, M., Kivelson, M.G., Khurana, K.K., 2001. Wave activity in Europa's wake: implications for ion pickup. *Journal of Geophysical Research* 106 (A11), 26033–26048.
- Winske, D., Wu, C.S., Li, Y.Y., Mou, Z.Z., Guo, S.Y., 1985. Coupling of newborn ions to the solar wind by electromagnetic instabilities and their interaction with the bow shock. *Journal of Geophysical Research* 90, 2713–2726.
- Wolf-Gladrow, D.A., Neubauer, F.M., Lussem, M., 1987. Io's interaction with the plasma torus: a self-consistent model. *Journal of Geophysical Research* 92, 9949–9961.