

The Atacama Cosmology Telescope: Dusty Star-Forming Galaxies and Active Galactic Nuclei in the Southern Survey

Danica Marsden^{1,2}, Megan Gralla³, Tobias A. Marriage³, Eric R. Switzer⁴, Bruce Partridge⁵, Marcella Massardi⁶, Gustavo Morales⁷, Graeme Addison⁸, J Richard Bond⁴, Devin Crichton³, Sudeep Das⁹, Mark Devlin², Rolando Dünner⁷, Amir Hajian⁴, Matt Hilton¹⁰, Adam Hincks⁴, John P. Hughes¹¹, Kent Irwin¹², Arthur Kosowsky¹³, Felipe Menanteau¹¹, Kavilan Moodley¹⁰, Michael Niemack¹⁴, Lyman Page¹⁵, Erik D. Reese², Benjamin Schmitt², Neelima Sehgal¹⁶, Jonathan Sievers^{4,15,10}, Suzanne Staggs¹⁵, Daniel Swetz¹², Robert Thornton¹⁷, Edward Wollack¹⁸

¹Department of Physics, University of California, Santa Barbara, CA 93106, USA

²Department of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA, USA 19104

³Dept. of Physics and Astronomy, The Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218-2686

⁴Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, ON, Canada M5S 3H8

⁵Department of Physics and Astronomy, Haverford College, 370 Lancaster Avenue, Haverford, PA, USA 19041

⁶INAF, Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

⁷Departamento de Astronomía y Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile

⁸Department of Physics and Astronomy, University of British Columbia, Vancouver, BC, Canada V6T 1Z4

⁹Berkeley Center for Cosmological Physics, LBL and Department of Physics, University of California, Berkeley, CA, USA 94720

¹⁰Astrophysics and Cosmology Research Unit, School of Mathematics, Statistics & Computer Science, University of KwaZulu-Natal, Durban, 4041, South Africa

¹¹Department of Physics and Astronomy, Rutgers, The State University of New Jersey, Piscataway, NJ USA 08854-8019

¹²NIST Quantum Devices Group, 325 Broadway Mailcode 817.03, Boulder, CO, USA 80305

¹³Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA, USA 15260

¹⁴Department of Physics, Cornell University, Ithaca, NY 14853

¹⁵Jadwin Hall, Princeton University, Princeton, NJ, USA 08544

¹⁶Physics and Astronomy Department, Stony Brook University, Stony Brook, NY 11794-3800, USA

¹⁷Department of Physics, West Chester University of Pennsylvania, West Chester, PA, USA 19383

¹⁸NASA/Goddard Space Flight Center, Greenbelt, MD, USA 20771

11 June 2013

ABSTRACT

We present a catalog of 191 extragalactic sources detected by the Atacama Cosmology Telescope (ACT) at 148 GHz and/or 218 GHz in the 2008 Southern survey. Flux densities span 14–1700 mJy, and we use source spectral indices derived using ACT-only data to divide our sources into two sub-populations: 167 radio galaxies powered by central active galactic nuclei (AGN), and 24 dusty star-forming galaxies (DSFGs). We cross-identify 97% of our sources (166 of the AGN and 19 of the DSFGs) with those in currently available catalogs. When combined with flux densities from the Australian Telescope 20 GHz survey and follow-up observations with the Australia Telescope Compact Array, the synchrotron-dominated population is seen to exhibit a steepening of the slope of the spectral energy distribution from 20 to 148 GHz, with the trend continuing to 218 GHz. The ACT dust-dominated source population has a median spectral index, $\alpha_{148-218}$, of $3.7^{+0.62}_{-0.86}$, and includes both local galaxies and sources with redshifts as great as 5.6. Dusty sources with no counterpart in existing catalogs likely belong to a recently discovered subpopulation of DSFGs lensed by foreground galaxies or galaxy groups.

Key words: galaxies: surveys – galaxies: active – millimeter: galaxies

1 INTRODUCTION

The technologies that enable observations of large numbers of millimeter and submillimeter sources were developed relatively recently, opening up a previously unexplored regime for study that has the power to reveal the evolution of underlying galaxy populations over cosmic time, and in particular through epochs of intense star formation. Instruments such as the Submillimeter Common User Bolometer Array (SCUBA, SCUBA-2; Holland et al. 1999; Holland, W. S. et al. 2013), the Balloon-borne Large Area Submillimeter Telescope (BLAST; Pascale, E. et al. 2008), the AzTEC millimeter wavelength camera (Wilson et al. 2008) and the Large Apex Bolometer Camera (LABOCA; Siringo et al. 2009), have mapped up to 10 deg^2 of the sky, but greater coverage was limited by the large amount of integration time required to conduct blind surveys to significant cosmological depth. Increased sky coverage and sensitivity motivated the construction of space-based observatories such as *Spitzer* (Werner, M. W. et al. 2004) and *Herschel* (Pilbratt et al. 2010), operating in the wavelength regime $4\text{--}500 \mu\text{m}$. At longer wavelengths, the Wilkinson Microwave Anisotropy Probe (*WMAP*; Wright, E. L. et al. 2009) covered the whole sky at frequencies up to 94 GHz, but observations of unresolved extragalactic sources were only complete above 2 Jy due to its large beam size (roughly $13'$ at 94 GHz). The *Planck* space telescope contains a high frequency instrument observing at frequencies spanning 100–857 GHz (Lamarre, J.-M. et al. 2010). At 143 and 217 GHz, the *Planck* beam sizes are approximately $7'$ and $5'$ respectively, allowing for a source catalog complete down to flux densities of about 1 Jy. Large-area ($> 100 \text{ deg}^2$) ground-based radio surveys probe with higher resolution than *WMAP* or *Planck*, but only as high in frequency as 20 GHz (e.g., the Australia Telescope 20 GHz survey; Murphy, T. et al. 2010). Thus there exists a niche for millimeter wavelength large-scale mapping experiments with spatial resolution superior to that of the space-based observatories.

The Atacama Cosmology Telescope (ACT; Swetz, D. S. et al. 2011) made temperature anisotropy maps of the Cosmic Microwave Background (CMB) with arcminute resolution from its 2007 to 2010 observing seasons (Dünner, R. et al. 2013) at 148, 218 and 277 GHz. These maps contain galaxies that are luminous at millimeter wavelengths. Measurements of the millimeter fluxes of a large sample of sources have great potential to discriminate among source population models and reveal new source populations. For example, recent millimeter surveys (e.g., Vieira, J. D. et al. 2010; Marriage, T. A. et al. 2011; Planck Collaboration VII 2013) have yielded source counts that led to updated source models (Tucci et al. 2011) and the discovery of a new class of high-flux dusty galaxies (this publication; Vieira, J. D. et al. 2010; Negrello, M. et al. 2010).

ACT-detected sources at flux densities greater than 20 mJy are predominantly blazars, which have spectral energy distributions (SEDs) that are synchrotron-dominated. These galaxies are powered by active galactic nuclei (AGN), black holes at their centers accreting nearby matter in a process that produces time-variable jets out of the plane of the accretion disk; in the case of blazars, the jet direction lies close to our line of sight. Ejected ionized particles spiraling around magnetic field lines in the jets create the observed synchrotron emission. The study of blazar SEDs and source counts provides information about AGN physics (e.g., de Zotti et al. 2010; Tucci et al. 2011; Planck Collaboration XV 2011).

The second population identified at ACT wavelengths is comprised of infrared-luminous, dusty star-forming galaxies (DSFGs),

which exhibit modified blackbody emission at sub-millimeter to millimeter wavelengths, diminishing toward longer wavelengths. The observed SED is dominated by thermal emission from dust grains that have been heated by the prodigious optical and ultraviolet flux produced by newly formed stars (e.g., Draine 2003, and references therein). As well as the thermal dust emission, some of these galaxies may have a significant non-thermal contribution from AGN at their cores.

Some of these DSFGs are local galaxies ($z \ll 1$), cross-identified with Infrared Astronomy Satellite (IRAS) sources ($\lambda = 12\text{--}100 \mu\text{m}$; Devereux & Young 1990). The IRAS wavebands are relatively insensitive to dust with temperatures below 30 K, a significant and largely unexplored component of many nearby galaxies (Planck Collaboration XVI 2011). Recent results from *Herschel* (Amblard, A. et al. 2010) and BLAST (Wiebe, D. V. et al. 2009) have begun to extend our picture of the cold dust in galaxies, but millimeter-wavelength experiments can, through probing dust in nearby galaxies, contribute to establishing a well-calibrated dust SED for typical, low-redshift DSFGs. This in turn has cosmological ramifications, as current analyses of the first generations of stars and galaxies that fuel the cosmic infrared background (CIB) rely on an understanding and extrapolation of dust SED templates.

High-redshift DSFGs were observed by SCUBA in the first systematic survey of these sources, which create a significant fraction of the CIB emission (Blain et al. 2002). Selected at $850 \mu\text{m}$, these sources came to be known as submillimeter galaxies (SMGs). This new population of galaxies subsequently became a focus of observations (e.g., Weiß, A. et al. 2009; Viero, M. P. et al. 2009; Austermann, J. E. et al. 2010), and stacking analyses that resolved more of the CIB into emission from discrete, dusty star-forming galaxies (e.g., Dole et al. 2006; Devlin, M. J. et al. 2009). Though most SMGs will be undetectable by current millimeter survey instruments, a new population of sources significantly brighter and rarer than the submillimeter-selected SMGs, similarly exhibiting dust-dominated spectral indices, has been identified at millimeter wavelengths (Vieira, J. D. et al. 2010). They do not have counterparts in the IRAS catalog, indicating that they are not members of the standard, local, ultraluminous infrared galaxy (ULIRG) population.

These sources have recently been shown to be a new, higher redshift ($z > 3$) sub-population of the progenitor galaxy background, brought to the fore in millimeter-wave surveys because they are lensed by foreground galaxies or galaxy groups (Negrello et al. 2007; Lima et al. 2010; Negrello, M. et al. 2010; Vieira, J. D. et al. 2013; Hezaveh, Y. D. et al. 2013). Such objects are extremely rare; at submillimeter wavelengths, see, e.g., Rex, M. et al. (2010); Lupu, R. E. et al. (2012), but wide area surveys will find more. The South Pole Telescope (SPT; Carlstrom, J. E. et al. 2011) has reported significant numbers of these sources. This population provides an avenue for follow-up research to study the details not only of lensed SMGs, but also of the lens systems (e.g., Ikarashi, S. et al. 2011; Scott, K. S. et al. 2011; Lupu, R. E. et al. 2012; Weiß, A. et al. 2013).

In this paper, we report for the first time, the discovery of DSFGs in the ACT data. This is the first multifrequency analysis of the ACT sources, and the second report on ACT extragalactic sources. Marriage, T. A. et al. (2011), hereafter M11, presented a catalog of sources at 148 GHz only from the 455 deg^2 of the 2008 ACT Southern survey with the best uniformity and coverage. Here we extend those results to include 218 GHz (and updated 148 GHz) flux densities for the same 2008 ACT Southern survey region from the new




Figure 1. Sensitivity maps with source detections at 148 GHz (top) and 218 GHz (bottom), showing the most uniform 455 deg^2 patch of the *Southern strip* with the greatest depth of coverage. It lies between right ascension $00^{\text{h}}12^{\text{m}}$ and $07^{\text{h}}07^{\text{m}}$, and declination $-56^{\circ}11'$ and $-49^{\circ}00'$. The deepest data correspond to an exposure time of 23.5 minutes per square arcminute and a 1σ sensitivity of 2.34 mJy at 148 GHz and 3.66 mJy at 218 GHz. White circles or squares mark the locations of ACT sources with a size proportional to the log of the associated source flux density. Circles denote sources designated as AGN, and squares denote sources with spectra indicative of DSGFs. Toward the edges of the map, the variation in local noise properties due to uneven coverage is more apparent.

(Dünner, R. et al. 2013, hereafter D13) data release maps. A final study (Gralla et al., in prep.) of sources detected in the ACT *Equatorial strip* is in preparation.

The layout of the paper is as follows. Section 2 describes the 2008 season ACT observations and the reduction of raw data into maps, as well as follow-up observations made with the Australian Telescope Compact Array (ATCA). Section 3 details our method of source extraction. The source catalog is discussed in Section 4, as well as its astrometric and flux density accuracy, and the catalog’s estimated completeness and purity. Section 5 compares our catalog with currently available data sets, and Section 6 gives the source number counts. Trends observed in the spectral indices of our source populations are analyzed in Section 7. We conclude in Section 8 with a summary of our results.

2 OBSERVATIONS AND DATA

2.1 ACT Observations

The ACT experiment (Swetz, D. S. et al. 2011) is situated on the slopes of Cerro Toco in the Atacama Desert of Chile¹ at an elevation of 5,190 m. ACT’s latitude gives access to both the northern and southern celestial hemispheres. Observations occurred simultaneously in three frequency bands, at 148 GHz (2.0 mm), 218 GHz (1.4 mm) and 277 GHz (1.1 mm) with angular resolutions of roughly $1.4'$, $1.0'$ and $0.9'$ respectively. Observations of Saturn were used to determine beam profiles and pointing (Hincks, A. D. et al. 2010; Hasselfield, M. et al. 2013). From 2007 to 2010, ACT targeted two survey regions: the *Southern strip* centered around $\delta = -52.5^\circ$ and the *Equatorial strip* centered around $\delta = 0^\circ$. Further information about the ACT observations can be found in D13.

¹ 22.9586° south latitude, 67.7875° west longitude.

2.2 ACT Data

The reduction of raw ACT data into maps is detailed in D13; we will briefly review that process here. Each ACT detector array (one per frequency band) is composed of 1,024 detectors, with each detector timestream first analyzed and then kept or rejected based on multiple criteria, such as telescope operation, weather conditions, cosmic ray hits, or other interference. Approximately 800 (700) hours of data from the 2008 *Southern strip* for 148 GHz (218 GHz) remain after these cuts.

Maximum likelihood maps of pixels $30''$ on a side are produced from the timestream data. From an initial estimate of the maps, source profiles are fit for. We then subtract the timestream models for the point sources from the data and re-map. This prevents the power from being aliased into the map, and improves the final flux density estimates for $S/N > 5$ sources. Similarly, an estimate of the CMB signal is also removed. The mapping equation (Tegmark 1997) is then solved iteratively using the preconditioned conjugate gradient method. Finally, the signals that had been subtracted are added back to the map.

The measured source flux densities have converged on a single value by the 25th iteration of the maps used in our study. For 148 GHz, the D13 release map is from iteration 1000 of the map-maker, and we use this map both for convenience and in order that this study be publicly reproducible. For our study of source flux densities at 218 GHz, we use iteration 200, which is more than adequate. Between the map iterations used in this study, the fluxes of the sources change less than 1%. One may ask whether the value to which the measured source flux density has converged is accurate. Based on end-to-end simulations in which mock sources are injected into the ACT time streams, we estimate this accuracy at 3%.

The 148 GHz map is calibrated in temperature through cross comparison of spectra over the range $400 < \ell < 1000$ with WMAP as in Hajian, A. et al. (2011) with an uncertainty of 2%. This calibration is transferred to the 218 GHz map (2.4% uncertainty) through cross-calibration with 148 GHz. Hasselfield, M. et al.

(2013) used this calibration in measurements of the temperatures of Uranus and Saturn. The resulting planet temperature estimates were consistent with previous analyses (Griffin & Orton 1993; Goldin et al. 1997), lending confidence that the WMAP-based calibration at large angular scales will also apply to the calibration of sources at small angular scales.

Photometry based on matched filtering (Section 3) relies on an accurate estimate of the instrument beam shape. The impact of beam error on photometry depends on the details of the filter (Equation 1), which depend on both the beam window function and the angular power spectrum of the background and noise. We have found that the error in the solid angle of the beam is a conservative proxy for the photometric error due to beam uncertainty, and we use this simpler quantity (solid angle uncertainty) in the beam-related photometry error estimates described here. Due to uncertainties in the beam measurements from Saturn we assign 1% photometric errors for both bands. The profiles of bright point sources in the survey maps suggest that the effective beam for the survey is slightly broader than that measured from Saturn. The survey maps consist of overlapping observations taken over the entire season, and the observed broadening may be attributed to a night-by-night jitter in the telescope pointing with an rms of $5 \pm 1''$ (Hasselfield, M. et al. 2013). The uncertainty in jitter correction corresponds to 1% (148 GHz) and 2% (218 GHz) photometric uncertainties, which are correlated between bands.

Finally, for our photometric uncertainty budget we account for the fact that AGN have a lower effective frequency band center (and thus slightly broader beam) than Saturn, and DSFGs have a higher effective frequency band center (and thus narrower beam; Swetz, D. S. et al. 2011, Table 4). We choose effective 148 GHz and 218 GHz frequency centers corresponding to halfway between a steep spectrum AGN and a DSFG: 148.65 GHz and 218.6 GHz. This choice introduces a photometric bias of less than 1.5% at 148 GHz and 1.1% at 218 GHz, which is positive for DSFGs and negative for steep spectrum AGN. We fold this photometric bias from the source spectrum in with uncertainties due to mapping (3%), WMAP calibration (2%, 2.4%) and beam shape (1.4%, 2.2%) to obtain an overall flux density calibration uncertainty of 4.1% at 148 GHz and 4.6% at 218 GHz. As shown in Section 5, the ACT flux densities agree with independent measurements to within this margin of error.

For this study, we have used the most uniform 455 deg² of the 2008 ACT *Southern strip* at 148 and 218 GHz. This region, shown in Figure 1, spans declination $-56.2^\circ < \delta < -49.0^\circ$ and right ascension $00^{\text{h}}12^{\text{m}} < \alpha < 07^{\text{h}}07^{\text{m}}$. At 148 GHz we use the data publicly released with D13². Typical white noise levels in this region of the map are 30 μK -arcminute at 148 GHz and 50 μK -arcminute for 218 GHz. As described in Section 3, when matched-filtered with the ACT beam, this white noise level results in a 1σ point source flux density sensitivity in the best covered regions of $\sigma_0 = 2.34$ mJy at 148 GHz and 3.66 mJy at 218 GHz. The sensitivity levels in Figure 1 are proportional to the square root of the number of observations at that map location, N_{obs} , with one observation per 0.005 seconds. Then the sensitivity level in a given portion of the map is $\sigma = \sigma_0 \sqrt{N_{\text{obs,max}}/N_{\text{obs}}}$.

2.3 ATCA Observations

Marriage, T. A. et al. (2011) found that the sample of ACT

148 GHz-detected sources cross-identified with the Australian Telescope 20 GHz survey (AT20G; Murphy, T. et al. 2010) is dominated by sources with peaked or falling SEDs using flux densities measured at 5, 20 and 148 GHz. The study also confirmed the findings of the AT20G study (Murphy, T. et al. 2010; Massardi, M. et al. 2011), that this population of radio sources is characterized, on average, by spectral steepening between 20–30 GHz and 148 GHz. However, this sample from M11 was incomplete, biased in a way that favored sources with negative spectral indices between 20 and 148 GHz due to the AT20G survey completeness level of 78% above 50 mJy at 20 GHz. ACT-selected sources with 148 GHz flux densities less than 50 mJy and flat or rising spectra may not have been detected by AT20G. Therefore a targeted set of measurements of flux densities at 20 GHz were made for the M11 sources not identified with any source in the AT20G catalog within a 30'' search radius in order to complete the M11 20–148 GHz spectral study.

Scheduling and weather constraints permitted us to observe 41 of these 48 sources with the 6×22 m antenna array of ATCA³ over 6.5 hours at 20 GHz on November 2010, when the array was in its East-West 750A configuration, with 15 baselines ranging from 77 to 3750 m. The synchrotron-dominated spectra of these sources, as revealed in M11, indicated that they were compact (AGN), and would be unresolved. The primary beam of each array telescope at 20 GHz is $2.3'$, with a resolution of $0.5''$ possible with the full array in this configuration. For these observations, the two 2 GHz-wide frequency bands of the new ATCA Compact Array Broadband Backend (CABB; Wilson, W. E. et al. 2011) digital array correlators were set to be adjacent by centering them at 19 GHz and 21 GHz. The reduced average flux densities over the whole bandwidth of the correlator corresponds to the “20 GHz” flux density. Good weather conditions prevailed throughout the observations, from 07:35:24.9 UT through 14:11:54.9 UT. A few data blocks were flagged and removed in order to minimize noise and any spurious effects.

The primary flux calibrator used for these observations was PKS B1934-638, and the bandpass calibrator used was PKS B1921-293. The target sources were expected to have flux densities at or just below the AT20G survey limit of 40 mJy. Each target source was observed once for 1.5 minutes for an rms noise level of ≤ 0.15 mJy/beam. Target observations were interleaved with observations of four secondary calibrators for pointing and phase corrections, chosen to lie close to the targets.

2.4 ATCA Data

The ATCA follow-up data was reduced using a fully automated, custom, shell script pipeline based on tasks from the MIRIAD aperture synthesis reduction package (Sault et al. 1995). An initial inspection of the data was performed to identify contamination or any problems in the data acquisition. Automatic procedures were used to identify and flag data for each frequency band affected by shadowing or known radio contamination, resulting in less than 1% of the band being flagged and cut. The pipeline then generated the calibration solutions for bandpass, flux density amplitude and phase based on the calibrators. After checking that results were consistent between the two frequency bands, we merged the calibrated target visibilities before extracting the flux densities to improve sensitivity.

² <http://lambda.gsfc.nasa.gov/product/act/>

³ <http://www.narrabri.atnf.csiro.au/>

The data analysis of the AT20G survey (Murphy, T. et al. 2010) and of other ATCA projects (Massardi et al. 2011; Bonavera et al. 2011) have shown that the triple correlation technique (Thompson et al. 1986) can be effectively used to determine source flux densities down to tenth of mJy scales for point-like sources unaffected by poor phase stability. This technique is effective for point sources even if they are not in the phase center and in the case of low S/N.

We manually inspected the 20 GHz deconvolved images, and wherever a source was clearly identifiable we measured its flux density and flux density error at its peak in the image and the rms of the image pixels in a region unaffected by the source emission. For sources with high S/N, deconvolution techniques were able to reconstruct images, despite the poor uv -coverage of our observations. In the few cases where the source appeared extended with respect to the synthesized beam we estimated the flux density by integrating over the source area. In some cases, the image showed a confused field with multiple peaks, primarily due to sidelobes; for these cases we extracted a flux density estimate using the triple correlation technique. The flux density errors were obtained from the rms of the visibilities of the V Stokes parameter. This assumes that these objects have negligible circular polarization, which is an overestimate in the cases where the source is not in the phase center. The same statistic (V Stokes rms) was used to indicate the noise level reached for fields where no source was identified at 20 GHz. Of the 41 ACT sources observed in the follow-up campaign described here, all had a measured flux density at 20 GHz. The final results of this study are included as part of Table 4 with an asterisk denoting that a source was part of the follow-up ATCA sample.

3 ACT SOURCE EXTRACTION

We use the same matched filtering method as described in M11 to produce an estimate of the amplitude of a source at the location of the source center. In this section we summarize the method with an emphasis on aspects for which this analysis differs from M11.

Initially, the data are weighted by the square root of the number of observations per pixel, which results in a map with an approximately constant white noise level and a natural apodization as the number of observations falls off toward the edge of the survey region. New to this study, we filter the entire D13 148 GHz release map and corresponding 218 GHz map, which extend beyond the deep 455 deg² area of this study, in order to eliminate any potential edge effects. We then use only the 455 deg² region for source identification.

The following filter is applied to the Fourier transform of the map:

$$\Phi(\mathbf{k}) = \frac{F_{k_0, k_x}(\mathbf{k}) \tilde{B}^*(\mathbf{k}) |\tilde{T}_{\text{other}}(\mathbf{k})|^{-2}}{\int \tilde{B}^*(\mathbf{k}') F_{k_0, k_x}(\mathbf{k}') |\tilde{T}_{\text{other}}(\mathbf{k}')|^{-2} \tilde{B}(\mathbf{k}') d\mathbf{k}'}, \quad (1)$$

where $\mathbf{k} = (k_x, k_y)$ is the angular wave number, with x and y referring to the right ascension and declination directions. $B(\mathbf{k})$ is the Fourier transform of the “effective” instrument beam. As described in Section 2, the “instantaneous” beam is derived from planet observations (Hasselfield, M. et al. 2013). The “effective beam” in the 2008 survey map is broadened by imperfect telescope repointing with mean deviation $\sigma_\theta = 5''$. This broadening is included in $B(\mathbf{k})$ by multiplying the instantaneous beam transform (released with D13 for 148 GHz) by $\exp(-\ell^2 \sigma_\theta^2 / 2)$. \tilde{T}_{other} is the Fourier transform of all components of the data besides point




Figure 2. Matched filters (solid lines) are band pass filters that smooth the map on the scale of the beam (dashed lines) and remove large scale structure associated with the CMB, other astrophysical signals (e.g., the Sunyaev-Zel’dovich effect) and residual contamination from atmospheric brightness fluctuations. The beam and filter functions are plotted with unit normalization at their peak ($\ell = 0$). The matched filters were binned in radius to better show the relevant angular scales. In the analysis, we use two dimensional filters to capture the anisotropic character of the noise.

sources (i.e., atmospheric or detector noise, CMB, etc.). The function $F_{k_0, k_x}(\mathbf{k})$ is a high-pass filter that removes undersampled large scale modes below $k_0 = 1000$ and modes with $|k_x| < 100$, which are contaminated in a fraction of ACT data by telescope scan-synchronous noise. While $B(\mathbf{k})$ is well approximated as azimuthally symmetric, we retain the full two dimensional power spectrum $|\tilde{T}_{\text{other}}(\mathbf{k}')|^2$ to downweight anisotropic noise in the maps. The azimuthally-binned 148 and 218 GHz real space filters and associated beam profiles are shown in Figure 2.

The power spectrum $|\tilde{T}_{\text{other}}(\mathbf{k}')|^2$ used in this analysis was constructed in a different manner from M11: we used the power spectrum of the data itself instead of the average of difference (noise) maps and models for the CMB and other contaminating sky emission. This estimate is robust since the total power from the extragalactic source signal is low compared to the CMB, atmospheric noise, and white noise. To avoid a noisy estimate of the power spectrum, we smooth the power spectrum $|\tilde{T}_{\text{other}}(\mathbf{k}')|^2$ with a Gaussian. The exact formulation of the smoothing does not significantly change the resulting filtered map.

Applying the matched filter can cause ringing in the maps about the very brightest sources which impacts source extraction for low S/N sources. Therefore we identify sources with $S/N > 50$ in an initial application of the filter, and mask them in the maps. Sources with $S/N < 50$ are then extracted by match filtering these masked maps. Once filtered, groups of map pixels with $S/N > 4.8$ are identified.

Finite map pixel size affects the measured flux densities, since it is rare that a source falls exactly in the center of a pixel, leading to a systematic negative bias and increased scatter. We increase the map pixel resolution by a factor of 16 in a region 0.03° on a side around each source through Fourier space zero-padding (e.g., Press et al. 2007) to allow for a more precise determination of the source peak location, and therefore its flux density. Properly centering the detection has the effect of boosting the S/N of typical




Figure 3. Filtered 148 GHz (top) and 218 GHz (bottom) submaps. The data have been match-filtered such that the grey-scale is in units of flux density (mJy) with white (black) corresponding to -10 mJy (30 mJy). Insets show the flux density distribution across the data as a grey histogram that has a standard deviation of 2.57 mJy (3.78 mJy) for 148 GHz (218 GHz). Several sources, both synchrotron and dust dominated, marked as circle and square outlines respectively, are apparent as black beam-sized flux excesses. The white extended object in the 148 GHz map, marked by a diamond, is the Sunyaev Zel’dovich effect decrement from Abell 3128 NE (ACT-CL J0330-5228).

$S/N=4.8$ sources to ≥ 5 . Lastly, to account for the convolution of sources with the map pixel, which acts as a low pass filter (the square pixels become a sinc function in Fourier space), we deconvolve the map pixel window function in the higher resolution map. We find that this method reduces systematic errors associated with pixelization to below 1%. Purity tests (Section 4.3) reveal a significant number of false detections below S/N of 5.0, due to local noise and striping artifacts. Therefore we impose a $S/N = 5.0$ threshold on sources detected in each map given no prior knowledge of source location from the other frequency.

Once a $S/N > 5.0$ catalog has been generated for each of the filtered 148 GHz and 218 GHz maps, we look for sources that have been identified at one frequency but not the other. The flux density for the source at the second frequency is then measured from the filtered map for the second frequency. We use the same prescription as for the flux measurement method just described, using a submap at the second frequency centered on the location of the detection as determined from the map at the first frequency.

4 THE 148 GHz AND 218 GHz CATALOG

The ACT-detected source catalog is given in Table 4. We find 169 sources selected at 148 GHz with $S/N > 5$, spanning two decades in flux density, from 14 to 1700 mJy. The 218 GHz map independently yielded 133 sources with $S/N > 5$. The combination of these two independent source lists gives a total count of 191, with 110 galaxies detected with $S/N > 5$ at both frequencies.

The catalog provides the IAU name, celestial coordinates (J2000), S/N and flux density estimation of each 148 and 218 GHz ACT-detected source. Raw flux densities are estimated directly from the map as described in Section 3. Deboosted flux densities as derived according to Section 4.2 are given with associated 68% confidence intervals. The raw and deboosted spectral indices,

$\alpha_{148-218}$, between 148 and 218 GHz for each source, are provided. The classification of sources as dust or synchrotron-dominated is based on the $\alpha_{148-218}$ spectral index criterion described in Section 4.2. If the source was cross-identified with an AT20G catalog source (see Section 5), the AT20G source ID is given. If, instead, the 20 GHz source flux density was measured during the November 2010 follow-up campaign (Section 2.3), the 20 GHz ID name has an asterisk next to it.

Correlating 148 GHz flux densities between this catalog and the catalog given in M11, we find an average agreement at the 2% level, with larger scatter for individual sources. This level of consistency is expected given changes in calibration, map-making, beam profile estimates, and deboosting procedure made for this updated study.

The reported flux densities are the average over approximately two months of observation for each source, many of which are AGN-driven radio galaxies and thus likely to have varied in that time. For example, of the three bright sources cross-identified with *Planck*, in the year between ACT and *Planck* observations one source varied by 20% in flux density whereas another did not vary at all within errors. However, our simultaneous multifrequency observations allow for a consistent internal spectral characterization between ACT bands. In a future study, multiple years of data will be used to quantify the effects of variability on individual source spectra.

The following sections provide details and context for the catalog values.

4.1 Astrometric Accuracy

Radio interferometers can achieve very precise positional accuracy for sources, so ACT-selected sources cross-identified with a robust radio catalog give a good measure of the positional accuracy of the ACT source detections. The AT20G catalog covers the Southern sky, and through pointing checks against Very Long Baseline Interferometer (VLBI) measurements of International Celestial Reference Frame calibrators, the positional accuracy of AT20G is shown to be accurate to better than $1''$ (Murphy, T. et al. 2010). We exclude nearby extended/resolved sources from this analysis, determined from cross-identification of our sources with currently available catalogs, as distant point-like sources will present a clearer picture of the overall accuracy of our pointing.

We compared the positions of the 34 ACT sources with $S/N_{148} > 16$ to positions of associated sources in the AT20G catalog using a search radius of half the ACT 148 GHz beam ($0.7''$). Figure 4 shows the offsets in location between the AT20G right ascension (RA) and declination (Dec) and the ACT-derived positions for the sources as they were detected in the 148 GHz (black points) and 218 GHz (blue points) maps. The results are also summarized in Table 1. The large overlaid errorbars are centered on the mean offsets, and extend as far as the rms of the offsets. For sources with lower S/N , the ACT location rms with respect to AT20G position becomes inflated by the effect of noise in the maps. Therefore the catalog uses coordinates derived from the 148 GHz map since there is a higher level of noise present in the 218 GHz map.

4.2 Flux Density Accuracy

Flux densities derived from the ACT maps have systematic uncertainties arising from five effects: the overall calibration uncertainty, the mapmaker, errors in the assumed source profile, error in the

Table 1. Astrometric Pointing Accuracy

	148 GHz	218 GHz
Mean RA Offset (")	0.6 ± 0.4	0.5 ± 0.6
Mean Dec Offset (")	-0.4 ± 0.3	0.0 ± 0.6
RMS in RA Offset (")	2.1	3.5
RMS in Dec Offset (")	1.8	3.4




Figure 4. Astrometric accuracy of the ACT source detections. The 34 filled black circles are the positional offsets of ACT sources with $S/N_{148} > 16$ from counterparts in the AT20G catalog as calculated from the 148 GHz map. The blue points are for the same sources, but using positions derived from the 218 GHz map, which has more noise. The large errorbars are centered on the mean offsets, and extend as far as the rms of the offsets. See Table 1.

assumed source spectrum, and flux boosting of lower significance candidates. The first four potential sources of flux density error were discussed in Section 2. Here we will discuss the last source of flux density error.

The differential counts of the sources selected in our sample fall steeply with increasing flux (Table 2, Figure 6). With no prior information about the source flux, the most likely scenario is that the measured flux is the sum of a dimmer intrinsic flux and a positive noise fluctuation. We use the two-band Bayesian method developed in Crawford et al. (2010), and report the 16, 50, and 84 percentiles (68% enclosed, equivalent to 1σ) of the posterior flux and spectral index distributions. For the source count priors in this calculation we use the sum of the models of de Zotti et al. (2010) for radio sources and Béthermin et al. (2011) for dust-dominated sources. Following Vieira, J. D. et al. (2010), we take a flat prior on the spectral index between -3 and 5 , consistent with the expected range for our populations. The two-band likelihood includes negligible correlation between bands and is consistent with background astrophysical emission rather than correlated atmospheric emission. The source populations in 148 and 218 GHz naturally split into sources having their emission dominated by synchrotron

(centered on $\alpha = -0.6$) or thermal dust (centered on $\alpha = 3.7$; see Figure 7, bottom panel). We use the threshold spectral index $\alpha = 1.66$ (Vieira, J. D. et al. 2010) to divide these populations in terms of their posterior spectral index populations, with $P(\alpha > 1.66) > 0.5$ classified as dusty and $P(\alpha > 1.66) < 0.5$ as synchrotron dominated. The classification is robust within $\alpha \pm 0.5$ of this threshold.

In addition to the bias from the steepness of the population, we also treat the fact that the source finder locates the maximum flux along RA and Dec, which provides an extra two degrees of freedom. This can be corrected by finding the flux after two degrees of freedom are subtracted from the detection significance, following Vanderlinde, K. et al. (2010). Note that this departs from the treatment in Crawford et al. (2010) and Vieira, J. D. et al. (2010). Calibration uncertainty is included in all quoted fluxes, and is assumed to be uncorrelated between bands.

4.3 Purity and Completeness

The number of false detections at each frequency was estimated by running the detection algorithm on an inverted (negative temperature) map in which the sources, and for 148 GHz (the Sunyaev-Zel'dovich effect decrement) all ACT-detected and optically confirmed galaxy clusters, were masked. With this approach, no spurious detections are found in the 148 GHz data down to a S/N of 5. In the 218 GHz data, four spurious detections at $S/N \leq 6$ were observed with raw flux densities in the 25–31 mJy range (Table 2).

The purity simulations appear to be consistent with our findings from source cross-identification (Section 5). At 148 GHz, all but one 5σ source is cross-identified with other catalogs, and at 218 GHz seven 5σ sources are not cross-identified, although we expect some of these to be real DSFGs.

In order to estimate completeness of the catalog for each band, we added 100 synthetic sources of a single flux density to the 455 deg^2 region of the ACT maps used in this study. The map filtering and source detection algorithm (Section 3) were run, and the resulting catalog checked for inclusion of the input sources. This procedure was repeated in each band in intervals of 10 mJy in synthetic source flux density in the range 10–100 mJy. The completeness at each flux density was interpolated to estimate the completeness in flux density bins given in Table 2.

5 COMPARISON TO OTHER CATALOGS

To further characterize ACT sources, we consider cross-identifications with other catalogs. Given the number of ACT-detected sources and the ACT beam size, around 1 out of 10,000 sources randomly placed in the 455 deg^2 area considered here would coincide with an ACT source. Figure 5 shows the ACT sources located in flux density space, and cross-identifications from some overlapping catalogs with extragalactic source detections in the southern hemisphere discussed in this section. The synchrotron-dominated sources occupy points near the power law $S_\nu \propto \nu^\alpha$ with spectral index $\alpha = -0.6$. Dust-dominated sources will follow a line closer to $\alpha = 3.7$, and are generally associated with sources detected with $S/N > 5$ at 218 GHz only.

5.1 General Statistics of Identifications

Of our 191 sources, 174 are identified with sources in the 0.84 GHz Sydney University Molonglo Sky Survey (SUMSS; Mauch et al. 2003), and 122 of those cross-identified sources also belong to the

Table 2. Number Counts, Purity, and Completeness^a.

Flux Range (Jy)	148 GHz			218 GHz			N_{sync}^b	N_{dust}^c
	N	Purity	Completeness	N	Purity	Completeness		
0.015 – 0.02	23	100.0±0.0%	47.1±3.2%	8	100.0±0.0%	10.7±2.8%	22	0
0.02 – 0.03	47	100.0±0.0%	75.3±3.4%	42	92.9±4.8%	34.2±3.5%	46	10
0.03 – 0.05	48	100.0±0.0%	96.8±1.9%	43	97.7±2.3%	80.2±3.0%	48	10
0.05 – 0.09	20	100.0±0.0%	99.6±0.6%	21	100.0±0.0%	98.7±0.9%	20	1
0.09 – 0.17	13	100.0±0.0%	100.0±0.0%	12	100.0±0.0%	100.0±0.0%	13	2
0.17 – 0.33	4	100.0±0.0%	100.0±0.0%	2	100.0±0.0%	100.0±0.0%	4	0
0.33 – 0.65	2	100.0±0.0%	100.0±0.0%	1	100.0±0.0%	100.0±0.0%	2	0
0.65 – 1.39	1	100.0±0.0%	100.0±0.0%	1	100.0±0.0%	100.0±0.0%	1	0
1.39 – 2.87	1	100.0±0.0%	100.0±0.0%	1	100.0±0.0%	100.0±0.0%	1	0

^aThe number of sources at each frequency with S/N=5 over 455 deg². Errors on number count are simply

Poisson. See Figure 6 for a graph of purity/completeness-corrected differential source counts.

^bCounts of synchrotron-dominated sources are taken relative to the 148 GHz flux density.

^cCounts of dust-dominated sources are binned according to their 218 GHz flux density.




Figure 5. The ACT sources cross-identified in flux density space. Sources have been selected with S/N > 5.0 in at least one frequency band (dotted lines). See text for a more thorough description of cross-identification statistics with other catalogs. The diagonal lines follow the power law $S_\nu \propto \nu^\alpha$ with $\alpha = 1.66$ for the dashed line, distinguishing the $\alpha = -0.6$ (dash-dot) synchrotron-dominated source population, and the $\alpha = 3.7$ (dash-dot-dot) dust-dominated source population.

4.85 GHz Parkes-MIT-NRAO (PMN) survey (Wright et al. 1994) radio catalog. Fourteen sources were cross-identified with the Infrared Astronomical Satellite (IRAS; Helou et al. 1988) at 12–100 μm . Sources were also identified with one or both of the Australian Telescope 20 GHz survey (AT20G; Murphy, T. et al. 2010) and the 1.4/2.0 mm SPT (Vieira, J. D. et al. 2010) catalogs. Two of the remaining unmatched sources were observed with the Australian Telescope Compact Array (ATCA) during a November 2010 follow-up campaign. This leaves 6 sources with no identification in any of the available catalogs, all but one of which have dust-dominated spectra. The single unidentified source with a syn-

chrotron spectrum, ACT-S J023600-530237, falls near our S/N=5 threshold and could be a false detection.

A NASA/IPAC Extragalactic Database (NED) search using a radius of 2' cross-identified 3 sources from the ACT catalog with galaxy clusters: Abell S0250, Abell 3391 and Sérsic 037/01. Increasing the search radius to 3' adds Abell 3128 and Abell 3395 to this list. Simulations of the microwave sky suggest that only $\approx 3\%$ of galaxy clusters have their 148 GHz signal contaminated at 20% or more (Sehgal et al. 2010).

5.2 Matches to AT20G

The AT20G survey, carried out from 2004 to 2008 at 20 GHz with follow-up at 5 and 8 GHz, covered 6.1 sr of the Southern sky to a flux density limit of 40 mJy at 20 GHz (Murphy, T. et al. 2010). Of our sources, 115 match sources in the AT20G catalog, listed in Table 4, all of which we classify as synchrotron-dominated. Given the AT20G completeness limit and the predominance of flat spectra for our 148 GHz-selected sources (Section 7), faint ACT sources may not have matches in AT20G. We received time on the Australia Telescope Compact Array to measure flux densities for sources in the M11 catalog that did not appear in AT20G, the results of which are described in Sections 2.3 and 7.

5.3 Matches to the Planck Compact Source Catalogs

The *Planck* satellite team has released two sets of all-sky catalogs in nine frequency channels, including two very close to those used by ACT. The first set of catalogs made up the Early Release Compact Source Catalog (ERCSC; Planck Collaboration XIII 2011). Very recently, a deeper catalog based on 1.6 complete *Planck* surveys has been released, the Planck Catalog of Compact Sources (PCCS; Planck Collaboration XXVIII 2013). At 143 and 217 GHz, the *Planck* beam sizes are approximately 7' and 5' respectively. Like WMAP, *Planck* makes great gains by observing the entire sky simultaneously at several frequencies, thus capturing all bright sources. On the other hand, due to their large beam sizes, these experiments detect *only* the brightest sources. In the case of the PCCS, the detection threshold at 143 and 217 GHz is ≈ 400 mJy; at lower flux densities the catalog completeness drops well below 80%. In

the area treated in this study, *Planck* detects 3 of the brightest ACT sources.

For a more complete comparison with the PCCS, we examined all bright ACT sources, combining the Equatorial and Southern regions mapped by ACT. Almost 50 matches were found in this wider comparison. At 148 GHz, the agreement of ACT flux densities and the PCCS flux densities, properly color-corrected and (slightly) extrapolated to match ACT's central frequency, is excellent, at the 1% level, despite scatter introduced by source variability. At 218 GHz, source variability plays at least as large a role, and PCCS flux densities are $\approx 3\%$ higher than ACT's. If we remove a couple of the most variable sources, however, the agreement of the 218 GHz flux density scales improves to $\approx 1\%$. At both frequencies, *Planck* fluxes are on average higher. Agreement at the $\approx 1\%$ level, however, is well within the expected uncertainty in the ACT flux densities (Section 2).

5.4 Matches to the SPT Catalog

The SPT study of compact sources (Vieira, J. D. et al. 2010) was based on observations of a square patch of sky of 87 deg^2 centered at 05^h right ascension, having only fractional ACT overlap. Nevertheless, 2,304 of the 3,496 SPT candidate sources (those with $S/N > 3$ and flux densities $> 4.4 \text{ mJy}$) fall within the ACT survey region. We find 32 cross-identifications with ACT sources when we search the SPT 1.4 mm (220 GHz) and 2.0 mm (145 GHz) catalogs. Twenty-five of these were 148 GHz-selected and categorized in Vieira, J. D. et al. (2010) as synchrotron-dominated. The remaining seven were detected by ACT with $S/N > 5$ only at 218 GHz; three of the seven also have SUMSS or IRAS cross-identifications. To make a direct comparison of the flux densities of ACT-SPT cross-identified sources, we used our derived flux densities, but without the final deboosting steps, as they were not incorporated by the SPT analysis (Crawford et al. 2010). We find an overall offset at the $7 \pm 5\%$ level at 148 GHz and $3 \pm 7\%$ at 218 GHz, with ACT flux densities typically higher than those of SPT. In future studies that include variability, the large errors on this comparison may be decreased by excluding sources for which there is known variability.

Recently, using spectroscopic follow-up observations with the Atacama Large Millimeter/submillimeter Array (ALMA), Vieira, J. D. et al. (2013) and Weiß, A. et al. (2013), derived robust redshifts for about 18 of the SPT-detected dust-dominated sources from a larger sky area than in Vieira, J. D. et al. (2010). One of these sources, found to be at redshift 5.66, matches ACT-S J034640-520505 which otherwise had no cross-identification. A second previously unmatched source, ACT-S J002707-500713, was imaged with ALMA at $870 \mu\text{m}$, is resolved at the $0.5''$ scale, likely due to gravitational lensing, but awaits spectroscopic follow-up. Four other ACT dust-dominated sources that were cross-identified with the Vieira, J. D. et al. (2010) catalog now also have robust redshifts. ACT-S J055139-505800 is at redshift 2.123, ACT-S J053250-504709 is at redshift 3.399, ACT-S J052903-543650 is a source at redshift 3.369, with a lens at redshift 0.140, and ACT-S J053817-503058 has a redshift of 2.782, and a lens at a redshift of 0.404. It is likely that the few ACT dust-dominated sources which lie outside the SPT footprint are also lensed DSFGs.

5.5 Matches to the IRAS Catalog

Given the ACT beam size, a normal galaxy will be unresolved at redshifts $z \geq 0.05$ or distances greater than 200 Mpc. Conse-

quently, only very nearby objects appear extended in our maps. The IRAS source population consists primarily of local ($z \ll 1$) dust-dominated ULIRGs; we thus expect our resolved sources to coincide with IRAS sources. We find 14 sources cross-identified with IRAS sources, out of a possible 829 IRAS sources that lie within the ACT survey area treated here. Two of these have synchrotron-dominated spectra, both of which are nearby galaxies. The twelve that show spectra dominated by dust re-emission are Galactic sources, in the Magellanic clouds, or are known nearby star-forming galaxies. For example, for two local resolved galaxies particularly bright at ACT frequencies, NGC1566 (ACT-S J041959-545622), a Seyfert two-arm spiral, and IC1954 (ACT-S J033133-515352), a late-type spiral with a short central bar, ACT observes a higher 218 GHz flux density than at 148 GHz or 20 GHz, confirming the dust-dominated nature of their spectra.

5.6 Dust-Dominated ACT Sources

Of the 24 ACT dust-dominated sources in the catalog presented here, 19 of these are cross-identified with either IRAS (12 sources) or SPT (9 sources). Of these 19 cross-identified sources, some have multiple cross-identifications beyond the IRAS or SPT match, including with the SUMSS catalog. The remaining 5 have no matching counterpart, and signal-to-noise at 218 GHz in the range $5.27 < S/N < 6.35$. Purity tests suggest that a couple of these are spurious, but given that three $S/N \approx 6$ sources are cross-identified with SPT, some of these sources are likely real detections, high redshift galaxies lensed by intervening structure.

6 SOURCE COUNTS

The completeness-corrected differential number counts for ACT sources based on the data in Table 2 are plotted in Figure 6. For comparison, completeness-corrected number counts from Vieira, J. D. et al. (2010) and Planck Collaboration XIII (2011) are plotted as well. The ACT data fill in the flux density gap between the SPT and *Planck* catalogs at these frequencies, caused by the differences between experiments in sky coverage and sensitivity to point sources. The effect of calibration error is to shift the flux bins by $\pm 2\%$ ($\pm 2.4\%$) at 148 GHz (218 GHz; Section 2).

The combined ACT, SPT and *Planck* total counts at 148 GHz and synchrotron-dominated source counts at 218 GHz are best fit by the recently developed Tucci et al. (2011) C2Ex radio source model. This model divides the \sim flat- or steep-spectrum blazar population into BL Lac objects, with a region close to the AGN core dominating the observed emission ($\leq 0.3 \text{ pc}$), and flat spectrum radio quasars (FSRQs) with a break frequency indicating that the source of emission arises from an emitting region further from the AGN core (0.3 to 10 pc). The de Zotti et al. (2005) model for radio source counts, while consistent with the counts below 0.1 Jy, over-predicts the counts at higher flux densities. *Planck* analysis finds that the de Zotti et al. (2005) model is consistent with their counts at frequencies up to 100 GHz, but over-predicts the counts at higher frequencies in the flux density region of $\approx 1 \text{ Jy}$, though they begin to suffer from incompleteness below $\approx 1 \text{ Jy}$. Our few $\approx 1 \text{ Jy}$ brightest sources appear to be consistent with this finding but lend little statistical significance.

As well as the models for counts of radio sources, the source counts predicted for dusty starburst galaxies from Toffolatti et al. (1998) are shown in Figure 6. The brightest infrared sources in this model are 10 mJy. Given that all sources in the ACT catalog




Figure 6. Differential number counts of ACT-selected sources. Derived from Table 2 and corrected for completeness, the ACT differential source counts are plotted together with models of radio and infrared source populations. The Planck Collaboration VII (2013) data points and the SPT data points of Vieira, J. D. et al. (2010) are also plotted, both of which are consistent with ACT counts. The ACT and Planck points have Poissonian errors (1σ), whereas the SPT points include measurement and independent calibration errors. The fIR models on this scale predicts number counts too low to be seen at 148 GHz. The ACT data are consistent with being dominated by radio sources at both frequencies.

have flux densities greater than 10 mJy, these models predict that the 148 GHz- and 218 GHz-selected ACT catalogs should have few or no detected infrared sources. However, we find 24 sources that have dust-dominated spectra with flux densities above 20 mJy, and Vieira, J. D. et al. (2010) find 36 dust-dominated sources with flux densities above 15 mJy. Furthermore, Toffolatti et al. (1998) does not incorporate the effects of lensing on observed number counts.

Cai et al. (2013) modelled dust-dominated galaxies together with lensed high-redshift dust-dominated galaxies. However, Figure 6 indicates that at the ACT frequencies, this model underpredicts the counts, an effect already observed at lower frequencies. This is due to a deficiency in the assumed galactic spectral energy distributions used [Cai, private communication]. The Lima et al. (2010) model shown is for dust-dominated galaxies at high redshift (i.e. excluding local, IRAS cross-identified, dusty galaxies), and also incorporates gravitational lensing.

7 SOURCE SPECTRA

Source spectral energy distributions (SEDs) can be used to differentiate source types by their dominant emission mechanisms. Assuming the commonly used simple power law model $S(\nu) \propto \nu^\alpha$, a negative α is indicative of sources dominated by synchrotron emission, such as radio galaxies. Sources with free-free emission dominating will have an index close to 0. The high-redshift SMG population will have spectra dominated by re-emission of their prodigious optical and UV flux by the surrounding dust in a greybody spectrum, with indices expected to be greater than 2 and more typically 3–4.

We can divide the ACT source population according to several broad spectral groups: classical steep (and steepening) spectrum sources, sources that peak within the frequency range under consideration, and sources that show flat, rising or upturned spectra. Though the ACT catalog is dominated by synchrotron-dominated sources which have variable flux densities that observe no strict periodicity, as an ensemble, such a study may give rise to insights about the average spectral behavior of the galaxy populations.

Table 3 summarizes the median spectral indices between pairs of frequencies for various subsets of the ACT data. In obtaining the spectral indices we compare the deboosted flux densities from ACT with the raw flux densities from AT20G. The whole sample includes data from the 20 GHz ATCA follow-up observations taken in November 2010 (Section 2.3).

Figure 7 shows the 5–20 GHz, 20–148 GHz and 148–218 GHz color-magnitude diagrams. The radio-selected AGN (black points) are predominantly characterized by steepening of the spectra, with only a few characterized by extremely flat or inverted spectra. Magenta points, clearly showing rising spectra between 148 and 218 GHz, denote sources ACT-classified as spectrally dust dominated. They do not show up in the top two plots, as these sources would have flux densities falling below the detection threshold of the AT20G catalog.

The average spectral indices between AT20G and ACT frequencies indicates an underlying source population made up of Flat-Spectrum Radio Quasars (FSRQ), a type of blazar, with AGN jet pointed along our line of sight (de Zotti et al. 2010). Ejected material flows through several shocked regions in the jet which locally enhance the radiation (Marscher & Gear 1985;

Table 3. Median Spectral Indices.

Spectral Index	Synchrotron (all) ^a	Synchrotron ($S_{148} > 50$ mJy)	Synchrotron ($S_{148} < 50$ mJy)	Dust-dominated
α_{5-20}	$-0.15 \pm 0.39 (\pm 0.04)$	$-0.07 \pm 0.37 (\pm 0.06)$	$-0.21 \pm 0.38 (\pm 0.04)$...
α_{20-148}	$-0.42 \pm 0.32 (\pm 0.03)$	$-0.36 \pm 0.26 (\pm 0.04)$	$-0.43 \pm 0.35 (\pm 0.03)$...
$\alpha_{148-218}$	$-0.55 \pm 0.62 (\pm 0.05)$	$-0.60 \pm 0.23 (\pm 0.04)$	$-0.51 \pm 0.70 (\pm 0.06)$	$3.7 \pm 0.70 (\pm 0.14)$

^aQuoted errors are the 68% confidence levels of the distribution and, in parentheses, the standard deviation of the mean.




Figure 7. Color-magnitude diagrams comparing the 5–20 GHz (top), 20–148 GHz (middle), and 148–218 GHz (bottom) spectral indices for ACT-ATCA cross-identified sources. The synchrotron-dominated radio galaxy population is dominated by sources which have consistently falling SEDs towards higher frequencies.

Valtaoja et al. 1992). The observed spectral flatness is the superposition of many components with different turnover frequencies. At frequencies greater than approximately 100 GHz, however, Marriage, T. A. et al. (2011), Vieira, J. D. et al. (2010), and Planck Collaboration XIII (2011) observe a steepening of the spectrum ($\alpha_{148-218} \approx -0.6$) that until now had not been conclusively shown (Tucci et al. 2011). This change is possibly due to electron energy losses in the jet (“electron ageing”) or the transition to the optically thin regime in the extended radio lobes. The underlying physical mechanisms have been contested for more than a decade and remain the subject of ongoing study, (e.g., Nieppola et al. 2008; Ghisellini & Tavecchio 2008; Sambruna et al. 2010).

All but one of the synchrotron-dominated ACT detections with flux density > 50 mJy have cross-identifications in AT20G. Below this flux density, the mean spectral indices of the population of ACT-AT20G cross-identified sources are biased toward the neg-




Figure 8. Radio color-magnitude diagram using 20–148 GHz spectral indices for ACT-AT20G cross-identified sources. The data are divided between flux densities at 50 mJy at 148 GHz. The low flux sample was incomplete and suffered from selection bias that favored sources with more negative spectral indices. Data from the ATCA follow-up study for the M11 ACT 148 GHz sources not cross-identified with AT20G are shown in blue. The high flux sample, denoted with black points, has a median spectral index -0.37 . Prior to ATCA follow-up observations, the lower flux ACT sources cross-identified with the AT20G catalog (grey points) had a median spectral index of -0.52 . Including the follow-up data, the more complete lower flux sample has a median spectral index of -0.43 .

ative by the incompleteness of the AT20G catalog below 100 mJy at 5 GHz and below 40 mJy at 20 GHz. It is therefore illustrative to further divide this subpopulation according to 148 GHz flux density.

Figure 8, a radio color-magnitude diagram, plots spectral index α_{20-148} against 148 GHz source flux density. Black and grey points identify the M11 sources that were cross-identified with the AT20G catalog. Black points are for sources with flux densities > 50 mJy, which represented a complete sample. Grey points denote the fainter, incomplete sample. The blue points were obtained by calculating the spectral index for the previously unmatched, lower flux density subsample, followed up with ATCA (see Table 4 for AT20G source IDs with an asterisk). The population represented by the blue points fills in the picture remarkably for the fainter (below 50 mJy flux density at 148 GHz, below 40 mJy at 20 GHz) population, especially in the region with spectral index of approximately zero.

The unbiased $S_{148} > 50$ mJy sample has a 20–148 GHz me-

dian spectral index of $-0.37 \pm 0.24 (\pm 0.04)$. For the $S_{148} < 50$ mJy subsample prior to follow-up with ATCA (grey points only), the 20–148 GHz spectral index was $\alpha_{20-148} = -0.52 \pm 0.23 (\pm 0.03)$. However, with the addition of the synchrotron-classified ACT-selected sources with 2010 ATCA follow-up data (grey and blue points), the full sample has 20–148 GHz spectral index $\alpha_{20-148} = -0.43 \pm 0.30 (\pm 0.03)$.

8 CONCLUSION

We have described the extragalactic source population at 148 and 218 GHz in a 455 deg^2 region of the ACT 2008 *Southern strip*, centered on declination -52.5° . This updates the ACT 148 GHz catalog to use the data released with Dünner, R. et al. (2013), and extends the results in Marriage, T. A. et al. (2011) to two bands and by a more localized treatment of noise which in turn yields a higher S/N. We detect 191 sources above $S/N=5$ in at least one of the ACT 148 and 218 GHz frequency bands, spanning flux densities 14–1700 mJy (Table 4) and redshifts 0.005–5.6. The catalog is estimated to be 100% pure and 96.8% complete above 30 mJy at 148 GHz, and 97.7% pure and 80.2% complete above 30 mJy at 218 GHz. We have confirmed flux recovery of the pipeline, and jointly deboosted the flux densities in both bands.

The multifrequency nature of our observations allows for internal classification of sources into three broad classes of sources based on their spectra: synchrotron-dominated sources (the vast majority of which are cross-identified with radio catalogs), low-redshift dust-dominated sources with IRAS counterparts (typically ULIRGs), and dust-dominated sources also observed by SPT or with no cross-identification either shown or assumed to be high redshift star-forming galaxies. This last class, only recently observed at millimeter wavelengths, are part of the progenitor population of massive, modern-day, elliptical galaxies, background SMGs whose flux has been magnified through gravitational lensing by a foreground galaxy or galaxy group, an interpretation that is being validated with follow-up observations.

Comparison to other catalogs shows that 97% of ACT-detected sources correspond to sources detected at lower or higher frequencies. The 148 GHz source counts are fit reasonably well by the C2Ex radio model of Tucci et al. (2011), the most current model for radio sources. The ACT data support the case for a spectral steepening toward higher frequencies above 100 GHz for AGN, by the analysis of the average spectral indices derived from the combined AT20G and ACT datasets. The ACT dust-dominated source population has a median spectral index, $\alpha_{148-218}$, of $3.7^{+0.62}_{-0.86}$. Properly linking these sources into the broader context of galaxy formation and evolution is of cosmological interest, and a goal of future work.

The analysis presented here uses only the 2008 data of ACT’s *Southern strip*, representing only $1/6^{\text{th}}$ of the data ACT obtained between 2007 and 2010. In future work, we will extend our analysis of the source population to include the full dataset integrating both the southern and equatorial regions observed by ACT. The ACT *Equatorial strip* overlaps with deep Sloan Digital Sky Survey Stripe 82 observations (Annis et al. 2011). Thus as well as increasing the sky coverage and number counts for the ACT sources, future work (Gralla et al. in prep.) will include joint analyses with optical data.

ACKNOWLEDGMENTS

This work was supported by the U.S. National Science Foundation through awards AST-0408698 and AST-0965625 for the ACT project, and PHY-0355328, PHY-0855887, PHY-1214379, AST-0707731 and PIRE-0507768 (award number OISE-0530095). Funding was also provided by Princeton University, the University of Pennsylvania, and a Canada Foundation for Innovation (CFI) award to UBC. ES acknowledges support by NSF Physics Frontier Center grant PHY-0114422 to the Kavli Institute of Cosmological Physics. The PIRE program enabled this research program through exchanges between Chile, South Africa, Spain and the US. Computations were performed on the GPC supercomputer at the SciNet HPC Consortium. SciNet is funded by: SciNet is funded by the Canada Foundation for Innovation (CFI) under the auspices of Compute Canada, the Government of Ontario, the Ontario Research Fund – Research Excellence, and the University of Toronto. Data acquisition electronics were developed with assistance from the CFI. ACT operates in the Parque Astronómico Atacama in northern Chile under the auspices of the Comisión Nacional de Investigación Científica y Tecnológica de Chile (CONICYT).

We thank the staff at the Australia Telescope Compact Array site, Narrabri (NSW), for the valuable support they provide in running the telescope. ATCA is part of the Australia Telescope National Facility which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO.

This research made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

ACT data products are publicly accessible through LAMBDA (<http://lambda.gsfc.nasa.gov/>) and the ACT website (<http://www.physics.princeton.edu/act/>).

REFERENCES

- Amblard, A. et al. 2010, *A&A*, 518, L9+
- Annis J., Soares-Santos M., Strauss M. A., Becker A. C., Dodelson S., Fan X., Gunn J. E., Hao J., Ivezić Z., Jester S., Jiang L., Johnston D. E., Kubo J. M., Lampeitl H., Lin H., Lupton R. H., Miknaitis G., Seo H.-J., Simet M., Yanny B., 2011, ArXiv e-prints
- Austermann, J. E. et al. 2010, *MNRAS*, 401, 160
- Béthermin M., Dole H., Lagache G., Le Borgne D., Penin A., 2011, *A&A*, 529, A4
- Blain A. W., Smail I., Ivison R. J., Kneib J., Frayer D. T., 2002, *Physics Reports*, 369, 111
- Bonavera L., Massardi M., Bonaldi A., González-Nuevo J., de Zotti G., Ekers R. D., 2011, *MNRAS*, 416, 559
- Cai Z.-Y., Lapi A., Xia J.-Q., De Zotti G., Negrello M., Gruppioni C., Rigby E., Castex G., Delabrouille J., Danese L., 2013, *ApJ*, 768, 21
- Carlstrom, J. E. et al. 2011, *PASP*, 123, 568
- Crawford T. M., Switzer E. R., Holzapfel W. L., Reichardt C. L., Marrone D. P., Vieira J. D., 2010, *ApJ*, 718, 513
- de Zotti G., Massardi M., Negrello M., Wall J., 2010, *A&AR*, 18, 1
- de Zotti G., Ricci R., Mesa D., Silva L., Mazzotta P., Toffolatti L., González-Nuevo J., 2005, *A&A*, 431, 893
- Devereux N. A., Young J. S., 1990, *ApJ*, 359, 42
- Devlin, M. J. et al. 2009, *Nature*, 458, 737

- Dole H., Lagache G., Puget J., Caputi K. I., Fernández-Conde N., Le Floc'h E., Papovich C., Pérez-González P. G., Rieke G. H., Blaylock M., 2006, *A&A*, 451, 417
- Draine B. T., 2003, *ARA&A*, 41, 241
- Dünner, R. et al. 2013, *ApJ*, 762, 10
- Ghisellini G., Tavecchio F., 2008, *MNRAS*, 387, 1669
- Goldin A. B., Kowitt M. S., Cheng E. S., Cottingham D. A., Fixsen D. J., Inman C. A., Meyer S. S., Puchalla J. L., Ruhl J. E., Silverberg R. F., 1997, *ApJL*, 488, L161
- Griffin M. J., Orton G. S., 1993, *Icarus*, 105, 537
- Hajian, A. et al. 2011, *ApJ*, 740, 86
- Hasselfield, M. et al. 2013, *ArXiv e-prints*
- Helou G., Khan I. R., Malek L., Boehmer L., 1988, *ApJS*, 68, 151
- Hezaveh, Y. D. et al. 2013, *ApJ*, 767, 132
- Hincks, A. D. et al. 2010, *ApJS*, 191, 423
- Holland W. S., Robson E. I., Gear W. K., Cunningham C. R., Lightfoot J. F., Jenness T., Ivison R. J., Stevens J. A., Ade P. A. R., Griffin M. J., Duncan W. D., Murphy J. A., Naylor D. A., 1999, *MNRAS*, 303, 659
- Holland, W. S. et al. 2013, *MNRAS*, 430, 2513
- Ikarashi, S. et al. 2011, *MNRAS*, 415, 3081
- Lamarre, J.-M. et al. 2010, *A&A*, 520, A9
- Lima M., Jain B., Devlin M., Aguirre J., 2010, *ApJL*, 717, L31
- Lupu, R. E. et al. 2012, *ApJ*, 757, 135
- Marriage, T. A. et al. 2011, *ApJ*, 731, 100
- Marscher A. P., Gear W. K., 1985, *ApJ*, 298, 114
- Massardi M., Bonaldi A., Bonavera L., López-Cañiego M., de Zotti G., Ekers R. D., 2011, *MNRAS*, pp 878–+
- Massardi, M. et al. 2011, *MNRAS*, 412, 318
- Mauch T., Murphy T., Buttery H. J., Curran J., Hunstead R. W., Piestrzynski B., Robertson J. G., Sadler E. M., 2003, *MNRAS*, 342, 1117
- Murphy, T. et al. 2010, *MNRAS*, 402, 2403
- Negrello M., Perrotta F., González-Nuevo J., Silva L., de Zotti G., Granato G. L., Baccigalupi C., Danese L., 2007, *MNRAS*, 377, 1557
- Negrello, M. et al. 2010, *Science*, 330, 800
- Nieppola E., Valtaoja E., Tornikoski M., Hovatta T., Kotiranta M., 2008, *A&A*, 488, 867
- Pascale, E. et al. 2008, *ApJ*, 681, 400
- Pilbratt G. L., Riedinger J. R., Passvogel T., Crone G., Doyle D., Gageur U., Heras A. M., Jewell C., Metcalfe L., Ott S., Schmidt M., 2010, *A&A*, 518, L1
- Planck Collaboration VII 2013, *A&A*, 550, A133
- Planck Collaboration XIII 2011, *A&A*, 536, A13
- Planck Collaboration XV 2011, *A&A*, 536, A15
- Planck Collaboration XVI 2011, *A&A*, 536, A16
- Planck Collaboration XXVIII 2013, *ArXiv e-prints*
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 2007, *Numerical recipes in C (3rd ed.): the art of scientific computing*. Cambridge University Press, New York, NY, USA
- Rex, M. et al. 2010, *A&A*, 518, L13
- Sambruna R. M., Donato D., Ajello M., Maraschi L., Tueller J., Baumgartner W., Skinner G., Markwardt C., Barthelmy S., Gehrels N., Mushotzky R. F., 2010, *ApJ*, 710, 24
- Sault R. J., Teuben P. J., Wright M. C. H., 1995, in R. A. Shaw, H. E. Payne, & J. J. E. Hayes ed., *Astronomical Data Analysis Software and Systems IV Vol. 77 of Astronomical Society of the Pacific Conference Series, A Retrospective View of MIRIAD*. pp 433–+
- Scott, K. S. et al. 2011, *ApJ*, 733, 29
- Sehgal N., Bode P., Das S., Hernandez-Monteagudo C., Hufferberger K., Lin Y., Ostriker J. P., Trac H., 2010, *ApJ*, 709, 920
- Siringo G., Kreysa E., Kovács A., Schuller F., Weiß A., Esch W., Gemünd H.-P., Jethava N., Lundershausen G., Colin A., Güsten R., Menten K. M., Beelen A., Bertoldi F., Beeman J. W., Haller E. E., 2009, *A&A*, 497, 945
- Swetz, D. S. et al. 2011, *ApJS*, 194, 41
- Tegmark M., 1997, *ApJL*, 480, L87+
- Thompson A. R., Moran J. M., Swenson G. W., 1986, *Interferometry and synthesis in radio astronomy*
- Toffolatti L., Argüeso Gomez F., de Zotti G., Mazzei P., Franceschini A., Danese L., Burigana C., 1998, *MNRAS*, 297, 117
- Tucci M., Toffolatti L., de Zotti G., Martínez-González E., 2011, *A&A*, 533, A57
- Valtaoja E., Terasranta H., Urpo S., Nesterov N. S., Lainela M., Valtonen M., 1992, *A&A*, 254, 71
- Vanderlinde, K. et al. 2010, *ApJ*, 722, 1180
- Vieira, J. D. et al. 2010, *ApJ*, 719, 763
- Vieira, J. D. et al. 2013, *Nature*, 495, 344
- Viero, M. P. et al. 2009, *ApJ*, 707, 1766
- Weiß, A. et al. 2009, *ApJ*, 707, 1201
- Weiß, A. et al. 2013, *ApJ*, 767, 88
- Werner, M. W. et al. 2004, *ApJS*, 154, 1
- Wiebe, D. V. et al. 2009, *ApJ*, 707, 1809
- Wilson G. W., Austermann J. E., Perera T. A., Scott K. S., Ade P. A. R., Bock J. J., Glenn J., Golwala S. R., Kim S., Kang Y., Lydon D., Maukopf P. D., Predmore C. R., Roberts C. M., Soucar K., Yun M. S., 2008, *MNRAS*, 386, 807
- Wilson, W. E. et al. 2011, *MNRAS*, 416, 832
- Wright A. E., Griffith M. R., Burke B. F., Ekers R. D., 1994, *ApJS*, 91, 111
- Wright, E. L. et al. 2009, *ApJS*, 180, 283

Table 4: ACT High Significance 148 GHz and 218 GHz Extragalactic Source Catalog

ID and Coordinates			148 GHz			218 GHz			Spectral Index			20 GHz	
ACT ID	RA (J2000) hh:mm:ss	Dec dd:mm:ss	S/N	S _m ^a (mJy)	S _{db} (mJy)	S/N	S _m ^b (mJy)	S _{db} (mJy)	α _m	α _{db}	Type	ATCA ID ^c	S _m (mJy)
ACT-S J001849-511506	00 : 18 : 49.7	-51 : 15 : 06.1	11.5	56.0	54.7 ^{+5.1} _{-5.1}	7.6	41.3	39.7 ^{+5.7} _{-5.6}	-0.8	-0.8 ^{+0.4} _{-0.5}	sync	J001849-511455	154.0±5.0
ACT-S J001853-492953	00 : 18 : 53.0	-49 : 29 : 53.6	7.3	49.8	46.9 ^{+7.1} _{-7.1}	4.9	44.8	40.7 ^{+9.5} _{-9.5}	-0.3	-0.4 ^{+0.7} _{-0.8}	sync
ACT-S J002233-515326	00 : 22 : 33.7	-51 : 53 : 26.6	5.4	21.7	20.0 ^{+4.1} _{-4.1}	5.5	25.3	21.7 ^{+5.1} _{-5.5}	0.4	0.2 ^{+0.8} _{-0.9}	sync	J002233-515316	77.0±9.0
ACT-S J002512-542744	00 : 25 : 12.5	-54 : 27 : 44.6	6.1	22.6	20.6 ^{+3.8} _{-3.9}	4.9	23.0	20.8 ^{+4.8} _{-4.8}	-0.0	0.0 ^{+0.8} _{-0.8}	sync	J002511-542739	59.0±3.0
ACT-S J002707-500713	00 : 27 : 07.3	-50 : 07 : 13.3	2.2	13.5	10.4 ^{+3.1} _{-3.3}	6.0	44.8	39.3 ^{+7.8} _{-8.1}	3.7	3.5 ^{+1.0} _{-1.2}	dust
ACT-S J003134-514317	00 : 31 : 34.3	-51 : 43 : 17.7	14.6	54.4	53.6 ^{+4.1} _{-4.1}	9.1	43.0	41.8 ^{+5.0} _{-5.0}	-0.6	-0.6 ^{+0.3} _{-0.4}	sync	J003134-514325	78.0±4.0
ACT-S J003735-530736	00 : 37 : 35.2	-53 : 07 : 36.7	12.6	39.1	38.3 ^{+3.4} _{-3.3}	7.8	35.9	34.6 ^{+4.7} _{-4.8}	-0.2	-0.3 ^{+0.4} _{-0.4}	sync	J003735-530733	81.0±4.0
ACT-S J004622-500630	00 : 46 : 22.6	-50 : 06 : 30.2	5.9	30.1	27.1 ^{+5.3} _{-5.4}	3.6	23.4	19.8 ^{+6.5} _{-6.1}	-0.8	-0.8 ^{+0.9} _{-1.0}	sync
ACT-S J004905-552112	00 : 49 : 05.9	-55 : 21 : 12.0	14.2	60.8	59.8 ^{+4.6} _{-4.6}	8.4	53.2	51.6 ^{+6.6} _{-6.7}	-0.4	-0.4 ^{+0.4} _{-0.4}	sync	J004905-552110	89.0±5.0
ACT-S J004949-540242	00 : 49 : 49.0	-54 : 02 : 42.0	7.0	22.0	20.2 ^{+3.2} _{-3.2}	3.0	12.0	10.3 ^{+3.6} _{-3.6}	-1.8	-1.8 ^{+0.9} _{-0.8}	sync	J004950-540255*	32.7±0.2
ACT-S J004957-554445	00 : 49 : 57.9	-55 : 44 : 45.1	5.6	24.6	21.6 ^{+3.8} _{-4.7}	2.9	17.2	13.7 ^{+5.7} _{-4.5}	-1.2	-1.2 ^{+1.1} _{-1.1}	sync
ACT-S J005240-531132	00 : 52 : 40.4	-53 : 11 : 32.4	6.0	18.8	16.9 ^{+3.2} _{-3.3}	3.6	14.9	12.5 ^{+4.2} _{-3.9}	-0.7	-0.8 ^{+0.9} _{-1.0}	sync	J005242-531142*	23.7±0.3
ACT-S J005244-494725	00 : 52 : 44.2	-49 : 47 : 25.7	7.5	40.1	37.8 ^{+5.5} _{-5.5}	4.3	30.5	27.1 ^{+7.2} _{-7.0}	-0.8	-0.9 ^{+0.7} _{-0.8}	sync	J005243-494732	76.0±4.0
ACT-S J005605-524156	00 : 56 : 05.3	-52 : 41 : 56.9	10.6	34.6	33.6 ^{+3.4} _{-3.4}	6.3	29.2	27.6 ^{+4.8} _{-4.8}	-0.5	-0.5 ^{+0.5} _{-0.5}	sync	J005604-524147*	21.7±0.1
ACT-S J005622-531848	00 : 56 : 22.0	-53 : 18 : 48.0	11.0	34.5	33.6 ^{+3.3} _{-3.3}	5.6	22.7	21.2 ^{+4.2} _{-4.1}	-1.1	-1.2 ^{+0.6} _{-0.6}	sync	J005622-531903*	12.9±0.1
ACT-S J005706-521425	00 : 57 : 06.1	-52 : 14 : 25.7	13.0	42.0	41.3 ^{+3.4} _{-3.4}	7.6	35.4	34.0 ^{+4.9} _{-4.8}	-0.5	-0.5 ^{+0.4} _{-0.4}	sync	J005705-521418	97.0±5.0
ACT-S J005855-521930	00 : 58 : 55.9	-52 : 19 : 30.7	21.9	71.2	70.7 ^{+3.9} _{-3.9}	10.3	47.5	46.6 ^{+4.9} _{-4.9}	-1.1	-1.1 ^{+0.3} _{-0.3}	sync	J005855-521926	170.0±9.0
ACT-S J010308-510917	01 : 03 : 08.1	-51 : 09 : 17.2	9.1	32.2	31.0 ^{+3.6} _{-3.7}	5.8	27.0	25.2 ^{+4.8} _{-4.1}	-0.5	-0.5 ^{+0.5} _{-0.6}	sync	J010306-510907	82.0±4.0
ACT-S J010330-513550	01 : 03 : 30.3	-51 : 35 : 50.9	13.1	43.5	42.7 ^{+3.5} _{-3.6}	10.2	48.5	47.5 ^{+5.0} _{-5.0}	0.3	0.3 ^{+0.3} _{-0.3}	sync	J010329-513551	87.0±4.0
ACT-S J011324-532945	01 : 13 : 24.3	-53 : 29 : 45.5	7.3	21.9	20.5 ^{+3.1} _{-3.1}	4.0	17.1	14.7 ^{+4.4} _{-4.2}	-0.8	-0.8 ^{+0.8} _{-0.9}	sync	J011323-532949	117.0±6.0
ACT-S J011650-544639	01 : 16 : 50.5	-54 : 46 : 39.2	5.5	19.4	17.1 ^{+3.8} _{-3.1}	4.4	20.3	17.9 ^{+4.7} _{-4.7}	0.1	0.1 ^{+0.9} _{-0.9}	sync
ACT-S J011829-511527	01 : 18 : 29.9	-51 : 15 : 27.3	14.4	49.5	48.8 ^{+3.7} _{-3.2}	10.5	52.1	51.0 ^{+5.2} _{-4.7}	0.1	0.1 ^{+0.3} _{-0.3}	sync	J011828-511524*	40.6±0.2
ACT-S J011950-535718	01 : 19 : 50.7	-53 : 57 : 18.9	52.9	158.7	158.4 ^{+5.4} _{-5.7}	30.7	129.8	129.2 ^{+6.2} _{-5.9}	-0.5	-0.5 ^{+0.1} _{-0.2}	sync	J011950-535717	245.0±12.0
ACT-S J012457-511315	01 : 24 : 57.3	-51 : 13 : 15.0	55.4	196.8	196.4 ^{+6.6} _{-6.6}	32.9	163.5	162.9 ^{+7.7} _{-7.4}	-0.5	-0.5 ^{+0.2} _{-0.1}	sync	J012457-511316	745.0±37.0
ACT-S J012623-510313	01 : 26 : 23.2	-51 : 03 : 13.2	7.0	24.1	22.5 ^{+3.5} _{-3.6}	5.0	22.8	20.7 ^{+4.7} _{-4.7}	-0.2	-0.2 ^{+0.7} _{-0.8}	sync	J012624-510309	111.0±6.0
ACT-S J012755-513646	01 : 27 : 55.8	-51 : 36 : 46.1	13.2	41.2	40.4 ^{+3.3} _{-3.3}	8.4	38.0	36.8 ^{+4.8} _{-4.7}	-0.2	-0.2 ^{+0.4} _{-0.4}	sync	J012756-513641	100.0±5.0
ACT-S J012756-532936	01 : 27 : 56.9	-53 : 29 : 36.2	10.0	27.7	26.8 ^{+2.9} _{-2.9}	7.0	30.1	28.7 ^{+4.5} _{-4.4}	0.2	0.2 ^{+0.5} _{-0.5}	sync	J012759-533007*	37.1±0.2
ACT-S J012835-525520	01 : 28 : 35.3	-52 : 55 : 20.6	6.2	17.2	15.7 ^{+2.9} _{-2.9}	5.1	19.4	17.7 ^{+3.9} _{-3.9}	0.3	0.3 ^{+0.7} _{-0.8}	sync	J012834-525519	175.0±9.0
ACT-S J013106-545714	01 : 31 : 06.6	-54 : 57 : 14.3	5.3	19.6	16.6 ^{+3.9} _{-4.2}	2.9	11.9	9.6 ^{+3.8} _{-3.1}	-1.6	-1.4 ^{+1.1} _{-1.0}	sync	J013108-545732*	31.6±0.2
ACT-S J013306-520006	01 : 33 : 06.0	-52 : 00 : 06.1	81.6	241.0	240.6 ^{+7.2} _{-7.7}	43.3	179.5	179.1 ^{+7.2} _{-7.7}	-0.8	-0.8 ^{+0.1} _{-0.1}	sync	J013305-520003	1198.0±60.0
ACT-S J013409-552615	01 : 34 : 09.7	-55 : 26 : 15.3	5.8	23.0	20.7 ^{+4.1} _{-4.2}	4.4	22.0	19.4 ^{+5.1} _{-5.1}	-0.2	-0.2 ^{+0.8} _{-0.9}	sync	J013408-552616	47.0±3.0
ACT-S J013540-514936	01 : 35 : 40.8	-51 : 49 : 36.0	7.3	20.6	19.3 ^{+2.9} _{-3.0}	5.1	19.0	17.3 ^{+3.8} _{-3.8}	-0.3	-0.3 ^{+0.7} _{-0.7}	sync	J013540-514945	60.0±3.0
ACT-S J013548-524417	01 : 35 : 48.7	-52 : 44 : 17.2	19.1	52.2	51.7 ^{+3.1} _{-3.1}	9.6	38.1	37.2 ^{+4.2} _{-4.2}	-0.8	-0.8 ^{+0.3} _{-0.3}	sync	J013548-524417	55.0±3.0
ACT-S J013727-543946	01 : 37 : 27.3	-54 : 39 : 46.3	7.9	25.9	24.6 ^{+3.4} _{-3.4}	4.8	19.2	17.3 ^{+4.1} _{-4.1}	-0.9	-0.9 ^{+0.6} _{-0.8}	sync	J013726-543940*	19.2±0.1
ACT-S J013949-521745	01 : 39 : 49.8	-52 : 17 : 45.9	14.4	39.8	39.2 ^{+3.0} _{-3.0}	6.6	26.1	24.8 ^{+4.1} _{-4.1}	-1.1	-1.2 ^{+0.4} _{-0.5}	sync	J013949-521746	91.0±5.0
ACT-S J014648-520236	01 : 46 : 48.8	-52 : 02 : 36.8	29.2	82.4	82.0 ^{+3.7} _{-3.7}	16.4	67.8	67.2 ^{+4.6} _{-4.8}	-0.5	-0.5 ^{+0.2} _{-0.2}	sync	J014648-520233	226.0±11.0
ACT-S J014744-524550	01 : 47 : 44.8	-52 : 45 : 50.2	4.6	12.5	11.2 ^{+2.8} _{-2.8}	7.2	28.4	26.4 ^{+4.2} _{-4.2}	2.2	2.2 ^{+0.8} _{-0.7}	dust
ACT-S J015358-540650	01 : 53 : 58.7	-54 : 06 : 50.5	10.8	30.5	29.6 ^{+3.0} _{-2.8}	6.4	25.7	24.3 ^{+4.1} _{-4.1}	-0.5	-0.5 ^{+0.5} _{-0.5}	sync	J015358-540653	50.0±3.0
ACT-S J015419-510752	01 : 54 : 19.9	-51 : 07 : 52.7	58.9	177.3	177.0 ^{+6.0} _{-6.0}	29.2	135.4	134.8 ^{+6.3} _{-6.3}	-0.7	-0.7 ^{+0.2} _{-0.1}	sync	J015419-510751	329.0±17.0
ACT-S J015558-512544	01 : 55 : 58.3	-51 : 25 : 44.2	6.4	18.9	16.4 ^{+3.1} _{-3.1}	1.8	6.9	7.0 ^{+2.3} _{-1.6}	-3.8	-2.2 ^{+0.8} _{-0.5}	sync	J015557-512545	115.0±6.0

ID and Coordinates			148 GHz			218 GHz			Spectral Index			20 GHz	
ACT ID	RA (J2000) hh:mm:ss	Dec dd:mm:ss	S/N	S _m ^a (mJy)	S _{db} (mJy)	S/N	S _m ^b (mJy)	S _{db} (mJy)	α _m	α _{db}	Type	ATCA ID ^c	S _m (mJy)
ACT-S J015649-543948	01 : 56 : 49.9	-54 : 39 : 48.2	13.6	42.1	41.2 ^{+3.3} _{-3.3}	5.2	22.8	21.1 ^{+4.4} _{-4.2}	-1.7	-1.7 ^{+0.5} _{-0.6}	sync	J015649-543949	278.0±14.0
ACT-S J015817-500419	01 : 58 : 17.5	-50 : 04 : 19.4	6.9	27.4	25.5 ^{+4.1} _{-4.1}	5.9	31.2	29.2 ^{+5.5} _{-5.5}	0.3	0.4 ^{+0.6} _{-0.7}	sync	J015817-500419	121.0±6.0
ACT-S J015914-530857	01 : 59 : 14.3	-53 : 08 : 57.7	10.2	26.5	25.5 ^{+2.8} _{-2.7}	4.0	14.5	13.0 ^{+3.5} _{-3.0}	-1.7	-1.8 ^{+0.7} _{-0.7}	sync	J015913-530853	69.0±4.0
ACT-S J020449-550258	02 : 04 : 49.0	-55 : 02 : 58.9	15.2	49.6	48.9 ^{+2.8} _{-3.5}	10.4	47.7	46.7 ^{+4.8} _{-4.9}	-0.1	-0.1 ^{+0.3} _{-0.3}	sync	J020454-550337*	29.9±0.3
ACT-S J020648-534536	02 : 06 : 48.2	-53 : 45 : 36.9	6.2	16.4	14.8 ^{+2.8} _{-2.8}	3.7	13.2	11.1 ^{+3.6} _{-3.4}	-0.7	-0.8 ^{+0.9} _{-1.0}	sync	J020647-534543	60.0±3.0
ACT-S J020921-522926	02 : 09 : 21.7	-52 : 29 : 26.2	13.8	35.3	34.7 ^{+2.8} _{-2.7}	9.2	34.7	33.8 ^{+4.0} _{-4.0}	-0.1	-0.1 ^{+0.3} _{-0.4}	sync	J020916-522949*	22.8±0.2
ACT-S J021046-510103	02 : 10 : 46.6	-51 : 01 : 03.3	532.9	1721.0	1718.3 ^{+48.4} _{-51.6}	301.5	1420.0	1418.2 ^{+48.4} _{-48.4}	-0.5	-0.5 ^{+0.1} _{-0.1}	sync	J021046-510101	3287.0±164.0
ACT-S J021238-500412	02 : 12 : 38.8	-50 : 04 : 12.6	5.2	21.0	17.5 ^{+4.4} _{-5.0}	2.7	13.8	10.6 ^{+4.8} _{-3.8}	-1.4	-1.2 ^{+1.2} _{-1.1}	sync	J021240-500445*	11.3±0.1
ACT-S J021520-510437	02 : 15 : 20.0	-51 : 04 : 37.5	5.6	16.5	14.5 ^{+3.1} _{-3.2}	3.2	13.0	10.4 ^{+4.0} _{-3.5}	-0.8	-0.8 ^{+1.1} _{-1.0}	sync	J021518-510426*	7.8±0.2
ACT-S J021603-520009	02 : 16 : 03.3	-52 : 00 : 09.7	13.7	34.6	33.9 ^{+2.7} _{-2.7}	7.3	28.8	27.6 ^{+4.1} _{-4.1}	-0.5	-0.5 ^{+0.4} _{-0.5}	sync	J021603-520012	211.0±11.0
ACT-S J021835-550350	02 : 18 : 35.4	-55 : 03 : 50.9	13.2	43.9	43.1 ^{+3.5} _{-3.6}	6.4	30.1	28.5 ^{+4.8} _{-4.8}	-1.0	-1.1 ^{+0.4} _{-0.5}	sync	J021834-550350	54.0±3.0
ACT-S J022215-510633	02 : 22 : 15.7	-51 : 06 : 33.5	10.7	31.2	30.2 ^{+3.1} _{-3.1}	4.6	19.0	17.1 ^{+4.2} _{-3.9}	-1.4	-1.5 ^{+0.6} _{-0.7}	sync	J022215-510629	75.0±3.0
ACT-S J022331-534742	02 : 23 : 31.0	-53 : 47 : 42.1	46.7	123.4	123.2 ^{+4.4} _{-4.4}	24.6	95.9	95.4 ^{+5.3} _{-5.1}	-0.7	-0.7 ^{+0.2} _{-0.2}	sync	J022330-534740	464.0±23.0
ACT-S J022530-522552	02 : 25 : 30.2	-52 : 25 : 52.3	16.7	41.3	40.8 ^{+2.7} _{-2.7}	8.2	31.5	30.5 ^{+4.0} _{-3.9}	-0.7	-0.8 ^{+0.4} _{-0.4}	sync	J022529-522555	52.0±3.0
ACT-S J022803-490126	02 : 28 : 03.7	-49 : 01 : 26.5	6.8	39.5	36.7 ^{+6.0} _{-6.0}	4.1	32.4	28.2 ^{+8.1} _{-7.9}	-0.6	-0.7 ^{+0.8} _{-0.9}	sync
ACT-S J022820-553738	02 : 28 : 20.6	-55 : 37 : 38.2	7.0	28.0	26.2 ^{+4.1} _{-4.2}	5.4	27.9	25.8 ^{+5.3} _{-5.3}	-0.0	-0.0 ^{+0.7} _{-0.7}	sync	J022820-553725	89.0±5.0
ACT-S J022821-554605	02 : 28 : 21.4	-55 : 46 : 05.6	26.3	109.0	108.5 ^{+5.1} _{-5.3}	14.4	83.8	83.0 ^{+6.3} _{-6.5}	-0.7	-0.7 ^{+0.2} _{-0.2}	sync	J022821-554603	391.0±18.0
ACT-S J022912-540325	02 : 29 : 12.8	-54 : 03 : 25.3	121.5	337.1	336.6 ^{+10.1} _{-10.1}	66.0	268.6	268.3 ^{+10.1} _{-10.1}	-0.6	-0.6 ^{+0.1} _{-0.1}	sync	J022912-540324	338.0±17.0
ACT-S J022925-523225	02 : 29 : 25.7	-52 : 32 : 25.8	19.0	47.0	46.5 ^{+2.8} _{-2.8}	9.7	37.3	36.4 ^{+4.0} _{-4.0}	-0.6	-0.6 ^{+0.3} _{-0.3}	sync	J022925-523226	182.0±9.0
ACT-S J023246-535635	02 : 32 : 46.1	-53 : 56 : 35.8	9.6	24.8	23.9 ^{+2.7} _{-2.7}	5.8	20.2	18.9 ^{+3.6} _{-3.6}	-0.6	-0.6 ^{+0.5} _{-0.6}	sync	J023246-535636*	32.0±0.2
ACT-S J023356-503021	02 : 33 : 56.8	-50 : 30 : 21.7	5.9	18.9	16.5 ^{+3.3} _{-3.4}	2.1	10.2	8.4 ^{+3.7} _{-3.4}	-2.3	-1.8 ^{+1.1} _{-0.8}	sync	J023356-503020	62.0±3.0
ACT-S J023600-530237 ^d	02 : 36 : 00.8	-53 : 02 : 37.1	5.1	12.6	10.1 ^{+2.8} _{-3.7}	2.5	9.1	6.4 ^{+3.4} _{-2.7}	-1.2	-1.0 ^{+1.4} _{-1.2}	sync
ACT-S J023923-510822	02 : 39 : 23.2	-51 : 08 : 22.6	7.3	20.5	19.2 ^{+2.9} _{-2.9}	4.9	20.2	18.3 ^{+4.2} _{-4.3}	-0.1	-0.1 ^{+0.7} _{-0.8}	sync	J023923-510817*	30.9±0.2
ACT-S J024039-542932	02 : 40 : 39.8	-54 : 29 : 32.9	7.2	21.7	20.3 ^{+3.1} _{-3.1}	5.1	21.0	19.1 ^{+4.2} _{-4.2}	-0.1	-0.1 ^{+0.6} _{-0.7}	sync	J024040-542933	58.0±3.0
ACT-S J024137-492447	02 : 41 : 37.6	-49 : 24 : 47.9	10.2	44.5	43.1 ^{+4.5} _{-4.6}	5.6	36.7	34.1 ^{+6.8} _{-6.9}	-0.6	-0.6 ^{+0.5} _{-0.6}	sync	J024136-492452	99.0±5.0
ACT-S J024154-534546	02 : 41 : 54.5	-53 : 45 : 46.2	6.9	17.6	16.3 ^{+2.7} _{-3.1}	5.0	18.1	16.4 ^{+3.7} _{-3.7}	0.0	0.0 ^{+0.7} _{-0.7}	sync	J024155-534548*	56.5±0.2
ACT-S J024313-510516	02 : 43 : 13.7	-51 : 05 : 16.2	13.9	40.1	39.4 ^{+3.1} _{-3.1}	6.7	31.6	30.0 ^{+4.9} _{-4.9}	-0.7	-0.7 ^{+0.4} _{-0.5}	sync	J024313-510517	189.0±7.0
ACT-S J024342-532803	02 : 43 : 42.2	-53 : 28 : 03.4	5.9	14.5	13.0 ^{+2.6} _{-2.7}	3.9	14.5	12.1 ^{+3.8} _{-3.8}	-0.1	-0.2 ^{+0.9} _{-1.0}	sync
ACT-S J024430-541605 ^d	02 : 44 : 30.3	-54 : 16 : 05.3	3.5	10.0	8.3 ^{+2.9} _{-2.8}	5.6	24.2	20.5 ^{+4.8} _{-5.2}	2.4	2.4 ^{+1.2} _{-1.1}	dust
ACT-S J024509-554419	02 : 45 : 09.2	-55 : 44 : 19.6	2.2	9.3	11.9 ^{+2.3} _{-1.5}	12.0	72.8	68.8 ^{+6.4} _{-6.5}	6.0	4.6 ^{+0.3} _{-0.5}	dust
ACT-S J024540-525758	02 : 45 : 40.4	-52 : 57 : 58.2	7.0	16.8	15.6 ^{+2.5} _{-2.5}	5.1	18.4	16.8 ^{+3.7} _{-3.7}	0.2	0.2 ^{+0.7} _{-0.7}	sync	J024539-525748*	24.0±0.2
ACT-S J024615-495354	02 : 46 : 15.1	-49 : 53 : 54.4	10.7	42.0	40.5 ^{+4.1} _{-4.1}	3.7	20.9	19.1 ^{+5.2} _{-4.1}	-2.0	-2.0 ^{+0.6} _{-0.6}	sync	J024614-495350	70.0±4.0
ACT-S J024615-552739	02 : 46 : 15.6	-55 : 27 : 39.0	3.2	11.8	9.5 ^{+3.7} _{-3.3}	5.5	32.1	27.6 ^{+6.3} _{-6.8}	2.8	2.7 ^{+1.2} _{-1.1}	dust
ACT-S J024647-502608	02 : 46 : 47.0	-50 : 26 : 08.4	5.5	18.4	16.3 ^{+3.5} _{-3.6}	4.9	24.1	21.8 ^{+5.0} _{-5.1}	0.7	0.8 ^{+0.8} _{-0.8}	sync
ACT-S J024948-555627	02 : 49 : 48.2	-55 : 56 : 27.2	5.5	23.2	20.6 ^{+4.4} _{-4.5}	5.4	29.1	26.8 ^{+5.5} _{-5.6}	0.6	0.7 ^{+0.8} _{-0.8}	sync	J024948-555615	95.0±3.0
ACT-S J025112-520812	02 : 51 : 12.9	-52 : 08 : 12.8	6.2	15.2	13.8 ^{+2.6} _{-2.6}	3.9	14.2	12.1 ^{+3.7} _{-3.6}	-0.3	-0.3 ^{+0.8} _{-1.0}	sync	J025109-520801*	55.9±0.2
ACT-S J025204-514560	02 : 52 : 04.0	-51 : 45 : 60.0	5.5	13.9	11.9 ^{+2.7} _{-2.9}	2.8	10.4	8.0 ^{+3.6} _{-2.8}	-1.0	-1.0 ^{+1.1} _{-1.1}	sync	J025201-514546*	4.8±0.9
ACT-S J025329-544152	02 : 53 : 29.3	-54 : 41 : 52.4	254.3	830.0	828.7 ^{+23.3} _{-24.9}	127.6	659.4	659.1 ^{+23.3} _{-23.3}	-0.6	-0.6 ^{+0.1} _{-0.1}	sync	J025329-544151	1933.0±96.0
ACT-S J025353-495321	02 : 53 : 53.0	-49 : 53 : 21.8	2.4	9.3	8.1 ^{+3.1} _{-2.1}	6.3	40.0	35.1 ^{+6.8} _{-6.8}	4.3	3.8 ^{+0.8} _{-1.0}	dust
ACT-S J025629-522724	02 : 56 : 29.9	-52 : 27 : 24.7	5.9	14.9	13.4 ^{+2.7} _{-2.7}	4.9	17.4	15.7 ^{+3.6} _{-3.7}	0.4	0.4 ^{+0.8} _{-0.8}	sync
ACT-S J025839-505203	02 : 58 : 39.1	-50 : 52 : 03.5	37.9	115.5	115.0 ^{+4.5} _{-4.3}	18.1	85.4	84.7 ^{+5.6} _{-5.6}	-0.8	-0.8 ^{+0.2} _{-0.2}	sync	J025838-505204	314.0±15.0
ACT-S J025849-533201	02 : 58 : 49.7	-53 : 32 : 01.0	6.9	17.7	16.4 ^{+2.7} _{-2.7}	4.9	19.0	17.1 ^{+4.0} _{-4.0}	0.1	0.1 ^{+0.7} _{-0.8}	sync	J025847-533205*	10.2±0.8

ID and Coordinates			148 GHz			218 GHz			Spectral Index			20 GHz	
ACT ID	RA	Dec	S/N	S _m ^a	S _{db}	S/N	S _m ^b	S _{db}	α _m	α _{db}	Type	ATCA ID ^c	S _m
	hh:mm:ss	dd:mm:ss		(mJy)	(mJy)		(mJy)	(mJy)					(mJy)
ACT-S J030056-510236	03 : 00 : 56.6	-51 : 02 : 36.3	11.2	34.3	33.4 ^{+3.3} _{-3.2}	6.0	27.2	25.5 ^{+4.7} _{-4.7}	-0.7	-0.7 ^{+0.5} _{-0.6}	sync	J030055-510229	104.0±5.0
ACT-S J030132-525604	03 : 01 : 32.6	-52 : 56 : 04.6	15.6	39.7	39.1 ^{+2.8} _{-2.7}	7.7	31.2	30.0 ^{+4.2} _{-4.1}	-0.7	-0.7 ^{+0.4} _{-0.4}	sync	J030132-525555*	34.8±0.1
ACT-S J030328-523431	03 : 03 : 28.3	-52 : 34 : 31.1	24.9	64.5	64.1 ^{+3.3} _{-3.1}	13.1	52.4	51.8 ^{+4.3} _{-4.3}	-0.5	-0.5 ^{+0.2} _{-0.3}	sync	J030328-523433	89.0±5.0
ACT-S J030615-552806	03 : 06 : 15.8	-55 : 28 : 06.3	14.1	54.1	53.2 ^{+4.1} _{-4.1}	8.4	53.3	51.6 ^{+6.7} _{-6.7}	-0.1	-0.1 ^{+0.4} _{-0.4}	sync	J030616-552808	109.0±6.0
ACT-S J031001-532003	03 : 10 : 01.3	-53 : 20 : 03.8	3.9	10.1	8.7 ^{+2.6} _{-2.4}	7.6	31.4	29.3 ^{+4.3} _{-4.4}	3.1	3.2 ^{+0.9} _{-0.8}	dust
ACT-S J031207-554137	03 : 12 : 07.0	-55 : 41 : 37.0	18.6	74.3	73.5 ^{+4.6} _{-4.4}	9.1	59.8	58.3 ^{+6.8} _{-6.9}	-0.6	-0.6 ^{+0.3} _{-0.3}	sync	J031207-554133	294.0±15.0
ACT-S J031426-510432	03 : 14 : 26.1	-51 : 04 : 32.2	31.2	93.4	93.0 ^{+4.0} _{-4.0}	18.2	84.7	84.1 ^{+5.4} _{-5.4}	-0.3	-0.3 ^{+0.2} _{-0.2}	sync	J031425-510431	112.0±6.0
ACT-S J031823-533148	03 : 18 : 23.5	-53 : 31 : 48.0	19.5	52.6	52.1 ^{+3.1} _{-3.0}	7.6	32.1	30.9 ^{+4.3} _{-4.4}	-1.3	-1.3 ^{+0.4} _{-0.4}	sync	J031823-533146*	28.8±0.2
ACT-S J031911-500027	03 : 19 : 11.8	-50 : 00 : 27.4	7.0	27.5	25.5 ^{+4.0} _{-4.1}	3.6	19.0	16.2 ^{+5.3} _{-4.7}	-1.1	-1.2 ^{+0.8} _{-0.9}	sync	J031910-500026*	28.1±0.3
ACT-S J032207-535422	03 : 22 : 07.6	-53 : 54 : 22.6	6.4	17.6	16.1 ^{+2.9} _{-2.9}	4.6	17.4	15.5 ^{+3.8} _{-3.9}	-0.1	-0.1 ^{+0.8} _{-0.8}	sync	J032210-535445*	19.1±0.2
ACT-S J032212-504232	03 : 22 : 12.8	-50 : 42 : 32.4	11.3	35.3	34.4 ^{+3.3} _{-3.3}	5.0	22.7	20.7 ^{+4.7} _{-4.5}	-1.2	-1.3 ^{+0.6} _{-0.7}	sync	J032212-504233*	40.7±0.3
ACT-S J032327-522632	03 : 23 : 27.5	-52 : 26 : 32.2	15.2	40.6	39.9 ^{+4.2} _{-2.9}	9.5	37.7	36.8 ^{+4.2} _{-4.3}	-0.2	-0.2 ^{+0.3} _{-0.4}	sync	J032327-522630	72.0±4.0
ACT-S J032540-524710	03 : 25 : 40.4	-52 : 47 : 10.6	2.8	7.1	5.9 ^{+2.3} _{-1.7}	6.5	26.2	23.3 ^{+4.3} _{-4.4}	3.7	3.6 ^{+0.9} _{-1.0}	dust
ACT-S J032650-533702	03 : 26 : 50.9	-53 : 37 : 02.3	17.9	47.5	47.0 ^{+2.9} _{-3.0}	9.5	41.5	40.5 ^{+4.6} _{-4.6}	-0.4	-0.4 ^{+0.3} _{-0.3}	sync	J032650-533701	91.0±5.0
ACT-S J033002-503517	03 : 30 : 02.3	-50 : 35 : 17.1	7.3	23.2	21.7 ^{+3.3} _{-3.2}	4.5	20.3	18.0 ^{+4.7} _{-4.6}	-0.4	-0.5 ^{+0.7} _{-0.8}	sync	J033002-503519	80.0±4.0
ACT-S J033115-524142	03 : 31 : 15.2	-52 : 41 : 42.3	10.6	27.2	26.5 ^{+2.7} _{-2.7}	6.7	27.1	25.7 ^{+4.2} _{-4.2}	-0.0	-0.1 ^{+0.5} _{-0.5}	sync	J033114-524148	55.0±3.0
ACT-S J033126-525829	03 : 31 : 26.2	-52 : 58 : 29.3	11.6	29.3	28.6 ^{+2.6} _{-2.7}	7.4	30.0	28.8 ^{+4.2} _{-4.2}	0.0	0.0 ^{+0.4} _{-0.5}	sync	J033126-525830	81.0±4.0
ACT-S J033134-515355	03 : 31 : 34.9	-51 : 53 : 55.2	6.1	15.9	15.0 ^{+2.6} _{-2.7}	11.9	48.4	47.3 ^{+4.4} _{-4.4}	2.9	3.0 ^{+0.5} _{-0.5}	dust
ACT-S J033444-521852	03 : 34 : 44.4	-52 : 18 : 52.7	7.7	19.9	18.8 ^{+2.7} _{-2.7}	5.2	18.7	17.1 ^{+3.7} _{-3.7}	-0.2	-0.2 ^{+0.6} _{-0.7}	sync	J033443-521900*	30.2±0.1
ACT-S J033554-543029	03 : 35 : 54.1	-54 : 30 : 29.9	15.1	43.5	42.9 ^{+3.2} _{-3.2}	5.9	25.2	23.6 ^{+4.4} _{-4.3}	-1.5	-1.5 ^{+0.5} _{-0.5}	sync	J033553-543025	305.0±15.0
ACT-S J034156-515146	03 : 41 : 56.1	-51 : 51 : 46.5	6.5	16.3	14.4 ^{+2.6} _{-2.6}	1.8	6.5	6.3 ^{+2.2} _{-2.2}	-3.5	-2.2 ^{+0.6} _{-0.6}	sync	J034154-515144*	17.0±0.1
ACT-S J034349-524116	03 : 43 : 49.5	-52 : 41 : 16.8	8.8	22.2	21.1 ^{+2.6} _{-2.7}	4.1	14.2	12.5 ^{+3.5} _{-3.2}	-1.3	-1.4 ^{+0.7} _{-0.8}	sync	J034349-524116	171.0±9.0
ACT-S J034640-520505	03 : 46 : 40.5	-52 : 05 : 05.9	7.0	17.5	16.7 ^{+2.5} _{-2.6}	12.3	48.2	47.1 ^{+4.3} _{-4.2}	2.7	2.7 ^{+0.5} _{-0.4}	dust
ACT-S J034940-540109	03 : 49 : 40.5	-54 : 01 : 09.9	6.5	18.8	17.3 ^{+3.0} _{-3.0}	5.3	20.6	18.9 ^{+4.0} _{-4.0}	0.2	0.2 ^{+0.7} _{-0.7}	sync	J034941-540106	80.0±9.0
ACT-S J035034-524801 ^d	03 : 50 : 34.2	-52 : 48 : 01.7	1.7	4.3	3.1 ^{+1.7} _{-1.9}	5.3	20.1	12.9 ^{+5.6} _{-11.2}	5.4	3.5 ^{+1.0} _{-2.1}	dust
ACT-S J035128-514256	03 : 51 : 28.3	-51 : 42 : 56.3	54.2	143.7	143.4 ^{+4.8} _{-4.8}	28.1	116.6	116.2 ^{+5.7} _{-5.9}	-0.5	-0.5 ^{+0.2} _{-0.2}	sync	J035128-514254	401.0±20.0
ACT-S J035700-495549	03 : 57 : 00.7	-49 : 55 : 49.3	14.3	56.9	56.0 ^{+4.4} _{-4.4}	7.1	45.1	43.2 ^{+6.6} _{-6.6}	-0.6	-0.7 ^{+0.4} _{-0.5}	sync	J035700-495547	74.0±4.0
ACT-S J035840-543405	03 : 58 : 40.8	-54 : 34 : 05.0	9.1	31.5	30.3 ^{+3.6} _{-3.6}	6.4	33.1	31.4 ^{+5.3} _{-5.4}	0.1	0.1 ^{+0.5} _{-0.5}	sync	J035842-543407*	21.6±0.2
ACT-S J040400-552021	04 : 04 : 00.1	-55 : 20 : 21.5	7.0	28.7	26.8 ^{+4.2} _{-4.2}	4.2	21.9	19.3 ^{+5.3} _{-5.2}	-0.8	-0.8 ^{+0.8} _{-0.9}	sync	J040400-552023	84.0±10.0
ACT-S J040403-540555	04 : 04 : 03.1	-54 : 05 : 55.8	2.7	8.5	8.3 ^{+2.3} _{-1.6}	9.4	43.5	40.9 ^{+4.9} _{-4.8}	4.6	4.2 ^{+0.6} _{-0.7}	dust
ACT-S J040622-503501	04 : 06 : 22.2	-50 : 35 : 01.4	5.5	19.6	17.0 ^{+3.8} _{-4.0}	2.9	13.7	10.8 ^{+4.6} _{-3.6}	-1.2	-1.1 ^{+1.1} _{-1.1}	sync	J040621-503504*	48.6±0.3
ACT-S J041137-514921	04 : 11 : 37.6	-51 : 49 : 21.7	22.4	70.1	69.6 ^{+3.7} _{-3.7}	12.4	55.3	54.5 ^{+4.7} _{-4.9}	-0.6	-0.6 ^{+0.3} _{-0.3}	sync	J041137-514923	259.0±12.0
ACT-S J041249-560053	04 : 12 : 49.4	-56 : 00 : 53.6	4.7	21.4	19.2 ^{+4.8} _{-4.8}	5.6	34.5	30.2 ^{+6.7} _{-7.0}	1.3	1.2 ^{+0.9} _{-0.9}	sync	J041247-560035	118.0±6.0
ACT-S J041313-533155	04 : 13 : 13.7	-53 : 31 : 55.4	12.7	37.7	36.9 ^{+3.2} _{-3.2}	8.0	36.1	34.9 ^{+4.7} _{-4.7}	-0.1	-0.1 ^{+0.4} _{-0.4}	sync	J041313-533200	163.0±7.0
ACT-S J041436-560349	04 : 14 : 36.3	-56 : 03 : 49.5	4.3	19.9	20.2 ^{+3.6} _{-2.6}	17.6	114.5	112.0 ^{+7.5} _{-7.5}	4.7	4.5 ^{+0.3} _{-0.5}	dust
ACT-S J042000-545622	04 : 20 : 00.5	-54 : 56 : 22.4	7.7	25.3	24.6 ^{+3.2} _{-2.9}	26.0	130.9	129.8 ^{+6.6} _{-6.6}	4.3	4.3 ^{+0.4} _{-0.3}	dust
ACT-S J042504-533159	04 : 25 : 04.2	-53 : 31 : 59.2	53.4	150.3	150.0 ^{+5.1} _{-5.4}	32.9	135.6	135.1 ^{+6.2} _{-6.2}	-0.3	-0.3 ^{+0.1} _{-0.1}	sync	J042504-533158	184.0±9.0
ACT-S J042842-500530	04 : 28 : 42.9	-50 : 05 : 30.0	35.0	162.9	162.2 ^{+6.7} _{-6.7}	18.3	122.6	121.6 ^{+7.9} _{-7.9}	-0.7	-0.8 ^{+0.2} _{-0.2}	sync	J042842-500534	238.0±11.0
ACT-S J042851-543005	04 : 28 : 51.8	-54 : 30 : 05.2	6.7	21.6	20.0 ^{+3.4} _{-3.3}	5.6	28.7	26.6 ^{+5.3} _{-5.3}	0.7	0.7 ^{+0.6} _{-0.7}	sync	J042852-543007	54.0±3.0
ACT-S J042907-534943	04 : 29 : 07.0	-53 : 49 : 43.6	31.3	87.9	87.5 ^{+3.8} _{-3.8}	16.7	69.4	68.7 ^{+4.8} _{-4.8}	-0.6	-0.6 ^{+0.2} _{-0.2}	sync	J042908-534940	145.0±4.0
ACT-S J043221-510927	04 : 32 : 21.3	-51 : 09 : 27.7	26.9	87.1	86.7 ^{+4.1} _{-4.1}	13.3	64.1	63.4 ^{+5.2} _{-5.4}	-0.8	-0.8 ^{+0.2} _{-0.2}	sync	J043221-510925	319.0±15.0
ACT-S J043651-521639	04 : 36 : 51.8	-52 : 16 : 39.1	14.4	38.8	38.2 ^{+2.9} _{-3.0}	8.3	35.9	34.8 ^{+4.5} _{-4.5}	-0.2	-0.2 ^{+0.4} _{-0.4}	sync	J043652-521639	74.0±4.0

ID and Coordinates			148 GHz			218 GHz			Spectral Index			20 GHz	
ACT ID	RA	Dec	S/N	S _m ^a	S _{db}	S/N	S _m ^b	S _{db}	α _m	α _{db}	Type	ATCA ID ^c	S _m
	hh:mm:ss	dd:mm:ss		(mJy)	(mJy)		(mJy)	(mJy)					(mJy)
ACT-S J044116-543849	04 : 41 : 16.2	-54 : 38 : 49.1	5.4	18.0	15.9 ^{+3.5} _{-3.6}	4.7	23.3	20.8 ^{+5.0} _{-5.1}	0.6	0.7 ^{+0.8} _{-0.9}	sync	J044115-543859*	0.0±0.3
ACT-S J044158-515454	04 : 41 : 58.2	-51 : 54 : 54.3	33.0	92.0	91.6 ^{+3.8} _{-3.8}	17.2	76.2	75.5 ^{+5.2} _{-5.2}	-0.5	-0.5 ^{+0.2} _{-0.2}	sync	J044158-515453	262.0±13.0
ACT-S J044502-523425	04 : 45 : 02.6	-52 : 34 : 25.6	9.8	26.7	25.8 ^{+2.9} _{-2.8}	4.8	19.1	17.3 ^{+4.0} _{-4.0}	-1.0	-1.0 ^{+0.6} _{-0.7}	sync	J044506-523448*	50.0±0.3
ACT-S J044702-510257	04 : 47 : 02.8	-51 : 02 : 57.4	5.3	17.9	14.9 ^{+3.7} _{-3.7}	2.5	12.2	9.1 ^{+4.4} _{-4.4}	-1.4	-1.2 ^{+1.2} _{-1.1}	sync	J044706-510304*	43.3±0.3
ACT-S J044747-515058	04 : 47 : 47.7	-51 : 50 : 58.2	9.6	26.5	25.6 ^{+2.9} _{-2.9}	4.9	19.9	18.1 ^{+4.2} _{-4.2}	-0.8	-0.9 ^{+0.6} _{-0.7}	sync	J044748-515100	156.0±8.0
ACT-S J044821-504138	04 : 48 : 21.9	-50 : 41 : 38.4	9.4	31.4	30.2 ^{+3.5} _{-3.5}	5.5	26.2	24.3 ^{+4.9} _{-4.9}	-0.5	-0.6 ^{+0.6} _{-0.6}	sync	J044822-504133	76.0±4.0
ACT-S J045029-534657	04 : 50 : 29.0	-53 : 46 : 57.2	8.2	22.6	21.3 ^{+2.8} _{-2.9}	3.3	12.8	11.1 ^{+3.6} _{-3.6}	-1.7	-1.7 ^{+0.8} _{-0.8}	sync	J045032-534655*	47.2±0.3
ACT-S J045103-493630	04 : 51 : 03.3	-49 : 36 : 30.2	13.4	60.5	59.4 ^{+4.9} _{-4.9}	7.4	52.2	50.2 ^{+7.2} _{-7.2}	-0.4	-0.4 ^{+0.4} _{-0.5}	sync	J045102-493626	147.0±7.0
ACT-S J045239-530637	04 : 52 : 39.2	-53 : 06 : 37.9	8.0	22.5	21.3 ^{+2.9} _{-3.0}	4.2	17.3	15.2 ^{+4.2} _{-4.0}	-0.8	-0.9 ^{+0.7} _{-0.8}	sync	J045238-530635	78.0±4.0
ACT-S J045504-553115	04 : 55 : 04.0	-55 : 31 : 15.4	10.1	38.9	37.7 ^{+4.0} _{-4.0}	6.3	39.2	37.0 ^{+6.4} _{-6.4}	-0.0	-0.0 ^{+0.5} _{-0.5}	sync	J045503-553112	67.0±4.0
ACT-S J045559-530237	04 : 55 : 59.4	-53 : 02 : 37.1	7.8	21.2	20.1 ^{+2.8} _{-2.8}	4.5	17.6	15.7 ^{+4.0} _{-4.0}	-0.6	-0.6 ^{+0.7} _{-0.8}	sync	J045558-530239	61.0±3.0
ACT-S J050019-532125	05 : 00 : 19.0	-53 : 21 : 25.1	11.0	30.0	29.2 ^{+2.8} _{-2.9}	6.4	27.2	25.7 ^{+4.3} _{-4.4}	-0.3	-0.3 ^{+0.5} _{-0.5}	sync	J050019-532121	135.0±7.0
ACT-S J050401-502311	05 : 04 : 01.6	-50 : 23 : 11.7	17.1	64.1	63.3 ^{+4.2} _{-4.1}	8.1	47.9	46.3 ^{+6.1} _{-6.2}	-0.8	-0.8 ^{+0.4} _{-0.4}	sync	J050401-502313	129.0±7.0
ACT-S J050746-515602	05 : 07 : 46.3	-51 : 56 : 02.3	6.1	17.0	15.3 ^{+3.0} _{-3.0}	3.5	13.9	11.5 ^{+4.0} _{-3.7}	-0.7	-0.8 ^{+0.9} _{-1.0}	sync	J050747-515607	85.0±4.0
ACT-S J051356-505549	05 : 13 : 56.0	-50 : 55 : 49.4	6.8	23.5	21.9 ^{+3.6} _{-3.6}	5.1	25.3	23.0 ^{+5.1} _{-5.2}	0.1	0.1 ^{+0.7} _{-0.8}	sync	J051355-505541	46.0±2.0
ACT-S J051506-534420	05 : 15 : 06.9	-53 : 44 : 20.3	1.7	4.8	5.4 ^{+1.1} _{-1.1}	7.7	32.3	28.7 ^{+4.3} _{-4.5}	6.3	4.4 ^{+0.8} _{-0.7}	dust
ACT-S J051812-514358	05 : 18 : 12.5	-51 : 43 : 58.7	9.5	28.3	27.3 ^{+3.1} _{-3.1}	6.2	30.1	28.4 ^{+4.9} _{-5.1}	0.1	0.1 ^{+0.5} _{-0.6}	sync	J051811-514404	57.0±3.0
ACT-S J051840-500545	05 : 18 : 40.0	-50 : 05 : 45.5	5.2	21.2	17.6 ^{+4.5} _{-5.4}	2.5	14.9	10.9 ^{+5.6} _{-4.3}	-1.3	-1.1 ^{+1.3} _{-1.1}	sync
ACT-S J052046-550823	05 : 20 : 46.3	-55 : 08 : 23.1	6.3	22.5	20.6 ^{+3.7} _{-3.8}	5.3	25.3	23.2 ^{+4.9} _{-5.0}	0.3	0.3 ^{+0.7} _{-0.8}	sync	J052045-550824	78.0±4.0
ACT-S J052139-491301	05 : 21 : 39.7	-49 : 13 : 01.6	5.9	33.8	30.3 ^{+6.0} _{-6.0}	2.9	27.9	21.8 ^{+9.2} _{-7.7}	-0.8	-0.9 ^{+1.1} _{-1.2}	sync
ACT-S J052317-530833	05 : 23 : 17.3	-53 : 08 : 33.8	8.4	24.1	22.9 ^{+3.0} _{-3.0}	3.7	15.6	13.5 ^{+4.1} _{-4.1}	-1.3	-1.4 ^{+0.8} _{-0.8}	sync	J052318-530837	47.0±2.0
ACT-S J052743-542609	05 : 27 : 43.4	-54 : 26 : 09.6	8.1	26.5	25.1 ^{+3.4} _{-3.4}	4.2	18.5	16.3 ^{+4.5} _{-4.3}	-1.0	-1.1 ^{+0.7} _{-0.8}	sync	J052743-542616	98.0±5.0
ACT-S J052903-543650	05 : 29 : 03.5	-54 : 36 : 50.8	1.9	6.8	6.1 ^{+2.4} _{-1.6}	6.1	32.2	27.4 ^{+5.6} _{-5.9}	5.0	4.0 ^{+0.7} _{-1.0}	dust
ACT-S J053117-550425	05 : 31 : 17.5	-55 : 04 : 25.6	12.7	47.4	46.4 ^{+4.1} _{-4.0}	6.9	38.1	36.4 ^{+5.6} _{-5.7}	-0.6	-0.6 ^{+0.4} _{-0.5}	sync	J053115-550423*	30.0±0.6
ACT-S J053208-531033	05 : 32 : 08.9	-53 : 10 : 33.5	20.1	56.7	56.2 ^{+3.3} _{-3.2}	12.2	54.6	53.8 ^{+4.8} _{-4.9}	-0.1	-0.1 ^{+0.3} _{-0.3}	sync	J053208-531035	88.0±4.0
ACT-S J053250-504709	05 : 32 : 50.5	-50 : 47 : 09.9	2.5	9.2	8.8 ^{+2.7} _{-1.8}	8.2	46.7	43.2 ^{+6.0} _{-6.0}	4.7	4.1 ^{+0.6} _{-0.8}	dust
ACT-S J053311-523827	05 : 33 : 11.9	-52 : 38 : 27.5	1.7	4.8	4.8 ^{+1.7} _{-1.2}	6.3	28.3	23.7 ^{+4.9} _{-5.1}	6.1	4.2 ^{+0.6} _{-0.9}	dust
ACT-S J053323-554935	05 : 33 : 23.8	-55 : 49 : 35.8	17.6	76.8	76.0 ^{+4.9} _{-4.9}	9.6	64.2	62.7 ^{+7.0} _{-7.0}	-0.5	-0.5 ^{+0.3} _{-0.3}	sync	J053324-554936	51.0±3.0
ACT-S J053458-543906	05 : 34 : 58.6	-54 : 39 : 06.2	8.2	28.3	27.0 ^{+3.6} _{-3.6}	5.6	27.9	26.0 ^{+5.1} _{-5.1}	-0.1	-0.1 ^{+0.6} _{-0.6}	sync	J053458-543901	61.0±3.0
ACT-S J053817-503058	05 : 38 : 17.1	-50 : 30 : 58.2	1.7	6.5	7.3 ^{+2.1} _{-1.5}	7.0	43.2	37.6 ^{+6.4} _{-6.4}	6.3	4.3 ^{+0.5} _{-0.7}	dust
ACT-S J053909-551055	05 : 39 : 09.3	-55 : 10 : 55.7	8.2	31.6	30.2 ^{+4.0} _{-4.0}	6.1	35.1	33.0 ^{+6.0} _{-6.0}	0.2	0.2 ^{+0.5} _{-0.5}	sync	J053909-551059	113.0±6.0
ACT-S J054025-530348	05 : 40 : 25.3	-53 : 03 : 48.7	10.5	30.6	29.7 ^{+3.1} _{-3.0}	7.6	34.2	32.9 ^{+4.7} _{-4.7}	0.3	0.3 ^{+0.4} _{-0.5}	sync	J054025-530346	193.0±10.0
ACT-S J054030-535626	05 : 40 : 30.0	-53 : 56 : 26.4	6.1	17.4	15.3 ^{+3.0} _{-3.1}	2.4	9.6	7.9 ^{+3.0} _{-2.2}	-2.0	-1.7 ^{+1.0} _{-0.8}	sync	J054029-535632	53.0±3.0
ACT-S J054045-541821	05 : 40 : 45.8	-54 : 18 : 21.5	133.1	435.1	434.4 ^{+13.1} _{-13.1}	68.3	341.6	341.4 ^{+12.2} _{-13.1}	-0.6	-0.6 ^{+0.1} _{-0.1}	sync	J054045-541821	1127.0±56.0
ACT-S J054223-514256	05 : 42 : 23.2	-51 : 42 : 56.3	25.2	79.6	79.1 ^{+3.9} _{-3.9}	14.0	70.1	69.3 ^{+5.5} _{-5.5}	-0.3	-0.3 ^{+0.2} _{-0.2}	sync	J054223-514257	128.0±7.0
ACT-S J054830-521836	05 : 48 : 30.2	-52 : 18 : 36.5	8.9	26.8	25.7 ^{+3.1} _{-3.1}	5.9	28.3	26.5 ^{+4.9} _{-4.9}	0.1	0.1 ^{+0.5} _{-0.6}	sync	J054833-521840*	31.5±0.3
ACT-S J054944-524626	05 : 49 : 44.1	-52 : 46 : 26.7	62.5	189.4	189.1 ^{+6.0} _{-6.4}	33.6	148.4	147.9 ^{+6.7} _{-6.7}	-0.6	-0.6 ^{+0.2} _{-0.1}	sync	J054943-524625	235.0±11.0
ACT-S J055047-530454	05 : 50 : 47.5	-53 : 04 : 54.9	8.7	25.9	24.8 ^{+3.1} _{-3.1}	5.5	25.0	23.1 ^{+4.7} _{-4.7}	-0.1	-0.2 ^{+0.6} _{-0.7}	sync	J055049-530501*	0.0±0.3
ACT-S J055115-533435	05 : 51 : 15.4	-53 : 34 : 35.3	2.5	7.5	5.7 ^{+3.7} _{-2.0}	5.5	25.1	20.3 ^{+4.3} _{-4.3}	3.5	3.2 ^{+1.1} _{-1.3}	dust
ACT-S J055139-505800	05 : 51 : 39.3	-50 : 58 : 00.8	1.9	7.0	7.0 ^{+2.4} _{-1.6}	6.7	39.3	34.3 ^{+6.2} _{-6.3}	5.4	4.2 ^{+0.6} _{-0.8}	dust
ACT-S J055152-552628	05 : 51 : 52.4	-55 : 26 : 28.9	7.6	31.3	29.6 ^{+4.3} _{-4.3}	5.1	29.5	27.0 ^{+5.9} _{-5.9}	-0.2	-0.2 ^{+0.7} _{-0.7}	sync	J055152-552632	81.0±4.0
ACT-S J055235-534926	05 : 52 : 35.5	-53 : 49 : 26.8	5.4	16.2	14.0 ^{+3.2} _{-3.3}	3.1	12.9	10.1 ^{+4.2} _{-3.6}	-0.8	-0.8 ^{+1.1} _{-1.2}	sync

ID and Coordinates			148 GHz			218 GHz			Spectral Index			20 GHz	
ACT ID	RA (J2000) hh:mm:ss	Dec dd:mm:ss	S/N	S _m ^a (mJy)	S _{db} (mJy)	S/N	S _m ^b (mJy)	S _{db} (mJy)	α _m	α _{db}	Type	ATCA ID ^c	S _m (mJy)
ACT-S J055811-502952	05 : 58 : 11.5	-50 : 29 : 52.2	15.7	61.8	61.0 ^{+4.3} _{-4.4}	8.8	55.6	54.1 ^{+6.6} _{-6.7}	-0.3	-0.3 ^{+0.4} _{-0.4}	sync	J055811-502948	381.0±19.0
ACT-S J055947-502646	05 : 59 : 47.2	-50 : 26 : 46.6	10.1	40.9	39.6 ^{+4.3} _{-4.3}	5.3	29.8	27.5 ^{+5.8} _{-5.8}	-0.9	-1.0 ^{+0.6} _{-0.6}	sync	J055947-502652	78.0±4.0
ACT-S J060212-542508	06 : 02 : 12.8	-54 : 25 : 08.9	10.5	35.8	34.7 ^{+3.6} _{-3.6}	6.9	35.5	33.8 ^{+5.4} _{-5.3}	-0.1	-0.1 ^{+0.5} _{-0.5}	sync	J060212-542507	114.0±5.0
ACT-S J060749-525743	06 : 07 : 49.0	-52 : 57 : 43.5	14.5	43.4	42.7 ^{+3.2} _{-3.2}	8.6	37.6	36.5 ^{+4.5} _{-4.6}	-0.4	-0.4 ^{+0.3} _{-0.4}	sync	J060749-525744	85.0±4.0
ACT-S J060849-545643	06 : 08 : 49.1	-54 : 56 : 43.5	39.6	144.3	143.7 ^{+5.7} _{-5.4}	21.6	120.3	119.6 ^{+7.0} _{-7.0}	-0.5	-0.5 ^{+0.2} _{-0.2}	sync	J060849-545642	386.0±19.0
ACT-S J061311-492604	06 : 13 : 11.3	-49 : 26 : 04.3	5.9	30.3	27.4 ^{+5.4} _{-5.4}	5.8	40.0	37.3 ^{+7.1} _{-7.1}	0.7	0.8 ^{+0.7} _{-0.7}	sync
ACT-S J061714-530608	06 : 17 : 14.9	-53 : 06 : 08.4	12.7	40.8	39.9 ^{+3.4} _{-3.4}	7.9	35.8	34.5 ^{+4.7} _{-4.7}	-0.4	-0.4 ^{+0.4} _{-0.4}	sync	J061714-530615*	26.8±0.4
ACT-S J061955-542716	06 : 19 : 55.5	-54 : 27 : 16.1	11.6	43.0	42.0 ^{+3.9} _{-3.9}	5.5	30.3	28.1 ^{+5.7} _{-5.6}	-1.0	-1.0 ^{+0.5} _{-0.5}	sync	J061955-542713	99.0±5.0
ACT-S J062143-524133	06 : 21 : 43.1	-52 : 41 : 33.2	39.6	125.3	124.8 ^{+4.9} _{-4.7}	22.2	104.1	103.4 ^{+6.1} _{-5.9}	-0.5	-0.5 ^{+0.2} _{-0.2}	sync	J062143-524132	266.0±9.0
ACT-S J062552-543852	06 : 25 : 52.1	-54 : 38 : 52.9	28.8	121.6	121.0 ^{+5.5} _{-5.5}	15.1	90.1	89.2 ^{+6.8} _{-6.6}	-0.8	-0.8 ^{+0.2} _{-0.2}	sync	J062552-543850	304.0±15.0
ACT-S J062621-534130	06 : 26 : 21.0	-53 : 41 : 30.5	13.9	45.1	44.3 ^{+3.5} _{-3.5}	7.0	35.4	33.8 ^{+5.2} _{-5.2}	-0.7	-0.7 ^{+0.4} _{-0.5}	sync	J062620-534151	253.0±4.0
ACT-S J062649-543231	06 : 26 : 49.5	-54 : 32 : 31.7	8.1	32.9	31.4 ^{+4.2} _{-4.2}	6.1	32.3	30.4 ^{+5.5} _{-5.5}	-0.1	-0.1 ^{+0.5} _{-0.6}	sync	J062648-543214	106.0±3.0
ACT-S J062747-512614 ^d	06 : 27 : 47.8	-51 : 26 : 14.3	2.5	8.5	6.2 ^{+3.2} _{-2.4}	5.3	27.1	21.5 ^{+6.0} _{-8.0}	3.4	3.1 ^{+1.2} _{-1.5}	dust
ACT-S J063159-540453	06 : 31 : 59.7	-54 : 04 : 53.7	7.4	27.5	25.9 ^{+3.8} _{-3.9}	6.4	31.2	29.5 ^{+5.0} _{-5.1}	0.3	0.3 ^{+0.6} _{-0.6}	sync	J063201-540455	68.0±4.0
ACT-S J063715-500414 ^d	06 : 37 : 15.5	-50 : 04 : 14.3	1.7	8.6	8.3 ^{+3.0} _{-2.2}	6.1	47.4	40.2 ^{+8.1} _{-8.6}	5.8	4.1 ^{+0.6} _{-0.9}	dust
ACT-S J063739-500733	06 : 37 : 39.5	-50 : 07 : 33.9	7.9	37.9	35.9 ^{+5.0} _{-5.0}	4.3	28.2	24.9 ^{+6.8} _{-6.6}	-0.9	-1.0 ^{+0.7} _{-0.8}	sync	J063738-500722	84.0±4.0
ACT-S J064110-520224	06 : 41 : 10.4	-52 : 02 : 24.2	6.5	22.8	20.7 ^{+3.6} _{-3.6}	3.2	15.0	12.4 ^{+4.5} _{-3.7}	-1.3	-1.3 ^{+0.9} _{-0.9}	sync	J064107-520225*	31.6±0.3
ACT-S J064150-555106	06 : 41 : 50.4	-55 : 51 : 06.2	5.5	27.0	23.0 ^{+5.3} _{-5.7}	2.3	16.4	12.8 ^{+6.0} _{-4.2}	-1.8	-1.5 ^{+1.2} _{-1.0}	sync	J064150-555103*	16.3±0.4
ACT-S J064320-535845	06 : 43 : 20.2	-53 : 58 : 45.3	30.0	101.7	101.2 ^{+4.4} _{-4.6}	17.1	87.2	86.6 ^{+5.9} _{-5.9}	-0.4	-0.4 ^{+0.2} _{-0.2}	sync	J064320-535846	130.0±7.0
ACT-S J064630-545111	06 : 46 : 30.2	-54 : 51 : 11.3	7.5	31.4	29.5 ^{+4.3} _{-4.3}	3.8	20.6	17.9 ^{+5.4} _{-4.9}	-1.2	-1.3 ^{+0.8} _{-0.8}	sync	J064629-545116	90.0±3.0
ACT-S J065020-501758	06 : 50 : 20.5	-50 : 17 : 58.5	6.2	34.4	31.5 ^{+5.7} _{-5.8}	4.3	31.8	27.9 ^{+7.7} _{-7.7}	-0.3	-0.3 ^{+0.8} _{-0.9}	sync
ACT-S J065207-551605 ^d	06 : 52 : 07.4	-55 : 16 : 05.5	3.6	16.3	13.8 ^{+4.5} _{-4.2}	6.4	46.0	41.9 ^{+7.6} _{-7.7}	2.8	2.9 ^{+1.0} _{-0.9}	dust
ACT-S J065518-495156	06 : 55 : 18.4	-49 : 51 : 56.1	6.6	41.2	38.2 ^{+6.5} _{-6.6}	4.7	49.8	44.6 ^{+10.8} _{-10.9}	0.4	0.4 ^{+0.7} _{-0.8}	sync	J065518-495206	66.0±3.0
ACT-S J070411-551420	07 : 04 : 11.6	-55 : 14 : 20.9	5.7	28.4	25.5 ^{+5.2} _{-5.3}	4.7	31.2	27.9 ^{+6.8} _{-6.9}	0.2	0.2 ^{+0.8} _{-0.9}	sync	J070412-551441	59.0±3.0
ACT-S J070700-502233	07 : 07 : 00.1	-50 : 22 : 33.6	7.2	46.0	41.4 ^{+6.5} _{-6.5}	2.2	17.4	17.6 ^{+5.3} _{-3.6}	-3.2	-2.3 ^{+0.8} _{-0.5}	sync	J070700-502226	114.0±6.0

^aFlux density as measured directly from the ACT 148 GHz map.

^bFlux density as measured directly from the ACT 218 GHz map.

^cAT20G or an asterisk denotes the November 2010 follow-up observations catalog.

^dAn ACT source not cross-identified with the catalogs specified in Section ??.