

NASA Mission Management Updates to the *Hinode* Science Working Group

September 2015



On behalf of the US instrument teams

Lead HOP information site: <http://www.isas.jaxa.jp/home/solar/guidance/index.html>

Guidance for Hinode scientific operations

July. 27, 2015

HOP PROPOSAL FORM

- [Message to accepted HOP proposers: Hinode Ground-based coordination protocol \(updated on Dec. 23, 2009\)](#)
 - [New policy regarding major flare watches, target of opportunity HOPs, and synoptic/long-term study HOPs \(updated on July 31, 2010\)](#)
 - [Prioritization of Flare observations for Hinode: \(updated on December 20, 2011\)](#)
 - [Prioritization of IRIS-Hinode Operations Plans \(IHOPs\) \(updated on June 18, 2015\)](#)
-
- [Introduction \(Announcement for Solar News on 1 July 2008\)](#)
 - [Guidance for proposal observations \(HOP\)](#)
 - [Deadline for submission](#)
 - [Information required in submitted proposals](#)
 - [The following types of observations are strongly recommended to be submitted as HOPs, regardless of whether the proposers are inside or outside the Hinode team:](#)
 - [Points of contact for HOP submissions](#)
 - [Information about the Hinode Instruments'](#)
 - [Remarks](#)
 - [Guideline for Hinode scientific operations](#)
 - [Message to accepted HOP proposers: Hinode Ground-based coordination protocol \(updated on Dec. 23, 2009\)](#)
 - [New policy regarding major flare watches, target of opportunity HOPs, and synoptic/long-term study HOPs \(updated on July 31, 2010\)](#)
 - [Prioritization of Flare observations for Hinode: \(updated on December 20, 2011\)](#)

2-2. HOP Observations

NASA site: <http://hinode.msfc.nasa.gov/hops.html>

HINODE OPERATION PLANS (HOPS)

Information needed for proposal development outlined below.
 Also refer to <http://www.isas.jaxa.jp/home/solar/guidance/index.html> as the lead reference.

After reviewing the material, ****SUBMIT A HOP PROPOSAL HERE****.

HOPs List & Guidance

Months/Status HOP Subsets HOP HOP list

HINODE Operation Plan (HOP) list

HOP No.	HOP Title
HOP-0012 web run 3 times	COSE VTT Hinode/SOT collaboration: Search of transient horizontal magnetic fields in the chromosphere
HOP-0013 web run 3 times	Relative structure of small magnetic elements: faculae, network and minor network in the quiet Sun
HOP-0014 web run 17 times TAD	COSE Two-Quasars Prominence Structure and Dynamics
HOP-0015 web run 3 times	Search for photospheric/chromospheric drivers of chromospheric/polar coronal eruptions
HOP-0016 web run 3 times	Coordinated Observations between Hinode Observatory & Hinode Satellite
HOP-0018 web run 2 times	Observation of an equatorial coronal hole
HOP-0019 web run 2 times	Support of SMM network flight
HOP-0020 web run 8 times TAD	The Study of Oscillatory Magnetic Connection in Unipolar Data
HOP-0021 web run 78 times	COSE (HOP040): Synoptic SOT Irradiance Scans
HOP-0022 web run 8 times	Polar Coronal Hole Observation

HOPs Coordination Calendar



POST-SUBMISSION PRODUCTIVITY UPDATES

SUBMIT HOP Results / Output

Hinode Operations Plan Productivity

Following productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina - Savage [at] nasa - gov.

* Required

HOP Number *

HOP Title *

HOP & Proposer Productivity Logs

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

137 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

[Abramenko](#)
[Aime](#)

CONTENTS

2-2. HOP Observations

Submission Form: https://docs.google.com/forms/d/1mvUqVsII EZ0ta4hbzkVqKKv_kW8x6IH1584IkvaOzX8/viewform

HINODE OPERATION PLANS (HOPS)

Information needed for proposal development outlined below.
 Also refer to <http://www.isas.jaxa.jp/home/solar/guidance/index.html> as the lead reference.

After reviewing the material, ****SUBMIT A HOP PROPOSAL HERE****

HOPs List & Guidance

Months/Events HOP Guidance HOP List HOP List

HINODE Operation Plan (HOP) List

HOP No.	HOP Title
HOP-0012 web run 3 times web run 3 times	CORE VTT Hinode/SOT collaboration: Search of transient horizontal magnetic fields in the chromosphere
HOP-0013 web run 3 times web run 3 times	Relative structure of small magnetic elements: faculae, network and minor network in the quiet Sun
HOP-0014 web run 17 times TSD	CORE Two-Quasars Prominence Structure and Dynamics
HOP-0015 web run 3 times web run 3 times	Search for photospheric/chromospheric drivers of chromospheric/penumbral changes
HOP-0016 web run 3 times web run 3 times	Coordinated Observations between Hinode Observatory & Hinode Satellite
HOP-0018 web run 2 times web run 2 times	Observation of an equatorial coronal hole
HOP-0019 web run 2 times web run 2 times	Support of SLM network flight
HOP-0020 web run 2 times TSD	The Study of Oscillatory Magnetic Connection in Unstable Data
HOP-0021 web run 78 times web run 8 times	CORE (HPOPHO): Synoptic SOT Irradiance Scans
HOP-0022 web run 8 times	Polar Coronal Hole Observation

HOPs Coordination Calendar

Hinode HOP Coordination Calendar

POST-SUBMISSION PRODUCTIVITY UPDATES

SUBMIT HOP Results / Output

Hinode Operations Plan Productivity

Following productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina . Savage [at] nasa . gov.

* Required

HOP Number *

HOP Title *

HOP & Proposer Productivity Logs

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

137 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

[Abramenko](#)

[Aime](#)

Hinode [IRIS] Operation Plan (I/HOP) Submission Form

Comments or questions about this form should be directed toward Sabrina . Savage (at) nasa . gov.

* Required

Submission Guidance & Helpful Links

Planning for Hinode operations is performed on a three month cycle that is updated monthly. At the end of every month a monthly meeting is held to confirm the observations for the coming month and to lay out the broad objectives for the second and third months.

The cut-off for consideration is the 14th day of each month. For example, requests for observations received between the 15th of June and the 14th of July will be presented and discussed at the monthly meeting held at the end of July.

It is recommended that proposers make their submissions as early as possible, so that the Science Schedule Coordinators (SSCs) have time to refine the proposals to fit the current Hinode situation.

Late submissions may be considered only exceptionally, if scheduling conflicts can be easily resolved in the operation planning meetings.

For more detailed information, refer to the following:

<http://www.isas.jaxa.jp/home/solar/guidance/index.html>

<http://hinode.msfc.nasa.gov/hops.html>

Title of Proposed Observation *

2-2. HOP Observations

Calendar: <http://hinode.msfc.nasa.gov/status.html>

Scientific Justification *

Short statement describing the observation and scientific justification. This should be as short and concise as possible, but it should still contain all the key details. This statement is important because Hinode's limited data volume situation may make it necessary to modify some planned observations on some days. The Hinode team will refer to this statement when setting priorities for which observations to perform.

Hinode Calendar

Please see <http://hinode.msfc.nasa.gov/status.html> for important planning information (e.g., Hinode & IRIS eclipse season, Focused Mode period, etc.)

Dates *

Enter "ToO" if not specific. 1. Provide the start and end dates with reason. 2. Provide the minimum number of observation days during the period. 3. Provide desires and requirements for continuity of observations, for example: "three consecutive days are desired, but not required," "three consecutive days are required," or "It is not necessary for observations to be on consecutive days."

Time window

Time window in day, if required. 1. Provide the "minimum" duration with the start and end times in UT, if it is a coordinated observation with ground-based or space-based observatories. 2. Specify whether any short interruptions (e.g., for ten-minute synoptics) are allowed over the observing periods.

TECHNICAL STATUS

- Hinode operations are nominal.
- Monthly Events
- Focused Mode Calendar (*Subject to Revision*)



XRT

- XRT operations are nominal.
- Bakeouts
 - Bakeout List
- Filter Wheel 1 is in full use.



2-2. HOP Observations

SSC planning site: http://hinode.msfc.nasa.gov/submitted_hops.html

HINODE OPERATION PLANS SUBMISSIONS -- AUGUST 2015 SSC MEETING

RECENT/UPCOMING SUBMISSIONS

HOP 289 Filament/Su : Aug 1-8 : 17-21 UT
 HOP 287 GREGOR/Verma : Aug 10-19 : 8:30-10:30 UT
 HOP 292 Brightenings/Kanoh : Aug 10+ : 3 hours min
 HOP 286 NST/Hong : Aug 17-19 : 18-21 UT
 HOP 291 Hida/Ueno : Aug 17-23 : 0-4 UT
 HOP 173 EPO/Yaji : Aug 24-29 : 2-6 UT
 HOP 252 MOSES-III/Kankelborg : **Aug 27 : 17:25-18:40 UT +/- (11:25 MDT)**
 HOP 288 Flares/Cheng : Aug 27-31 : 17-23 UT
 HOP 290 CLASP/Ishikawa : Sep 3** : 17:16-18:16 UT +/-
 HOP 257 SST/Tarbell : Sep 3-Oct 14 : 8-11 UT

NEW/UPDATED SUBMISSIONS

HOP206 -- PolarMap/Shimojo : September : Once every 3 days for 6 hours (SAA-free)
 Submitted HOP #1: Jets/Savcheva : Oct 19-25 : 6 hrs/day

Ongoing HOP # 206/81:

HOP description

Title: Polar Panorama Map for understanding Polar Reversal in Cycle 24
 ToO: No
 Proposer: Shimojo, Tsuneta, Shiota, Sako, Anjali
 Dates: September
 Times: Once every 3 days for 6 hours
 Target/Pointing: North Pole

Comments:

September is the month for the North pole, and we need to schedule runs of HOP 81 (6-days) and the Polar Panorama HOP 206 (every 2-3 days for a month). You may recall that he made the standing request at the SWG meeting this year. Both take 6-7 hours of SAA-free time per run.

Submitted HOP # 1:

Full Proposal

Title: Coordinated UV/EUV/X-Ray observations of coronal jets

Main Objective: Identify the structure and dynamics of the jet acceleration region and the mechanism for supplying plasma and acceleration

ToO: No

Proposer: Antonia Savcheva, Paola Testa, Katharine Reeves

Previous Submissions:

Savcheva: None

Testa

Reeves

Dates: 2015 Oct 19-25

Times: 6 hours per day, time of day not important

Target/Pointing: AR with some inclusion polarities on the outskirts, or near a coronal hole

SOT:

SP maps of the photospheric magnetic field

EIS:

A study similar to ISSI_jet_1, but add a few more cool lines (e.g., FeVIII, IX) with shorter exposure times, and a smaller raster size to overlap better with the IRIS FOV.

XRT:

384"x384" FOV, thin-Be filter, fast cadence (~20 sec) for as much of the observing period as telemetry allows.

Other instruments:

IRIS: medium 8-step dense raster, 4 sec exposure time, 1330 & 1400 SJIs, large linelist (OBS ID: 3600257226).

Comments:

2-2. HOP Observations

****New****

Productivity Form: https://docs.google.com/forms/d/1KeBH_RCLjzChGvra-flyKKYvtouqJ3M00Acj3SrR5_E/viewform

HINODE OPERATION PLANS (HOPS)

Information needed for proposal development outlined below.
Also refer to <http://www.isas.jaxa.jp/home/solar/guidance/index.html> as the lead reference.

After reviewing the material, ****SUBMIT A HOP PROPOSAL HERE****.

HOPs List & Guidance

HOP No.	HOP Title
HOP-0012 web run 3 times	COSE VTT Hinode/SOT collaboration: Search of transient horizontal magnetic fields in the chromosphere
HOP-0013 web run 3 times	Relative structure of small magnetic elements: faculae, network and minor network in the quiet Sun
HOP-0014 web run 17 times TSD	COSE Two-Quasars Prominence Structure and Dynamics
HOP-0015 web run 3 times	Search for photospheric/chromospheric drivers of chromospheric penumbral microflares
HOP-0016 web run 3 times	Coordinated Observations between Hinode Observatory & Hinode Satellite
HOP-0018 web run 2 times	Observation of an equatorial coronal hole
HOP-0019 web run 2 times	Support of SLM network Right
HOP-0020 web run 8 times TSD	The Study of Oscillatory Magnetic Connection in Unipolar Data
HOP-0021 web run 78 times	COSE (HOP040): Synoptic SOT Irradiance Scans
HOP-0022 web run 8 times	Polar Coronal Hole Observation

HOPs Coordination Calendar



POST-SUBMISSION PRODUCTIVITY UPDATES

SUBMIT HOP Results / Output

Hinode Operations Plan Productivity
Follow-up productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina .Savage [at] nasa . gov.

* Required

HOP Number *

HOP Title *

HOP & Proposer Productivity Logs

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

137 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

[Abramenko](#)
[Aime](#)

CONTENTS

Hinode Operations Plan Productivity
Follow-up productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina .Savage [at] nasa . gov.

* Required

HOP Number *

Abbreviated HOP Title *

Your name *

Email *

Number of runs
If the run dates listed are incorrect, please provide corrections - http://www.isas.jaxa.jp/home/solar/hinode_op/hop_list.php

Target(s) *
Brief target description (i.e., Active Region, Flare, Quiet Sun, Polar Coronal Hole, Jets, CMEs, Prominences, etc.)

2-2. HOP Observations

HOP Productivity Log: http://hinode.msfc.nasa.gov/operations/hop_assessment/HOP_Productivity_Log.html

HINODE OPERATION PLANS (HOPS)

Information needed for proposal development outlined below.
 Also refer to <http://www.isax.jaxa.jp/home/solar/guidance/index.html> as the lead reference.

After reviewing the material, ****SUBMIT A HOP PROPOSAL HERE****.

HOPs List & Guidance

HOP No.	HOP Title
HOP 0012	CORE VTT/InnovISOT collaboration: Search of transient horizontal magnetic fields in the chromosphere
HOP 0013	Relative structure of small magnetic elements: faculae, network and minor network in the quiet Sun
HOP 0014	CORE Two-Quasars Prominence Structure and Dynamics
HOP 0015	Search for photospheric/chromospheric drivers of chromospheric penumbral microflares
HOP 0016	Coordinated Observations between HinE Observatory & Hinode Satellite
HOP 0017	Observation of an equatorial coronal hole
HOP 0018	Support of SLM network Right
HOP 0019	The Study of Oribital Magnetic Connection in Unipolar Data
HOP 0020	CORE (HOP040): Synoptic SOT Irradiance Scans
HOP 0021	Polar Coronal Hole Observation

HOPs Coordination Calendar

POST-SUBMISSION PRODUCTIVITY UPDATES

SUBMIT HOP Results / Output

Hinode Operations Plan Productivity

Following productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina . Savage [at] nasa . gov.

* Required

HOP Number *

HOP Title *

HOP & Proposer Productivity Logs

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

137 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

[Abramenko](#)
[Aime](#)

CONTENTS

Questions / Comments: [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov)

[HOP INSTRUCTION PAGE](#)

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

233 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

105 Refereed Publications [Histogram Per Year](#) [Histogram Per HOP](#)

25 Non-refereed Publications [Histogram Per Year](#) [Histogram Per HOP](#)

60 EPO Activities [Histogram Per Year](#) [Histogram Per HOP](#)

43 Other Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP ####: [# Refereed Publications; # Non-refereed Publications; # EPO activities; # Other outputs] -- Total

**** -- Reason for lack of productivity noted (e.g., insufficient observations, PI relocation, etc.)**


[HOP 0072](#): [0; 0; 0; 0] -- 0
[HOP 0073](#): [16; 0; 1; 2] -- 19
[HOP 0074](#): [0; 0; 0; 0] -- 0 **
[HOP 0075](#): [0; 0; 0; 0] -- 0
[HOP 0076](#): [0; 0; 0; 0] -- 0
[HOP 0077](#): [0; 0; 0; 0] -- 0
[HOP 0078](#): [0; 0; 0; 0] -- 0
[HOP 0079](#): [3; 0; 0; 0] -- 3
[HOP 0080](#): [1; 0; 0; 0] -- 1
[HOP 0081](#): [9; 4; 1; 0] -- 14
[HOP 0082](#): [0; 0; 0; 0] -- 0
[HOP 0083](#): [0; 0; 0; 0] -- 0
[HOP 0084](#): [0; 0; 0; 0] -- 0
[HOP 0085](#): [0; 0; 0; 0] -- 0
[HOP 0086](#): [0; 0; 0; 0] -- 0
[HOP 0087](#): [0; 0; 0; 0] -- 0
[HOP 0088](#): [0; 0; 0; 0] -- 0
[HOP 0089](#): [0; 0; 0; 0] -- 0
[HOP 0090](#): [0; 0; 0; 0] -- 0
[HOP 0091](#): [0; 0; 0; 0] -- 0 **
[HOP 0092](#): [0; 0; 0; 0] -- 0
[HOP 0093](#): [4; 0; 0; 0] -- 4








** Lack of productivity reason given (e.g., insufficient observations, bad seeing, not aligned, etc.)

2-2. HOP Observations

Total Products Per Year



Total Products Per HOP



Current report (as of Sept. 10, 2015):

84 HOPs reporting (~37% for HOPs < 71 + 2 < 71)

259 Total Productivity Outputs

115 Refereed Publications

27 Non-refereed Publications [e.g., Conf. Proceedings]

67 EPO Activities

50 Other Outputs [e.g., Talks, Posters]

Notes:

1. HOP run dates not always correct on HOP listing page.

2. Complaint from Proposer (Antolin):

“The HOP number does not seem to be included in the details of the observation for the Hinode instruments (it is included in the IRIS run details). This makes it very hard and time consuming to gather all the information for a specific HOP.”

2-2. HOP Observations

HOP Proposer Log: http://hinode.msfc.nasa.gov/operations/hop_assessment/HOP_Proposer_Productivity_Log.html

HINODE OPERATION PLANS (HOPS)

Information needed for proposal development outlined below.
 Also refer to <http://www.isas.jaxa.jp/home/solar/guidance/index.html> as the lead reference.

After reviewing the material, ****SUBMIT A HOP PROPOSAL HERE****.

HOPs List & Guidance

Months/Starts HOP Guidance HOP List HOP List

HINODE Operation Plan (HOP) list

HOP No.	HOP Title
HOP-0012 web run 3 times	COSE VTT Hinode/SOT collaboration: Search of transient horizontal magnetic fields in the chromosphere
HOP-0013 web run 3 times	Relative structure of small magnetic elements: Faculae, network and minor networks in the quiet Sun
HOP-0014 web run 17 times TSD	COSE Two-Quasars Prominence Structure and Dynamics
HOP-0015 web run 3 times	Search for photospheric/chromospheric drivers of chromospheric penumbral microflares
HOP-0016 web run 3 times	Coordinated Observations between HinE Observatory & Hinode Satellite
HOP-0018 web run 2 times	Observation of an equatorial coronal hole
HOP-0019 web run 2 times	Support of SLM network Right
HOP-0020 web run 8 times TSD	The Study of Oribital Magnetic Connection in Umlauf Data
HOP-0021 web run 78 times	COSE (HOP040): Synoptic SOT Irradiance Scans
HOP-0022 web run 8 times	Polar Coronal Hole Observation

HOPs Coordination Calendar

Hinode HOP Coordination Calendar

POST-SUBMISSION PRODUCTIVITY UPDATES

SUBMIT HOP Results / Output

Hinode Operations Plan Productivity

Following productivity information used to better assess the quality of the HOP program and the value of future HOP submissions. Direct any questions or comments to Sabrina . Savage [at] nasa . gov.

* Required

HOP Number *

HOP Title *

HOP & Proposer Productivity Logs

HOP Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

137 Total Productivity Outputs [Histogram Per Year](#) [Histogram Per HOP](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

[Abramenko](#)
[Aime](#)

CONTENTS

Questions / Comments: [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov)

[HOP INSTRUCTION PAGE](#)

HOP Proposer Productivity Log

Log reliant on proposer-provided information and is therefore not complete. Please contact [sabrina.savage \[at\] nasa.gov](mailto:sabrina.savage@nasa.gov) to have any information updated.

- [Abramenko](#)
- [Aime](#)
- [Anan](#)
- [Anjali](#)
- [Antolin](#)
- [Anzer](#)
- [Arnaud](#)
- [Asai](#)
- [Asensio Ramos](#)
- [Attie](#)
- [Atrill](#)
- [Aulanier](#)
- [Bai](#)
- [Baker](#)
- [Balmaceda](#)
- [Balthasar](#)
- [Bando](#)
- [Banerjee](#)
- [Bastian](#)
- [Becher](#)
- [Beck](#)
- [Bellot Rubio](#)
- [Bemporad](#)
- [Berger](#)
- [Berghmans](#)
- [Bewsher](#)
- [Blanco](#)
- [Bogachev](#)
- [Brajša](#)
- [Brooks](#)
- [Brosius](#)
- [Cabello](#)
- [Cadavid](#)
- [Carlsson](#)
- [Carnonell](#)
- [Casini](#)
- [Cauzzi](#)
- [Centeno](#)
- [Centeno](#)



2-2. HOP Observations

SSC planning site: http://hinode.msfc.nasa.gov/submitted_hops.html

HINODE OPERATION PLANS SUBMISSIONS -- AUGUST 2015 SSC MEETING

RECENT/UPCOMING SUBMISSIONS

HOP 289 Filament/Su : Aug 1-8 : 17-21 UT
 HOP 287 GREGOR/Verma : Aug 10-19 : 8:30-10:30 UT
 HOP 292 Brightenings/Kanoh : Aug 10+ : 3 hours min
 HOP 286 NST/Hong : Aug 17-19 : 18-21 UT
 HOP 291 Hida/Ueno : Aug 17-23 : 0-4 UT
 HOP 173 EPO/Yaji : Aug 24-29 : 2-6 UT
 HOP 252 MOSES-II/Kankeiborg : **Aug 27 : 17:25-18:40 UT +/- (11:25 MDT)**
 HOP 288 Flares/Cheng : Aug 27-31 : 17-23 UT
 HOP 290 CLASP/Ishikawa : Sep 3** : 17:16-18:16 UT +/-
 HOP 257 SST/Tarbell : Sep 3-Oct 14 : 8-11 UT

NEW/UPDATED SUBMISSIONS

HOP206 -- PolarMap/Shimojo : September : Once every 3 days for 6 hours (SAA-free)
 Submitted HOP #1: Jets/Savcheva : Oct 19-25 : 6 hrs/day

Submitted HOP # 1:

Full Proposal

Title: Coordinated UV/EUV/X-Ray observations of coronal jets

Main Objective: Identify the structure and dynamics of the jet acceleration region and the mechanism for supplying plasma and acceleration

ToO: No

Proposer: Antonia Savcheva, Paola Testa, Katharine Reeves

Previous Submissions:

Savcheva: None

Testa

Reeves

Dates: 2015 Oct 19-25

Times: 6 hours per day, time of day not important

Target/Pointing: AR with some inclusion polarities on the outskirts, or near a coronal hole

SOT:

SP maps of the photospheric magnetic field

EIS:

A study similar to ISSI_jet_1, but add a few more cool lines (e.g., FeVIII, IX) with shorter exposure times, and a smaller raster size to overlap better with the IRIS FOV.

XRT:

384"x384" FOV, thin-Be filter, fast cadence (~20 sec) for as much of the observing period as telemetry allows.

Other instruments:

IRIS: medium 8-step dense raster, 4 sec exposure time, 1330 & 1400 SJs, large linelist (OBS ID: 3600257226).

Comments:

Ongoing HOP # 206/81:

HOP description

Title: Polar Panorama Map for understanding Polar Reversal in Cycle 24

ToO: No

Proposer: Shimojo, Tsuneta, Shiota, Sako, Anjali

Dates: September

Times: Once every 3 days for 6 hours

Target/Pointing: North Pole

Comments:

September is the month for the North pole, and we need to schedule runs of HOP 81 (6-days) and the Polar Panorama HOP 206 (every 2-3 days for a month). You may recall that he made the standing request at the SWG meeting this year. Both take 6-7 hours of SAA-free time per run.



2-2. HOP Observations

Hinode regularly coordinates with both ground- and space-based observatories and complements several regularly scheduled data-collecting observatories(*). Hinode also co-observes with sounding rocket and balloon technology development demonstrations. Much of the coordination is scheduled through the Hinode Operations Plan (HOP) program. Since 2008, partnering sites and instrumentation include (but not limited to):

Ground-based:

- Atacama Large Millimeter/Submillimeter Array (ALMA) – Chile
- Bialkow Observatory – Poland
- Big Bear Solar Observatory (BBSO) [NST/FISS/IRIM] – New Jersey
- Dunn Solar Telescope (DST/NSO) [IBIS/ROSA/SHAZAM/FIRS] – New Mexico
- Dutch Open Telescope (DOT) – La Palma
- Fuxian Lake Solar Observatory – China
- GREGOR Solar Telescope [GRIS] – Tenerife
- Haleakala Observatory – Hawaii
- Hida Observatory [DST] – Japan
- Iitate Radio Telescope (IPRT) – Tohoku University/Japan
- Kanzelhöhe Solar Observatory (KSO) – Austria
- Lomnický Peak Observatory [CoMP] – Czech Republic
- Mauna Loa Solar Observatory (MLSO) [CoMP] – Hawaii
- McMath-Pierce Telescope (NSO) – New Mexico
- Meudon Solar Tower – Paris
- Ondrejov Observatory – Czech Republic
- Pic du Midi Observatory – France
- Solar Magnetic Activity Research Telescope (SMART) – Japan
- Solar Terrestrial Laboratory [IPS] – Nagoya University/Japan
- Solar Tower Telescope of Nanjing University – China
- Swedish Solar Telescope (SST) [CRISP/TRIPPEL] – La Palma
- Synoptic Optical Long-term Investigations of the Sun (SOLIS/NSO) – New Mexico
- Vacuum Tower Telescope (VTT) – Tenerife
- Very Large Array (VLA)

(Note: Several High Schools and Science Museums in Japan)

Space-based:


- Active Cavity Radiometer Irradiance Monitor Satellite (ACRIM)
- *Advanced Composition Explorer (ACE)
- Akatsuki (Venus probe)
- Cassini (Saturn mission)
- Hubble Space Telescope (HST) [WFPC3]
- Interface Region Imaging Spectrograph (IRIS)
- Mercury Surface, Space Environment, Geochemistry, and Ranging (Mercury mission)
- Nuclear Spectroscopic Telescope Array (NuSTAR)
- Project for OnBoard Autonomy 2 (PROBA2) [SWAP]
- *Ramaty High Energy Solar Spectroscopic Imager (RHESSI)
- Solar and Heliospheric Observatory (SOHO) [SUMER/EIT/CDS/UVCS/MDI/LASCO]
- *Solar Dynamics Observatory (SDO) [AIA/EVE/HMI]
- Solar Radiation and Climate Experiment (SORCE) [TIM]
- *Solar Terrestrial Relations Observatory (STEREO) [EUVI]
- Telescopes for EUV Spectral Imaging of the Sun (TESIS)
- Time History of Events and Macroscale Interactions during Substorms (THEMIS)
- Transient Region and Coronal Explorer (TRACE)
- *Wind: Comprehensive Solar Wind Laboratory for Long-Term Solar Wind Measurements

Technology Demonstrations:

SUMI	HIC (1)	CLASP
RAISE	FOXSI	
SUNRISE (Balloon)	MOSES (1 & 2)	
EUNIS	VAULT	

2-3. Hinode Scientific Activities

Hinode: A Comprehensive Mission to Study the Variable Sun




A proposal to the Senior Review of Heliophysics Operating Missions
March 2015

Table 1: Recent Extended Mission Science results linked to relevant 2014 Heliophysics Roadmap Research Focus Areas (RFAs) and 2012 Decadal Survey Challenges (DSCs). *Additional example results are described in the text.*

RFAs & DSCs	§	SCIENCE OBJECTIVE	Example Results
§ 2.1: Understand the structure and stability of the magnetized atmosphere			
RFA H1	2.1.1	Observe the emergence of flux ropes	- Emerging flux shown to have connectivity in the corona through twist
	2.1.2	Study the evolution of polar crown filaments	- Transverse waves in a quiescent prominence imply chromospheric driving of high-frequency waves
	2.1.3	Map topological structures and the origin of the slow solar wind	- Slow solar wind sources strongly rooted in active region boundaries
DSCs SH 2 & 3			
§ 2.2: Measure the storage and release of mass and energy from the corona			
RFA F5	2.2.1	Determine the correlation between chromospheric and coronal heating	- Rapid brightenings in the chromosphere and transition region link to accelerated particles and heating
RFA F1	2.2.2	Characterize the properties of flare-forming regions and reconnection sites	- Reconnection likely produces plasma heating via turbulence
RFA F2	2.2.3	Isolate the locations of CME initiation	- Compact pre-eruption brightenings observed at the base of flaring active regions and above the erupting loop tops
RFA H1	2.2.4	Determine the role of magnetic field line braiding in atmospheric heating	- Transient brightening heating events favor stranded coronal loop models.
DSCs SH 1 & 2			
§ 2.3: Characterize energy and mass transfer from the photosphere to the corona			
RFA H1	2.3.1	Determine the role of the chromosphere in providing hot plasma to the corona	- Emerging flux quiescently reconnecting with pre-existing fields leads to total radiated energy comparable to M-class flares
RFA F5	2.3.2	Study the propagation of Alfvén waves through the solar atmosphere	- Reflected energy measurements indicate the generation of Alfvén wave turbulence in active region loops
DSCs SH 2 & 3			
§ 2.4: Quantify variations over the solar cycle			
RFA F4	2.4.1	Monitor dynamo and magnetic field changes	- Intermittent fields transfer flux to the magnetic network at a rate that could replace the entire network in 18-24 hours.
RFA H1	2.4.2	Survey irradiance variability	- The XRT solar irradiance time profile can provide quantitative correction to the <i>GOES</i> flux during low activity periods.
DSCs SH 2 & 3			

2-3. Hinode Scientific Activities

Hinode: A Comprehensive Mission to Study the Variable Sun





In response to the 2012 NRC Decadal Survey Science Challenges and 2014 Heliophysics Roadmap Research Focus Areas, the *Hinode* mission has set forth four **Prioritized Science Goals (PSGs)**:

- Study the sources and evolution of highly energetic dynamic events.
- Characterize cross-scale magnetic field topology and stability.
- Trace mass and energy flow from the photosphere to the corona.
- Continue long term synoptic support to quantify cycle variability.

Table 2: Prioritized Science Goals and required observations mapped to the 2014 Heliophysics Roadmap Research Focus Areas (RFAs) and 2012 Decadal Survey Challenges (DSCs).


RFAs & DSCs	§	SCIENCE OBJECTIVE	SAMPLE OBSERVATIONS
§ 3.1: PSG1 – Study the sources and evolution of highly energetic dynamic events			
RFA H1	3.1.1	Observe large-scale eruptive events from flare to particle acceleration.	- Coordinated radio, EUV, and X-ray observations of thermal & non-thermal eruption processes
RFA F2	3.1.2	Characterize the energetics of nanoflares and properties of non-thermal electrons.	- Nanoflare electron beam studies with <i>NuSTAR</i>
RFA F1	3.1.3	Probe magnetic reconnection flux transfer and energy release during flaring events.	- Spectrally probing chromospheric evaporation/condensation in flare loops
DSCs SH 2 & 3			Focused Mode opportunity: CME watch Synergies: <i>IRIS, ALMA VLA, EOVSA, NuSTAR, SDO, RHESSI, STEREO</i>
§ 3.2: PSG2 – Characterize the cross-scale magnetic field topology and stability.			
RFA H1	3.2.1	Study active region energy storage, topology, and evolution.	- Monitor the temperature stratification above ARs to potentially predict energy releases
RFA F2	3.2.2	Determine the impact of small-scale magnetic fields on the solar atmosphere.	- Specifically-designed sequences for <i>IRIS</i> coordination to observe unresolved fine structure in the transition region
DSCs SH 2 & 3	3.2.3	Determine the impact of large-scale magnetic field variations on the heliosphere.	- Test S-web model predictions concerning magnetic field topology in the outer corona with deep, wide-field SXR imaging
			Focused Mode opportunity: AR evolution Synergies: <i>IRIS, SDO</i>
§ 3.3: PSG3 – Trace mass and energy flow from the photosphere to the corona.			
RFA H1	3.3.1	Isolate solar wind sources and measure their mass supply.	- New EIS observing modes scanning for slow solar wind sources
RFA F5	3.3.2	Characterize the heating of the chromosphere.	- Link chromospheric heating diagnostic ion transport with Poynting flux estimates
DSCs SH 2 & 3	3.3.3	Quantify the generation, dissipation, and impact of magnetic waves.	- Vorticity measurements tracing chromospheric twist into the corona
			Focused Mode opportunity: EIS scans Synergies: <i>ACE, IRIS, SDO</i>
§ 3.4: PSG4 – Continue long term synoptic support to quantify cycle variability.			
RFA F4	3.4.1	Understand solar irradiance variations.	- Derive continuum contrast from SOT continuum bands and <i>IRIS</i> Mg II index
RFA H1	3.4.2	Monitor solar cycle evolution and stability.	- Polar magnetic fields, X-ray bright points, and magnetic activity band progression as indicators of solar cycle evolution and activity
DSCs SH 1 & 3	3.4.3	Relate solar variability to stellar evolution.	- Application of “Sun-as-a-star” methods to synoptic data for stellar abundance profiles
			Synergies: <i>IRIS, SDO</i>





2014 March 29 X-flare

EIS and *IRIS* spectral line profiles

Energy release at chromosphere sends cooler plasma down to surface; heated, 15 MK plasma rises into corona



2011 Sep 25 M-flare




Evaporation in solar flare footpoints observed by *Hinode*/EIS and *RHESSEI*. Upflow velocities in excess of 600 km/s were found for multi-million degree (~12 MK) lines of Fe XXIV and Fe XXIII. The data are compared with predictions of 1D hydrodynamic simulations. The arrows indicate footpoint regions (Doschek et al., to be submitted, 2014).

2-3. Hinode Scientific Activities


Coronal Non-Thermal Velocity in Polar Regions from Solar Minimum to Cycle 24 Solar Maximum

- The **non-thermal velocities** (Vnt) (e.g., turbulence, multiple flow sites) have been measured in the corona using the spectral line of Fe XII at 195.12 \AA observed by *Hinode*/EIS.
- The magnetic nulls at high latitude regions were determined. These show changes during the cycle induced by the opposite polarity streams reaching the poles.
- The coronal Vnt does not show such variation – this lack of variation during the cycle indicates that a **local dynamo may exist**.

Fe XII – 195.12 \AA



Magnetic Nulls






Harra et al., Solar Phys., submitted 2014



2-3. Hinode Scientific Activities

Hinode/EIS Coordinated Observations with the NRL VAULT Rocket on 2014 Sept 30


- What is the connection between the chromosphere, transition region, and corona? Coordinated observations lead us towards the answers.

EIS raster images in different spectral lines






“The Hinode satellite’s X-Ray Telescope recently observed the solar eclipse that crossed over North America on October 23rd. This rare event occurred as Hinode monitored solar flares from the largest active region seen on the Sun in over two decades.”



Hinode XRT – Thin-Be

C1.0 flare on Oct 10th shows clear signatures of supra-arcade downflowing loops (SADLs) in XRT and the hot AIA channels. SADLs are thought to be closely related to magnetic reconnection. (Savage et al. 2012)

SDO/AIA – 171, 193, 94, 131



2-3. Hinode Scientific Activities



ARTICLE

Received 24 Jun 2014 | Accepted 24 Nov 2014 | Published 6 Jan 2015

DOI: [10.1038/ncomms6947](https://doi.org/10.1038/ncomms6947)

Full-Sun observations for identifying the source of the slow solar wind


David H. Brooks^{1,†}, Ignacio Ugarte-Urra¹ & Harry P. Warren²

Goal: Identify sources of the flow of energy and matter throughout the solar system.

Solar wind has two components:

1. Fast: coronal holes
2. Slow: ??

Plasma composition used to distinguish the two components and link to solar structures.





EIS mosaics (2-day spectrometer scan during January 2013):

- Temperatures: $\sim 0.5 - 3$ MK

- Pure temperature images of the full Sun at the highest spatial resolution yet achieved



- Observation not normally practical due to time and telemetry

2-3. Hinode Scientific Activities



- Full Sun plasma composition map
 - *Required 16 million calculations to create!*
 - Darker → Photospheric abundances
 - **Lighter → Coronal abundances**

Slow solar wind trends toward coronal abundances.








- Doppler radial velocity map (2 MK)
 - Red → Inflow (toward Sun)
 - **Blue → Outflow (away from Sun)**



Magnetic field modeling of large-scale coronal field


- Subset of ~1.6 million lines shown
- Dotted → Closed
- **Solid → Open**





Slow solar wind source map


- Shear line due to solar rotation during 2-day scan
- Red/Green:
 - * Coronal composition
 - * Outflowing plasma
 - * Open field that extends down to ecliptic plane
- **Red** → **Active region (AR) boundaries**
- **Green** → **~Holes b/t ARs**
- [Other unknown source high in corona]



In situ measurements from ACE

- Stable slow solar wind
- EIS source map potentially accounts for **50-80%** of the flux

Note:




~90% of the solar wind flux originates below 40° latitude

Optimal candidate campaign for **Hinode Focused Mode operations**, except for significant tech. issue:

- **Reduced number of commands available**

Possible solution: Reduce number of pointings to encompass this 90% region.

Telemetry would still be a challenge.




One of the best sigmoidal active regions of the Hinode mission was observed on December 6/7, 2007. It is still providing a wealth of information about these kinds of structures today.

2-3. Hinode Scientific Activities




A. Savcheva et al., 2015a, in preparation


Magnetic field model



Doppler velocity




Nonthermal width



Non-linear force-free magnetic field (NLFFF) models are constructed from photospheric magnetograms and constrained by XRT data. When compared with EIS observations, the field model shows that the twisted core loops contain downflowing plasma, and provides evidence that large non-thermal widths are caused by additional heating due to reconnection that takes place at the edge of the flux rope.

2-3. Hinode Scientific Activities



A. Savcheva et al., 2015b, in preparation




The above figure shows maps of the squashing factor Q (left) and an overlay of $Q > 1E3$ on a STEREO EUVI 171 image (right) showing flare ribbons. High Q indicates highly divergent magnetic field structures.

Although the NLFFF models were optimized to match XRT observations, the Q maps derived from these models also correspond to observed flare ribbons in the EUV, indicating a connection between the magnetic topology and observed flare features.

The largest active regions usually appear in the declining phase of the Solar cycle, after the peak of the sunspot number has passed.





The sunspot group on 24 October 2014 (left) was the largest seen since 1990. It is larger than the Hinode SOT/BFI Field of View (220 x 110 arcseconds)



This region on 3 February 2015 has perhaps the largest area of any region this cycle. This vector magnetic map was made using the newly-perfected SOT/SP mosaic observing mode (600 x 160 arcsec)



2-3. Hinode Scientific Activities

Large flares continue to occur well past the sunspot maximum, providing many X- and M-flares for Hinode, RHESSI, and the rest of the HSO to observe. This X3.1 flare on October 24, 2014, was in AR 12192, which produced 6 X-flares (right) but no CMEs. The reason for this lack of eruptions may be the relatively low “magnetic free energy” in this region, despite its huge sunspots and magnetic flux.




2-3. Hinode Scientific Activities

After reversing their polarity around Solar Maximum, the polar magnetic fields strengthen (Shiota et al, 2015, in prep). Polar coronal holes begin to form, intermittently at first, and some traces of the next solar cycle begin to appear at high latitudes.





“Band-O-Gram,” showing the latitude migration of EUV bright points and the presence of two cycles on the Sun simultaneously. The new cycle points can be seen in high Northern latitudes starting in 2011. One of the Hinode 2015 Senior Review science goals is to repeat these analyses with a full cycle of XRT images. From McIntosh et al, 2014 ApJ, 792, 12.





2-3. Hinode Scientific Activities

THE INVERSE FIRST IONIZATION POTENTIAL (FIP) EFFECT IS A PROCESS WHICH RESULTS IN DIFFERENT ELEMENT ABUNDANCES IN THE SOLAR CORONA THAN IN THE PHOTOSPHERE.

- ◆ The inverse FIP effect has never been observed before in the Sun.
- ◆ The inverse FIP effect has just been observed in our star by the Extreme-ultraviolet Imaging Spectrometer (EIS) on *Hinode*
- ◆ **Argon XIV** is enhanced by a factor of about 30 over its normal coronal abundance in the example shown here (yes, it's the best example!).
- ◆ The inverse FIP enhancement is seven times larger than the photospheric abundance!
- ◆ Other enhancements in other regions are a factor of two or three greater than photospheric.
- ◆ The enhancements are seen close to sunspots. The precise association is not known.
- ◆ The solar coronal electron density is $\sim 4 \times 10^{10}$ up to about 10^{12} cm^{-3} .






Summary:

****** New study indicates inverse FIP effect observed during flare.

- Indicative of chromospheric evaporation in sunspot
- Observed in other stars but not before on Sun



2-3. Hinode Scientific Activities



Hinode captured a spectacular filament eruption on April 28, 2015.


XRT Al-mesh images show the prominence better than the Al-poly images because of the low temperature response of that filter.

Bright loops and streamers that are the consequences of reconnection in the wake of the eruption are prominently seen in XRT data, and can also be seen (though more faintly) in the AIA 193 data.




IRIS caught the filament liftoff, allowing for a detailed analysis of the dynamics of the eruption.

2-3. Hinode Scientific Activities






EIS scans in multiple wavelengths before and after the eruption show the changes in the corona caused by the disturbance. Post eruption loops are clearly seen in the Fe XV scan.

2-3. Hinode Scientific Activities



- Active Regions (AR) are regions where the sun's magnetic field is strong and hot and cool plasma is trapped.
- A filament (or filament channel, a.k.a. prominence on limb) is cool, thick plasma suspended above the surface where the magnetic field is changing. They appear shrouded in X-rays.
- A coronal cavity is a tube-like hole in the corona that typically sits above filament channels and are best viewed on the limb.
- Prominence arcade loops are the hot coronal loops that prevent prominences from erupting. When a prominence erupts, it has to tear these loops, which usually causes a coronal mass ejection with a solar flare.
- The south polar coronal hole is visible for this image. The poles are regions where weak magnetic fields exist of the same polarity. The fast solar wind originates in coronal holes.
- The filament sigmoid is a highly unpredictable filament that is likely to erupt. They are usually near or inside active regions are are closely monitored for space weather activity.



XRT

Soft X-ray: Corona
0.5 – 20 Million degrees

Kanzelhöhe Solar Observatory
(ESA – Ground-based)

H-alpha: Chromosphere
~10,000 degrees

Using the transit of Venus to probe the upper planetary atmosphere


Nature Communications, June 2015, DOI: 10.1038/ncomms8563

<http://www.nasa.gov/feature/goddard/scientists-study-venus-atmosphere-through-transit>

<http://www.csmonitor.com/Science/2015/0623/Venus-Does-this-sunlight-make-me-look-fat>

<http://www.natureasia.com/en/research/highlight/9988/>

Venus transit images from *Hinode* and *SDO* show that the planet's atmosphere is more extended than previously thought, implications for exoplanetary transit measurements with different wavelengths and for planetary probes encountering and/or using atmospheric drag.



“Seering Sun seen in X-rays”

<https://www.ras.org.uk/news-and-press/2676-searing-sun-seen-in-x-rays>


<http://www.nasa.gov/jpl/nustar/searing-sun-seen-in-x-rays>

Blue = *NuSTAR* (highest energy)

Green = *Hinode*/XRT

Red/Yellow = *SDO*/AIA

Combined data used to search for nanoflare heating of the corona.



Hinode/SOT/Ca II

IRIS Si IV Mg II k

Lower part


long threads (SOT)
small oscillations (SOT)
long lifetimes (SOT)

Upper part

short threads (SOT)
large oscillations (SOT)
short lifetimes (SOT)
higher V_{Los} (IRIS)
broad line widths (IRIS)



Evidence for heating by waves in the upper part




SOT observation:
Ca images, 8 sec cadence

IRIS observation:
4-step raster & slit jaw
images, 20 sec cadence


Joten Okamoto+15 coauthors, submitted to ApJ

2-3. Hinode Scientific Activities



Hinode SOT Ca II



IRIS Mg II k



Histogram of v_{LOS} and line width with height





Upper region:

- higher v_{LOS}
- large non-thermal comp.


40 km/s
= thermal velocity of 250,000 K

SOT Ca II (10,000 K)



IRIS Si IV (80,000 K)

x-t plot




time


horizontal direction

2-3. Hinode Scientific Activities

Hinode and IRIS observations




Numerical simulation



Resonant Absorption triggers azimuthal flows on the surface of flux tube

(e.g., Ionson 1978, Sakurai+1991, Goossens+2002)



This mechanism can explain the observed features:

- in-phase velocity patterns at 2 slit locations (green and red)
- changing velocity sign with time
- 180-degree difference between POS displacement and LOS velocity
- Temperature transition

3-1, 3-2. Focused Mode Coordination

- Status: Operable
 - No major issues reported from the teams.
 - (*Report from SOT upcoming.*)

- Request for FM HOPs sent to community via SolarNews.


- Focused Mode coordination
 - Priority list circulating with weekly meetings
 - Active Region evolution (flux emergence, waves in sunspots, flare monitoring)
 - Coronal Holes
 - Prominence / Filament
 - Disk-center (long baseline synoptic scans)
 - Polar magnetic network
 - Focused Mode Liaison no longer needed?

- Status:
 - SOT report:
 - Pure FM week easier and quicker than a normal week
 - Transition week is a little harder
 - Harder to stay on best AR target
 - Long continuous observations are good, but requires a decent target during planning
 - Not clear if it saves funds
 - Hours spent for mission ops is about the same in FM and normal mode (pourquoi?) and slightly higher in transitions--differences are not statistically significant; more significant effects are who is CO and if the shift shared by 2 people.
 - CO comments
 - It takes longer to do a 7 day plan but not 3 times longer. Health monitoring, data checking, coordination, reports take the same amount of time.
 - Yugma and pointing tool problems were highlighted with one planning opportunity per week

- HOP 79 could not be run in June due to a combination of focused mode and major flare watch restrictions
- HOP 285 (with VTT and IRIS) was limited due to availability of tracking curves in the 7-day plan
 - *This should be expected by or explained to the HOP proposers during FM.*
- Took longer to assemble a full set of SP/FG flat fields prior to and during eclipse season
- A pointing table upload had to be made during one of the 7-day plans, changing the SOT center pointing from a flare site between 2 emerging active regions to the leading spot of the westernmost region when activity returned to a lower level.
 - *Did this go smoothly?*

3-1, 3-2. Focused Mode Coordination

- Flare WatchDog
 - Yumi Bamba-san has been filling this position very well while working on her thesis
 - Suggestions for successor? Continue with a graduate student?
 - Does the procedure need to be sent around to the COs or just the FWD?





3-3. Focused Mode Calendar

2015

January						
S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

April						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

July						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

October						
S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

February						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28

May						
S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

August						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

November						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

March						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

June						
S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

September						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			

December						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

Focused Mode
(Tuesday upload only)

Normal
(Tuesday, Thursday, Saturday uploads)

IRIS Coordination

Hinode Eclipse

IRIS Eclipse

Rocket Launch / Notable Campaign

~ HOP 130

November 24th is Thanksgiving week (less US support already) and well into IRIS eclipse season. FM during Dec/Jan holidays.

Before looking ahead to 2016....

Fractional reduction in timeline creation (R):

$$R = \frac{N_plans_{Norm} - N_plans_{Norm+FM}}{N_plans_{Norm}}$$

$$N_plans_{Norm} \approx (52 \text{ weeks}) * (3 \frac{\text{plans}}{\text{week}}) = 156 \text{ plans}$$

$$N_plans_{Norm+FM} \approx N_plans_{Norm} - (N_weeks_{FM} * 2)$$

$$R \approx \frac{N_weeks_{FM} * 2}{156}$$

18 weeks of FM (2015) → 35% of the year; 23% reduction in number of plans generated
15 weeks of FM → 28% of the year; 19% reduction

Keep FM frequency as proposed or reduce by one 3 week run until the next review cycle (2018)? [i.e., phasing]



3-3. Focused Mode Calendar

PROPOSED PROPOSED PROPOSED 2016 PROPOSED PROPOSED PROPOSED

January						
S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

April						
S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30

July						
S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

October						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

February						
S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29					

May						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

August						
S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

November						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			

March						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

June						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

September						
S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	

December						
S	M	T	W	T	F	S
					1	2
					3	
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

Focused Mode
(Tuesday upload only)

Normal
(Tuesday, Thursday, Saturday uploads)

IRIS Coordination

Hinode Eclipse

IRIS Eclipse

Rocket Launch / Notable Campaign

~ HOP 130

*** Hi-C II launch in June/July?? **

November 22ND is Thanksgiving week (less US support already) and well into IRIS eclipse season (to be confirmed). FM during Dec/Jan holidays.

Hinode Overall Assessment and Findings

The 2015 Senior Review panel finds that the scientific return in recent years from *Hinode* is exceptional. The mission returns data in a unique niche of HSO data space, and the mission is in fair overall health with adequate data archiving efforts. Continuing the *Hinode* mission would ensure that a unique HSO asset is supported. The panel finds the proposal to demonstrate an excellent outlook for future collaborative work with other HSO missions (in particular, with IRIS, RHESSI, and SDO). The proposal meets the in-guide budget, and the team has developed new operating modes to be able to operate in an efficient manner while continuing to enable broad scientific discovery.

The Panel notes that the proposal includes approximately \$2.6M/year for Science Data Analysis, which in our opinion should include funding both for routine data quality analysis and to perform relevant scientific investigations by the team. The Panel was surprised upon comments from the presenters that for the preceding extended mission phase, no mission-related science was performed under extended-phase funding. The Panel is concerned that this approach is the result of a misinterpretation of the 2013 Senior Review findings, and wishes here to clarify that there is no restriction on using part of that SDA funding to support *Hinode*-related science research by the team. This appears to be common practice across the other mission teams, with the caveat that this has to be balanced with the priority to meet data production commitments under a constrained overall budget which is likely to reduce with time.

5. NASA situation



The panel found that the major concerns raised in 2010 and 2013 SR cycles have been addressed: costs have been reduced, the instrument teams have developed new, efficient observing modes to further streamline operations while ensuring that high-quality data are obtained. There is still some minor concern regarding a high cost/science ratio in light of the contributions from international partners. The proposed science available to additional *Hinode* operations is compelling, and there is recognition that this is an extremely complex mission. The panel praises the *Hinode* team's proactive engagement in cross-mission collaborations. The scientific output has been very satisfactory, especially in the context of addressing fundamental outstanding questions and testing model predictions, with generally quantitative methods and results.

The *Hinode* extended mission proposal received an 8/10 median score for extended mission science, and 9/10 median score for contributions to Heliophysics Systems Observatory; both scores are in the highest, "compelling" category. The panel finds that continuing the *Hinode* mission would ensure continued systems-approach research to outstanding topics in solar and heliospheric physics.

Hinode Proposal Weaknesses

While the scientific achievements from *Hinode* data were compelling and fully exploited the possibility of “system” science, the panel found the presentation of the science to be somewhat fragmented. The proposal adequately describes the links between PSGs, the 2014 Heliophysics Roadmap Research Focus Areas and 2012 Decadal Survey Challenges, but the narrative could have better mirrored the “system science” approach to bring continuity between at times seemingly disparate investigation topics.

3. Mission Grades

	Mission Median	Science Std Dev		HSO Median	Science Std Dev
IBEX	9	1.6	ACE	9	1.9
IRIS	9	1.3	SDO	9	1.5
Van Allen	9	1.5	Hinode	9	1.8
AIM	8	1.2	IBEX	8	1.2
Hinode	8	1.2	IRIS	8	1.2
RHESSI	8	1.0	STEREO (2)	8	1.1
STEREO (2)	8	1.2	THEMIS	8	1.3
TIMED	8	1.2	Wind	8	1.3
Voyager	8	1.6	RHESSI	8	1.6
ACE	7	1.5	STEREO (1)	8	1.5
SDO	7	1.5	Voyager	7	1.6
STEREO (1)	7	1.4	AIM	7	1.3
THEMIS	7	1.8	TIMED	7	1.9
TWINS	7	1.3	TWINS	7	1.3
Wind	7	1.2	Van Allen	7	1.4
CINDI	7	1.2	CINDI	6	1.2

The mission grades are as follows: Grades (scale of 0 to 10, 10 is best)

10–8 Future contributions promise to be compelling

7–4 Excellent, but less compelling

3–0 Future contributions relatively modest



-