



Use of GCMs in constraining climate sensitivity

Ringberg
March 2015


Gavin Schmidt, Kate Marvel, Ron
Miller and Larissa Nazarenko


NASA GISS



... and climate response...

Goddard Institute for Space Studies





Goddard Institute for
Space Studies

GISS-E2 Climate Sensitivities

GISS-E2 (Qflux) (NINT): 2.7°C (**Charney**)


GISS-E2 (Qflux) (TCADI): 2.9°C

GISS-E2-R (NINT): TCR=1.4°C ; ECS=2.3°C

Effective CS (Gregory): 2.1°C

GISS-E2-R (TCADI): TCR=1.6°C ; ECS=2.4°C

Effective CS (Gregory): 2.4°C



Goddard Institute for
Space Studies

Historical forcings by concentration

WMGHG: CO₂, CFCs, HFC, N₂O

CH₄ (incl. stratospheric
water vapor)

Ozone: tropospheric and
stratospheric


Anthropogenic Aerosols: SO₄, NO₃,
BC, OC

Indirect Effect: Tuned (~ -1 W/m²)

Land Use/Land over: Crops/Pasture

Solar (incl spectral irradiance)

Volcanic: Aerosol Optical Depth



Miller et al, 2014


All forcings also used singly in 5 member ensemble




Goddard Institute for Space Studies

Do forcings add linearly?

By concentration



By emissions







Irrigation as an historical forcing

Goddard Institute for Space Studies

Annual Irrigation Rates (mm day^{-1})




Annual Irrigation, km^3



Irrigation applied based on historical data.
Large-scale cooling (More LH, less SH)

Cook et al, 2014




Goddard Institute for
Space Studies

What is the role of GCMs?

(Collectively) as 'expert prior' (not pdf)

Estimates of forcings





Estimates of mapping between observables, derived parameters and actual emergent properties.





Single forcing runs CMIP5 GISS-E2-R

Goddard Institute for Space Studies






Perfect model framework


Goddard Institute for Space Studies

Known OHC, SAT, Forcing => defn. $\lambda_{TCR/ECS}$ from

$$\Delta F = \lambda_{TCR(ECS)} \Delta T (+ \Delta Q)$$

$$\Rightarrow TCR = \Delta F_{2xCO2} / \lambda_{TCR} \quad ; \quad ECS = \Delta F_{2xCO2} / \lambda_{ECS}$$





Goddard Institute for
Space Studies

Summary of inference from single forcing runs (GISS-E2-R)

Forcing	TCR (°C)	ECS (°C)
GHG	1.5	2.2
AA	2.0	3.4
LU	5.4	3.8
Oz	0.8	1.1
Historical	1.2	1.6
Actual	1.4	2.3


Forcings add vectorially in $\Delta F/\Delta T$ space




Goddard Institute for Space Studies

PDFs from historical record adjusted by single-forcing results

TCR




ECS



***Median, 17-83% range**

Uncertainties in E derived from ensemble spread



Goddard Institute for
Space Studies

Conclusions

TCR/ECS constraints based on historical transients are biased low.

Results from GISS ModelE suggest that this is related to higher efficacy of aerosols and LU

Accounting for this largely reconciles historical period estimates with paleo spread.

Even if a single model not definitive, not basis to assume $E_i=0$ with no uncertainty.

Increase in median TCR/ECS \sim 35-60%